

Package ‘BSDA’

July 21, 2025

Type Package

Title Basic Statistics and Data Analysis

Version 1.2.2

Date 2023-09-14

LazyData yes

Maintainer Alan T. Arnholt <arnholtat@appstate.edu>

Description Data sets for book ``Basic Statistics and Data Analysis" by
Larry J. Kitchens.

Depends lattice, R (>= 2.10)

Imports e1071

License GPL-3

Suggests ggplot2 (>= 2.1.0), dplyr, tidyr

RoxygenNote 7.2.3

Encoding UTF-8

URL <https://github.com/alanarnholt/BSDA>,
<https://alanarnholt.github.io/BSDA/>

BugReports <https://github.com/alanarnholt/BSDA/issues>

NeedsCompilation no

Author Alan T. Arnholt [aut, cre],
Ben Evans [aut]

Repository CRAN

Date/Publication 2023-09-18 17:50:05 UTC

Contents

Abbey	9
Abc	10
Abilene	10
Ability	11

Abortion	12
Absent	13
Achieve	14
Adsales	14
Aggress	15
Aid	16
Aids	17
Airdisasters	18
Airline	19
Alcohol	20
Allergy	20
Anesthet	21
Anxiety	22
Apolipop	22
Append	23
Appendec	24
Aptitude	25
Archaeo	25
Arthriti	26
Artifici	27
Asprin	28
Asthmati	28
Attorney	29
Autogear	30
Backtoback	31
Bbsalaries	31
Bigten	32
Biology	33
Birth	33
Blackedu	34
Blood	35
Board	36
Bones	37
Books	37
Bookstor	38
Brain	39
Bumpers	40
Bus	41
Bypass	41
Cabinets	42
Cancer	43
Carbon	44
Cat	45
Censored	45
Challeng	46
Chemist	47
Chesapea	48
Chevy	48

Chicken	49
Chipavg	50
Chips	51
Cigar	52
Cigarett	52
CIsim	53
Citrus	55
Clean	55
Coaxial	56
Coffee	57
Coins	57
Combinations	58
Commute	59
Concept	60
Concrete	60
Corn	61
Correlat	62
Counsel	62
Cpi	63
Crime	64
Darwin	65
Dealers	66
Defectiv	66
Degree	67
Delay	68
Depend	69
Detroit	69
Develop	70
Devmath	71
Dice	71
Diesel	72
Diplomat	73
Disposal	74
Dogs	75
Domestic	76
Dopamine	77
Dowjones	77
Drink	78
Drug	79
Dyslexia	80
Earthqk	81
EDA	81
Educat	82
Eggs	83
Elderly	84
Energy	85
Engineer	86
Entrance	86

Epaminicompact	87
Epatwoseater	88
Executiv	89
Exercise	90
Fabric	90
Faithful	91
Family	92
Ferraro1	93
Ferraro2	93
Fertility	94
Firstchi	95
Fish	96
Fitness	97
Florida2000	98
Fluid	99
Food	100
Framingh	100
Freshman	101
Funeral	102
Galaxie	103
Gallup	103
Gasoline	104
German	105
Golf	106
Governor	107
Gpa	108
Grades	109
Graduate	109
Greenriv	110
Gmnriv2	111
Groupabc	111
Groups	112
Gym	113
Habits	113
Haptoglo	114
Hardware	115
Hardwood	116
Heat	117
Heating	118
Hodgkin	118
Homes	119
Homework	120
Honda	121
Hostile	122
Housing	122
Hurrican	123
Iceberg	124
Income	125

Independent	126
Indian	127
Indiapol	128
Indy500	128
Inflatio	129
Inletoil	130
Inmate	131
Inspect	132
Insulate	133
Iqgpa	134
Irises	134
Jdpower	135
Jobsat	136
Kidsmoke	137
Kilowatt	138
Kinder	138
Laminect	139
Lead	140
Leader	141
Lethal	141
Life	142
Lifespan	143
Ligntmonth	143
Lodge	144
Longtail	145
Lowabil	146
Magnesium	146
Malpract	147
Manager	148
Marked	148
Math	149
Mathcomp	150
Mathpro	151
Maze	152
Median	152
Mental	153
Mercury	154
Metrent	154
Miller	155
Miller1	156
Moisture	156
Monoxide	157
Movie	158
Music	159
Name	160
Nascar	161
Nervous	161
Newsstand	162

Nfldraf2	163
Nfldraft	163
Nicotine	164
normarea	165
nsiz	166
ntester	167
Orange	168
Orioles	168
Oxytocin	169
Parented	170
Patrol	171
Pearson	172
Phone	172
Poison	173
Politic	174
Pollutio	175
Porosity	175
Poverty	176
Precinct	177
Prejudic	178
Presiden	178
Press	179
Prognost	180
Program	181
Psat	181
Psych	182
Puerto	183
Quail	183
Quality	184
Rains	185
Randd	186
Rat	186
Ratings	187
Reaction	188
Reading	189
Readiq	189
Referend	190
Region	191
Register	192
Rehab	192
Remedial	193
Rentals	194
Repair	195
Retail	195
Ronbrown1	196
Ronbrown2	197
Rural	197
Salary	198

Salinity	199
Sat	199
Saving	200
Scales	201
Schizop2	202
Schizoph	203
Seatbelt	203
Selfdefe	204
Senior	205
Sentence	206
Shkdrug	206
Shock	207
Shoplift	208
Short	209
Shuttle	209
SIGN.test	210
Simpson	213
Situp	214
Skewed	214
Skin	215
Slc	216
Smokyph	216
Snore	217
Snow	218
Soccer	219
Social	220
Sophomor	220
South	221
Speed	222
Spellers	222
Spelling	223
Sports	224
Spouse	224
SRS	225
Stable	226
Stamp	226
Statclas	227
Statelaw	228
Statisti	229
Step	229
Stress	230
Study	231
Submarin	232
Subway	233
Sunspot	233
Superbowl	234
Supercar	235
Tablrock	236

Teacher	237
Tenness	238
Tensile	239
Test1	240
Thermal	240
Tiaa	241
Ticket	242
Toaster	242
Tonsils	243
Tort	244
Toxic	245
Track	246
Track15	247
Treatments	247
Trees	248
Trucks	249
tsum.test	249
Tv	253
Twin	253
Undergrad	254
Vacation	255
Vaccine	256
Vehicle	257
Verbal	257
Victoria	258
Viscosit	259
Visual	260
Vocab	260
Wastewat	261
Weather94	262
Wheat	263
Windmill	264
Window	264
Wins	265
Wool	266
Yearsunspot	267
z.test	267
zsum.test	270

Abbey	<i>Daily price returns (in pence) of Abbey National shares between 7/31/91 and 10/8/91</i>
-------	--

Description

Data used in problem 6.39

Usage

Abbey

Format

A data frame/tibble with 50 observations on one variable

price daily price returns (in pence) of Abbey National shares

Source

Buckle, D. (1995), Bayesian Inference for Stable Distributions, *Journal of the American Statistical Association*, 90, 605-613.

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
qqnorm(Abbey$price)
qqline(Abbey$price)
t.test(Abbey$price, mu = 300)
hist(Abbey$price, main = "Exercise 6.39",
     xlab = "daily price returns (in pence)",
     col = "blue")
```

Abc	<i>Three samples to illustrate analysis of variance</i>
-----	---

Description

Data used in Exercise 10.1

Usage

Abc

Format

A data frame/tibble with 54 observations on two variables

response a numeric vector

group a character vector A, B, and C

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
boxplot(response ~ group, col=c("red", "blue", "green"), data = Abc )
anova(lm(response ~ group, data = Abc))
```

Abilene	<i>Crimes reported in Abilene, Texas</i>
---------	--

Description

Data used in Exercise 1.23 and 2.79

Usage

Abilene

Format

A data frame/tibble with 16 observations on three variables

crimetype a character variable with values Aggravated assault, Arson, Burglary, Forcible rape, Larceny theft, Murder, Robbery, and Vehicle theft.

year a factor with levels 1992 and 1999

number number of reported crimes

Source

Uniform Crime Reports, US Dept. of Justice.

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
par(mfrow = c(2, 1))
barplot(Abilene$number[Abilene$year=="1992"],
names.arg = Abilene$crimetype[Abilene$year == "1992"],
main = "1992 Crime Stats", col = "red")
barplot(Abilene$number[Abilene$year=="1999"],
names.arg = Abilene$crimetype[Abilene$year == "1999"],
main = "1999 Crime Stats", col = "blue")
par(mfrow = c(1, 1))

## Not run:
library(ggplot2)
ggplot2::ggplot(data = Abilene, aes(x = crimetype, y = number, fill = year)) +
  geom_bar(stat = "identity", position = "dodge") +
  theme_bw() +
  theme(axis.text.x = element_text(angle = 30, hjust = 1))

## End(Not run)
```

Ability

Perceived math ability for 13-year olds by gender

Description

Data used in Exercise 8.57

Usage

Ability

Format

A data frame/tibble with 400 observations on two variables

gender a factor with levels girls and boys

ability a factor with levels hopeless, belowavg, average, aboveavg, and superior

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
CT <- xtabs(~gender + ability, data = Ability)
CT
chisq.test(CT)
```

Abortion

Abortion rate by region of country

Description

Data used in Exercise 8.51

Usage

Abortion

Format

A data frame/tibble with 51 observations on the following 10 variables:

state a character variable with values alabama, alaska, arizona, arkansas, california, colorado, connecticut, delaware, dist of columbia, florida, georgia, hawaii, idaho, illinois, indiana, iowa, kansas, kentucky, louisiana, maine, maryland, massachusetts, michigan, minnesota, mississippi, missouri, montana, nebraska, nevada, new hampshire, new jersey, new mexico, new york, north carolina, north dakota, ohio, oklahoma, oregon, pennsylvania, rhode island, south carolina, south dakota, tennessee, texas, utah, vermont, virginia, washington, west virginia, wisconsin, and wyoming

region a character variable with values midwest northeast south west

regcode a numeric vector

rate1988 a numeric vector

rate1992 a numeric vector

rate1996 a numeric vector

provide1988 a numeric vector

provide1992 a numeric vector

lowhigh a numeric vector

rate a factor with levels Low and High

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
T1 <- xtabs(~region + rate, data = Abortion)
T1
chisq.test(T1)
```

Absent

Number of absent days for 20 employees

Description

Data used in Exercise 1.28

Usage

Absent

Format

A data frame/tibble with 20 observations on one variable

days days absent

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
CT <- xtabs(~ days, data = Absent)
CT
barplot(CT, col = "pink", main = "Exercise 1.28")
plot(ecdf(Absent$days), main = "ECDF")
```

Achieve	<i>Math achievement test scores by gender for 25 high school students</i>
---------	---

Description

Data used in Example 7.14 and Exercise 10.7

Usage

Achieve

Format

A data frame/tibble with 25 observations on two variables

score mathematics achievement score

gender a factor with 2 levels boys and girls

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
anova(lm(score ~ gender, data = Achieve))
t.test(score ~ gender, var.equal = TRUE, data = Achieve)
```

Adsales	<i>Number of ads versus number of sales for a retailer of satellite dishes</i>
---------	--

Description

Data used in Exercise 9.15

Usage

Adsales

Format

A data frame/tibble with six observations on three variables

month a character vector listing month

ads a numeric vector containing number of ads

sales a numeric vector containing number of sales

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
plot(sales ~ ads, data = Adsales, main = "Exercise 9.15")
mod <- lm(sales ~ ads, data = Adsales)
abline(mod, col = "red")
summary(mod)
predict(mod, newdata = data.frame(ads = 6), interval = "conf", level = 0.99)
```

Aggress	<i>Aggressive tendency scores for a group of teenage members of a street gang</i>
---------	---

Description

Data used in Exercises 1.66 and 1.81

Usage

Aggress

Format

A data frame/tibble with 28 observations on one variable

aggres measure of aggressive tendency, ranging from 10-50

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
with(data = Aggress,
      EDA(aggres))
# OR
IQR(Aggress$aggres)
diff(range(Aggress$aggres))
```

Aid	<i>Monthly payments per person for families in the AFDC federal program</i>
-----	---

Description

Data used in Exercises 1.91 and 3.68

Usage

Aid

Format

A data frame/tibble with 51 observations on two variables

state a factor with levels Alabama, Alaska, Arizona, Arkansas, California, Colorado, Connecticut, Delaware, District of Columbia, Florida, Georgia, Hawaii, Idaho, Illinois, Indiana, Iowa, Kansas, Kentucky, Louisiana, Maine, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Montana, Nebraska, Nevada, New Hampshire, New Jersey, New Mexico, New York, North Carolina, North Dakota, Ohio, Oklahoma, Oregon, Pennsylvania, Rhode Island, South Carolina, South Dakota, Tennessee, Texas, Utah, Vermont, Virginia, Washington, West Virginia, Wisconsin, and Wyoming

payment average monthly payment per person in a family

Source

US Department of Health and Human Services, 1993.

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
hist(Aid$payment, xlab = "payment", main =
"Average monthly payment per person in a family",
col = "lightblue")
boxplot(Aid$payment, col = "lightblue")
dotplot(state ~ payment, data = Aid)
```

Aids	<i>Incubation times for 295 patients thought to be infected with HIV by a blood transfusion</i>
------	---

Description

Data used in Exercise 6.60

Usage

Aids

Format

A data frame/tibble with 295 observations on three variables

duration time (in months) from HIV infection to the clinical manifestation of full-blown AIDS

age age (in years) of patient

group a numeric vector

Source

Kalbsleich, J. and Lawless, J., (1989), An analysis of the data on transfusion related AIDS, *Journal of the American Statistical Association*, 84, 360-372.

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
with(data = Aids,
  EDA(duration)
)
with(data = Aids,
  t.test(duration, mu = 30, alternative = "greater")
)
with(data = Aids,
  SIGN.test(duration, md = 24, alternative = "greater")
)
```

Airdisasters

Aircraft disasters in five different decades

Description

Data used in Exercise 1.12

Usage

Airdisasters

Format

A data frame /tibble with 141 observations on the following seven variables

year a numeric vector indicating the year of an aircraft accident

deaths a numeric vector indicating the number of deaths of an aircraft accident

decade a character vector indicating the decade of an aircraft accident

Source

2000 *World Almanac and Book of Facts*.

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
par(las = 1)
stripchart(deaths ~ decade, data = Airdisasters,
           subset = decade != "1930s" & decade != "1940s",
           method = "stack", pch = 19, cex = 0.5, col = "red",
           main = "Aircraft Disasters 1950 - 1990",
           xlab = "Number of fatalities")
par(las = 0)
```

Airline	<i>Percentage of on-time arrivals and number of complaints for 11 airlines</i>
---------	--

Description

Data for Example 2.9

Usage

Airline

Format

A data frame/tibble with 11 observations on three variables

airline a character variable with values Alaska, Amer West, American, Continental, Delta, Northwest, Pan Am, Southwest, TWA, United, and USAir

ontime a numeric vector

complaints complaints per 1000 passengers

Source

Transportation Department.

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
with(data = Airline,  
      barplot(complaints, names.arg = airline, col = "lightblue",  
              las = 2)  
)  
plot(complaints ~ ontime, data = Airline, pch = 19, col = "red",  
      xlab = "On time", ylab = "Complaints")
```

Alcohol

Ages at which 14 female alcoholics began drinking

Description

Data used in Exercise 5.79

Usage

Alcohol

Format

A data frame/tibble with 14 observations on one variable

age age when individual started drinking

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
qqnorm(Alcohol$age)
qqline(Alcohol$age)
SIGN.test(Alcohol$age, md = 20, conf.level = 0.99)
```

Allergy

Allergy medicines by adverse events

Description

Data used in Exercise 8.22

Usage

Allergy

Format

A data frame/tibble with 406 observations on two variables

event a factor with levels insomnia, headache, and drowsiness

medication a factor with levels seldane-d, pseudoephedrine, and placebo

Source

Marion Merrel Dow, Inc. Kansas City, Mo. 64114.

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
T1 <- xtabs(~event + medication, data = Allergy)
T1
chisq.test(T1)
```

Anesthet

Recovery times for anesthetized patients

Description

Data used in Exercise 5.58

Usage

Anesthet

Format

A with 10 observations on one variable

recover recovery time (in hours)

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
qqnorm(Anesthet$recover)
qqline(Anesthet$recover)
with(data = Anesthet,
  t.test(recover, conf.level = 0.90)$conf
)
```

Anxiety

Math test scores versus anxiety scores before the test

Description

Data used in Exercise 2.96

Usage

Anxiety

Format

A data frame/tibble with 20 observations on two variables

anxiety anxiety score before a major math test

math math test score

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
plot(math ~ anxiety, data = Anxiety, ylab = "score",
     main = "Exercise 2.96")
with(data = Anxiety,
     cor(math, anxiety)
)
linmod <- lm(math ~ anxiety, data = Anxiety)
abline(linmod, col = "purple")
summary(linmod)
```

Apolipop

Level of apolipoprotein B and number of cups of coffee consumed per day for 15 adult males

Description

Data used in Examples 9.2 and 9.9

Usage

Apolipop

Format

A data frame/tibble with 15 observations on two variables

coffee number of cups of coffee per day

apolipB level of apolipoprotein B

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
plot(apolipB ~ coffee, data = Apolipop)
linmod <- lm(apolipB ~ coffee, data = Apolipop)
summary(linmod)
summary(linmod)$sigma
anova(linmod)
anova(linmod)[2, 3]^0.5
par(mfrow = c(2, 2))
plot(linmod)
par(mfrow = c(1, 1))
```

Append

Median costs of an appendectomy at 20 hospitals in North Carolina

Description

Data for Exercise 1.119

Usage

Append

Format

A data frame/tibble with 20 observations on one variable

fee fees for an appendectomy for a random sample of 20 hospitals in North Carolina

Source

North Carolina Medical Database Commission, August 1994.

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```

fee <- Appendec$fee
ll <- mean(fee) - 2*sd(fee)
ul <- mean(fee) + 2*sd(fee)
limits <- c(ll, ul)
limits
fee[fee < ll | fee > ul]

```

Appendec	<i>Median costs of appendectomies at three different types of North Carolina hospitals</i>
----------	--

Description

Data for Exercise 10.60

Usage

Appendec

Format

A data frame/tibble with 59 observations on two variables

cost median costs of appendectomies at hospitals across the state of North Carolina in 1992

region a vector classifying each hospital as rural, regional, or metropolitan

Source

Consumer's Guide to Hospitalization Charges in North Carolina Hospitals (August 1994), North Carolina Medical Database Commission, Department of Insurance.

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```

boxplot(cost ~ region, data = Appendec, col = c("red", "blue", "cyan"))
anova(lm(cost ~ region, data = Appendec))

```

Aptitude

Aptitude test scores versus productivity in a factory

Description

Data for Exercises 2.1, 2.26, 2.35 and 2.51

Usage

Aptitude

Format

A data frame/tibble with 8 observations on two variables

aptitude aptitude test scores

product productivity scores

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
plot(product ~ aptitude, data = Aptitude, main = "Exercise 2.1")
model1 <- lm(product ~ aptitude, data = Aptitude)
model1
abline(model1, col = "red", lwd=3)
resid(model1)
fitted(model1)
cor(Aptitude$product, Aptitude$aptitude)
```

Archaeo

Radiocarbon ages of observations taken from an archaeological site

Description

Data for Exercises 5.120, 10.20 and Example 1.16

Usage

Archaeo

Format

A data frame/tibble with 60 observations on two variables

age number of years before 1983 - the year the data were obtained

phase Ceramic Phase numbers

Source

Cunliffe, B. (1984) and Naylor and Smith (1988).

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
boxplot(age ~ phase, data = Archaeo, col = "yellow",
        main = "Example 1.16", xlab = "Ceramic Phase", ylab = "Age")
anova(lm(age ~ as.factor(phase), data= Archaeo))
```

Arthriti

Time of relief for three treatments of arthritis

Description

Data for Exercise 10.58

Usage

Arthriti

Format

A data frame/tibble with 51 observations on two variables

time time (measured in days) until an arthritis sufferer experienced relief

treatment a factor with levels A, B, and C

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
boxplot(time ~ treatment, data = Arthriti,  
col = c("lightblue", "lightgreen", "yellow"),  
ylab = "days")  
anova(lm(time ~ treatment, data = Arthriti))
```

Artifici

Durations of operation for 15 artificial heart transplants

Description

Data for Exercise 1.107

Usage

Artifici

Format

A data frame/tibble with 15 observations on one variable

duration duration (in hours) for transplant

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
stem(Artifici$duration, 2)  
summary(Artifici$duration)  
values <- Artifici$duration[Artifici$duration < 6.5]  
values  
summary(values)
```

 Asprin

Dissolving time versus level of impurities in aspirin tablets

Description

Data for Exercise 10.51

Usage

Asprin

Format

A data frame/tibble with 15 observations on two variables

time time (in seconds) for aspirin to dissolve

impurity impurity of an ingredient with levels 1%, 5%, and 10%

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
boxplot(time ~ impurity, data = Asprin,
        col = c("red", "blue", "green"))
```

 Asthmati

Asthmatic relief index on nine subjects given a drug and a placebo

Description

Data for Exercise 7.52

Usage

Asthmati

Format

A data frame/tibble with nine observations on three variables

drug asthmatic relief index for patients given a drug

placebo asthmatic relief index for patients given a placebo

difference difference between the placebo and drug

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
qqnorm(Asthmati$difference)
qqline(Asthmati$difference)
shapiro.test(Asthmati$difference)
with(data = Asthmati,
      t.test(placebo, drug, paired = TRUE, mu = 0, alternative = "greater")
)
```

Attorney

Number of convictions reported by U.S. attorney's offices

Description

Data for Example 2.2 and Exercises 2.43 and 2.57

Usage

Attorney

Format

A data frame/tibble with 88 observations on three variables

staff U.S. attorneys' office staff per 1 million population

convict U.S. attorneys' office convictions per 1 million population

district a factor with levels Albuquerque, Alexandria, Va, Anchorage, Asheville, NC, Atlanta, Baltimore, Baton Rouge, Billings, Mt, Birmingham, Al, Boise, Id, Boston, Buffalo, Burlington, Vt, Cedar Rapids, Charleston, WVA, Cheyenne, Wy, Chicago, Cincinnati, Cleveland, Columbia, SC, Concord, NH, Denver, Des Moines, Detroit, East St. Louis, Fargo, ND, Fort Smith, Ark, Fort Worth, Grand Rapids, Mi, Greensboro, NC, Honolulu, Houston, Indianapolis, Jackson, Miss, Kansas City, Knoxville, Tn, Las Vegas, Lexington, Ky, Little Rock, Los Angeles, Louisville, Memphis, Miami, Milwaukee, Minneapolis, Mobile, Ala, Montgomery, Ala, Muskogee, Ok, Nashville, New Haven, Conn, New Orleans, New York (Brooklyn), New York (Manhattan), Newark, NJ, Oklahoma City, Omaha, Oxford, Miss, Pensacola, Fl, Philadelphia, Phoenix, Pittsburgh, Portland, Maine, Portland, Ore, Providence, RI, Raleigh, NC, Roanoke, Va, Sacramento, Salt Lake City, San Antonio, San Diego, San Francisco, Savannah, Ga, Scranton, Pa, Seattle, Shreveport, La, Sioux Falls, SD, South Bend, Ind, Spokane, Wash, Springfield, Ill, St. Louis, Syracuse, NY, Tampa, Topeka, Kan, Tulsa, Tyler, Tex, Washington, Wheeling, WVa, and Wilmington, Del

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
par(mfrow=c(1, 2))
plot(convict ~ staff, data = Attorney, main = "With Washington, D.C.")
plot(convict[-86] ~staff[-86], data = Attorney,
main = "Without Washington, D.C.")
par(mfrow=c(1, 1))
```

Autogear

Number of defective auto gears produced by two manufacturers

Description

Data for Exercise 7.46

Usage

Autogear

Format

A data frame/tibble with 20 observations on two variables

defectives number of defective gears in the production of 100 gears per day

manufacturer a factor with levels A and B

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
t.test(defectives ~ manufacturer, data = Autogear)
wilcox.test(defectives ~ manufacturer, data = Autogear)
t.test(defectives ~ manufacturer, var.equal = TRUE, data = Autogear)
```

Backtoback	<i>Illustrates inferences based on pooled t-test versus Wilcoxon rank sum test</i>
------------	--

Description

Data for Exercise 7.40

Usage

Backtoback

Format

A data frame/tibble with 24 observations on two variables

score a numeric vector

group a numeric vector

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
wilcox.test(score ~ group, data = Backtoback)
t.test(score ~ group, data = Backtoback)
```

Bbsalaries	<i>Baseball salaries for members of five major league teams</i>
------------	---

Description

Data for Exercise 1.11

Usage

Bbsalaries

Format

A data frame/tibble with 142 observations on two variables

salary 1999 salary for baseball player

team a factor with levels Angels, Indians, Orioles, Redsoxs, and Whitesoxs

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
stripchart(salary ~ team, data = Bbsalaries, method = "stack",
           pch = 19, col = "blue", cex = 0.75)
title(main = "Major League Salaries")
```

Bigten	<i>Graduation rates for student athletes and nonathletes in the Big Ten Conf.</i>
--------	---

Description

Data for Exercises 1.124 and 2.94

Usage

Bigten

Format

A data frame/tibble with 44 observations on the following four variables

school a factor with levels Illinois, Indiana, Iowa, Michigan, Michigan State, Minnesota, Northwestern, Ohio State, Penn State, Purdue, and Wisconsin

rate graduation rate

year factor with two levels 1984–1985 and 1993–1994

status factor with two levels athlete and student

Source

NCAA Graduation Rates Report, 2000.

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
boxplot(rate ~ status, data = subset(Bigten, year = "1993-1994"),
        horizontal = TRUE, main = "Graduation Rates 1993-1994")
with(data = Bigten,
     tapply(rate, list(year, status), mean)
)
```

Biology

Test scores on first exam in biology class

Description

Data for Exercise 1.49

Usage

Biology

Format

A data frame/tibble with 30 observations on one variable

score test scores on the first test in a beginning biology class

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
hist(Biology$score, breaks = "scott", col = "brown", freq = FALSE,  
main = "Problem 1.49", xlab = "Test Score")  
lines(density(Biology$score), lwd=3)
```

Birth

Live birth rates in 1990 and 1998 for all states

Description

Data for Example 1.10

Usage

Birth

Format

A data frame/tibble with 51 observations on three variables

state a character with levels Alabama, Alaska, Arizona, Arkansas, California, Colorado, Connecticut, Delaware, District of Columbia, Florida, Georgia, Hawaii, Idaho, Illinois, Indiana, Iowa, Kansas, Kentucky, Louisiana, Maine, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Montana, Nebraska, Nevada, New Hampshire, New Jersey, New Mexico, New York, North Carolina, North Dakota, Ohio, Oklahoma, Oregon, Pennsylvania, Rhode Island, South Carolina, South Dakota, Tennessee, Texas, Utah, Vermont, Virginia, Washington, West Virginia, Wisconsin, and Wyoming

rate live birth rates per 1000 population

year a factor with levels 1990 and 1998

Source

National Vital Statistics Report, 48, March 28, 2000, National Center for Health Statistics.

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
rate1998 <- subset(Birth, year == "1998", select = rate)
stem(x = rate1998$rate, scale = 2)
hist(rate1998$rate, breaks = seq(10.9, 21.9, 1.0), xlab = "1998 Birth Rate",
      main = "Figure 1.14 in BSDA", col = "pink")
hist(rate1998$rate, breaks = seq(10.9, 21.9, 1.0), xlab = "1998 Birth Rate",
      main = "Figure 1.16 in BSDA", col = "pink", freq = FALSE)
lines(density(rate1998$rate), lwd = 3)
rm(rate1998)
```

Blackedu

Education level of blacks by gender

Description

Data for Exercise 8.55

Usage

Blackedu

Format

A data frame/tibble with 3800 observations on two variables

gender a factor with levels Female and Male

education a factor with levels High school dropout, High school graudate, Some college, Bachelor's degree, and Graduate degree

Source

Bureau of Census data.

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
T1 <- xtabs(~gender + education, data = Blackedu)
T1
chisq.test(T1)
```

Blood

Blood pressure of 15 adult males taken by machine and by an expert

Description

Data for Exercise 7.84

Usage

Blood

Format

A data frame/tibble with 15 observations on the following two variables

machine blood pressure recorded from an automated blood pressure machine

expert blood pressure recorded by an expert using an at-home device

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
DIFF <- Blood$machine - Blood$expert
shapiro.test(DIFF)
qqnorm(DIFF)
qqline(DIFF)
rm(DIFF)
t.test(Blood$machine, Blood$expert, paired = TRUE)
```

Board

*Incomes of board members from three different universities***Description**

Data for Exercise 10.14

Usage

Board

Format

A data frame/tibble with 7 observations on three variables

salary 1999 salary (in \$1000) for board directors

university a factor with levels A, B, and C

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
boxplot(salary ~ university, data = Board, col = c("red", "blue", "green"),
        ylab = "Income")
tapply(Board$salary, Board$university, summary)
anova(lm(salary ~ university, data = Board))
## Not run:
library(dplyr)
dplyr::group_by(Board, university) %>%
  summarize(Average = mean(salary))

## End(Not run)
```

Bones	<i>Bone density measurements of 35 physically active and 35 non-active women</i>
-------	--

Description

Data for Example 7.22

Usage

Bones

Format

A data frame/tibble with 70 observations on two variables

density bone density measurements

group a factor with levels active and nonactive

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
t.test(density ~ group, data = Bones, alternative = "greater")
t.test(rank(density) ~ group, data = Bones, alternative = "greater")
wilcox.test(density ~ group, data = Bones, alternative = "greater")
```

Books	<i>Number of books read and final spelling scores for 17 third graders</i>
-------	--

Description

Data for Exercise 9.53

Usage

Books

Format

A data frame/tibble with 17 observations on two variables

book number of books read

spelling spelling score

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
plot(spelling ~ book, data = Books)
mod <- lm(spelling ~ book, data = Books)
summary(mod)
abline(mod, col = "blue", lwd = 2)
```

Bookstor

Prices paid for used books at three different bookstores

Description

Data for Exercise 10.30 and 10.31

Usage

Bookstor

Format

A data frame/tibble with 72 observations on two variables

dollars money obtained for selling textbooks

store a factor with levels A, B, and C

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
boxplot(dollars ~ store, data = Bookstor,
        col = c("purple", "lightblue", "cyan"))
kruskal.test(dollars ~ store, data = Bookstor)
```

Brain

Brain weight versus body weight of 28 animals

Description

Data for Exercises 2.15, 2.44, 2.58 and Examples 2.3 and 2.20

Usage

Brain

Format

A data frame/tibble with 28 observations on three variables

species a factor with levels African elephant, Asian Elephant, Brachiosaurus, Cat, Chimpanzee, Cow, Diplodocus, Donkey, Giraffe, Goat, Gorilla, Gray wolf, Guinea Pig, Hamster, Horse, Human, Jaguar, Kangaroo, Mole, Mouse, Mt Beaver, Pig, Potar monkey, Rabbit, Rat, Rhesus monkey, Sheep, and Triceratops

bodyweight body weight (in kg)

brainweight brain weight (in g)

Source

P. Rousseeuw and A. Leroy, *Robust Regression and Outlier Detection* (New York: Wiley, 1987).

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
plot(log(brainweight) ~ log(bodyweight), data = Brain,
     pch = 19, col = "blue", main = "Example 2.3")
mod <- lm(log(brainweight) ~ log(bodyweight), data = Brain)
abline(mod, lty = "dashed", col = "blue")
```

Bumpers

Repair costs of vehicles crashed into a barrier at 5 miles per hour

Description

Data for Exercise 1.73

Usage

Bumpers

Format

A data frame/tibble with 23 observations on two variables

car a factor with levels Buick Century, Buick Skylark, Chevrolet Cavalier, Chevrolet Corsica, Chevrolet Lumina, Dodge Dynasty, Dodge Monaco, Ford Taurus, Ford Tempo, Honda Accord, Hyundai Sonata, Mazda 626, Mitsubishi Galant, Nissan Stanza, Oldsmobile Calais, Oldsmobile Ciere, Plymouth Acclaim, Pontiac 6000, Pontiac Grand Am, Pontiac Sunbird, Saturn SL2, Subaru Legacy, and Toyota Camry

repair total repair cost (in dollars) after crashing a car into a barrier four times while the car was traveling at 5 miles per hour

Source

Insurance Institute of Highway Safety.

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
EDA(Bumpers$repair)
stripchart(Bumpers$repair, method = "stack", pch = 19, col = "blue")
library(lattice)
dotplot(car ~ repair, data = Bumpers)
```

Bus*Attendance of bus drivers versus shift*

Description

Data for Exercise 8.25

Usage

Bus

Format

A data frame/tibble with 29363 observations on two variables

attendance a factor with levels absent and present

shift a factor with levels am, noon, pm, swing, and split

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
T1 <- xtabs(~attendance + shift, data = Bus)
T1
chisq.test(T1)
```

Bypass*Median charges for coronary bypass at 17 hospitals in North Carolina*

Description

Data for Exercises 5.104 and 6.43

Usage

Bypass

Format

A data frame/tibble with 17 observations on two variables

hospital a factor with levels Carolinas Med Ct, Duke Med Ct, Durham Regional, Forsyth Memorial, Frye Regional, High Point Regional, Memorial Mission, Mercy, Moore Regional, Moses Cone Memorial, NC Baptist, New Hanover Regional, Pitt Co. Memorial, Presbyterian, Rex, Univ of North Carolina, and Wake County

charge median charge for coronary bypass

Source

Consumer's Guide to Hospitalization Charges in North Carolina Hospitals (August 1994), North Carolina Medical Database Commission, Department of Insurance.

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
EDA(Bypass$charge)
t.test(Bypass$charge, conf.level=.90)$conf
t.test(Bypass$charge, mu = 35000)
```

Cabinets

Estimates of costs of kitchen cabinets by two suppliers on 20 prospective homes

Description

Data for Exercise 7.83

Usage

Cabinets

Format

A data frame/tibble with 20 observations on three variables

home a numeric vector

supplA estimate for kitchen cabinets from supplier A (in dollars)

supplB estimate for kitchen cabinets from supplier A (in dollars)

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
DIF <- Cabinets$supplA - Cabinets$supplB
qqnorm(DIF)
qqline(DIF)
shapiro.test(DIF)
with(data = Cabinets,
      t.test(supplA, supplB, paired = TRUE)
)
with(data = Cabinets,
      wilcox.test(supplA, supplB, paired = TRUE)
)
rm(DIF)
```

Cancer

Survival times of terminal cancer patients treated with vitamin C

Description

Data for Exercises 6.55 and 6.64

Usage

Cancer

Format

A data frame/tibble with 64 observations on two variables

survival survival time (in days) of terminal patients treated with vitamin C

type a factor indicating type of cancer with levels breast, bronchus, colon, ovary, and stomach

Source

Cameron, E and Pauling, L. 1978. "Supplemental Ascorbate in the Supportive Treatment of Cancer." *Proceedings of the National Academy of Science*, 75, 4538-4542.

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```

boxplot(survival ~ type, Cancer, col = "blue")
stomach <- Cancer$survival[Cancer$type == "stomach"]
bronchus <- Cancer$survival[Cancer$type == "bronchus"]
boxplot(stomach, ylab = "Days")
SIGN.test(stomach, md = 100, alternative = "greater")
SIGN.test(bronchus, md = 100, alternative = "greater")
rm(bronchus, stomach)

```

Carbon

*Carbon monoxide level measured at three industrial sites***Description**

Data for Exercise 10.28 and 10.29

Usage

Carbon

Format

A data frame/tibble with 24 observations on two variables

CO carbon monoxide measured (in parts per million)

site a factor with levels SiteA, SiteB, and SiteC

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```

boxplot(CO ~ site, data = Carbon, col = "lightgreen")
kruskal.test(CO ~ site, data = Carbon)

```

Cat	<i>Reading scores on the California achievement test for a group of 3rd graders</i>
-----	---

Description

Data for Exercise 1.116

Usage

Cat

Format

A data frame/tibble with 17 observations on one variable

score reading score on the California Achievement Test

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
stem(Cat$score)
fivenum(Cat$score)
boxplot(Cat$score, main = "Problem 1.116", col = "green")
```

Censored	<i>Entry age and survival time of patients with small cell lung cancer under two different treatments</i>
----------	---

Description

Data for Exercises 7.34 and 7.48

Usage

Censored

Format

A data frame/tibble with 121 observations on three variables

survival survival time (in days) of patients with small cell lung cancer

treatment a factor with levels armA and armB indicating the treatment a patient received

age the age of the patient

Source

Ying, Z., Jung, S., Wei, L. 1995. "Survival Analysis with Median Regression Models." *Journal of the American Statistical Association*, 90, 178-184.

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
boxplot(survival ~ treatment, data = Censored, col = "yellow")
wilcox.test(survival ~ treatment, data = Censored, alternative = "greater")
```

Challeng	<i>Temperatures and O-ring failures for the launches of the space shuttle Challenger</i>
----------	--

Description

Data for Examples 1.11, 1.12, 1.13, 2.11 and 5.1

Usage

Challeng

Format

A data frame/tibble with 25 observations on four variables

flight a character variable indicating the flight

date date of the flight

temp temperature (in fahrenheit)

failures number of failures

Source

Dalal, S. R., Fowlkes, E. B., Hoadley, B. 1989. "Risk Analysis of the Space Shuttle: Pre-Challenger Prediction of Failure." *Journal of the American Statistical Association*, 84, No. 408, 945-957.

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```

stem(Challeng$temp)
summary(Challeng$temp)
IQR(Challeng$temp)
quantile(Challeng$temp)
fivenum(Challeng$temp)
stem(sort(Challeng$temp)[-1])
summary(sort(Challeng$temp)[-1])
IQR(sort(Challeng$temp)[-1])
quantile(sort(Challeng$temp)[-1])
fivenum(sort(Challeng$temp)[-1])
par(mfrow=c(1, 2))
qqnorm(Challeng$temp)
qqline(Challeng$temp)
qqnorm(sort(Challeng$temp)[-1])
qqline(sort(Challeng$temp)[-1])
par(mfrow=c(1, 1))

```

Chemist

Starting salaries of 50 chemistry majors

Description

Data for Example 5.3

Usage

Chemist

Format

A data frame/tibble with 50 observations on one variable

salary starting salary (in dollars) for chemistry major

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
EDA(Chemist$salary)
```

Chesapea	<i>Surface salinity measurements taken offshore from Annapolis, Maryland in 1927</i>
----------	--

Description

Data for Exercise 6.41

Usage

Chesapea

Format

A data frame/tibble with 16 observations on one variable

salinity surface salinity measurements (in parts per 1000) for station 11, offshore from Annapolis, Maryland, on July 3-4, 1927.

Source

Davis, J. (1986) *Statistics and Data Analysis in Geology, Second Edition*. John Wiley and Sons, New York.

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
qqnorm(Chesapea$salinity)
qqline(Chesapea$salinity)
shapiro.test(Chesapea$salinity)
t.test(Chesapea$salinity, mu = 7)
```

Chevy	<i>Insurance injury ratings of Chevrolet vehicles for 1990 and 1993 models</i>
-------	--

Description

Data for Exercise 8.35

Usage

Chevy

Format

A data frame/tibble with 67 observations on two variables

year a factor with levels 1988-90 and 1991-93

frequency a factor with levels much better than average, above average, average, below average, and much worse than average

Source

Insurance Institute for Highway Safety and the Highway Loss Data Institute, 1995.

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
T1 <- xtabs(~year + frequency, data = Chevy)
T1
chisq.test(T1)
rm(T1)
```

Chicken

Weight gain of chickens fed three different rations

Description

Data for Exercise 10.15

Usage

Chicken

Format

A data frame/tibble with 13 observations on three variables

gain weight gain over a specified period

feed a factor with levels ration1, ration2, and ration3

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
boxplot(gain ~ feed, col = c("red", "blue", "green"), data = Chicken)
anova(lm(gain ~ feed, data = Chicken))
```

Chipavg

Measurements of the thickness of the oxide layer of manufactured integrated circuits

Description

Data for Exercises 6.49 and 7.47

Usage

Chipavg

Format

A data frame/tibble with 30 observations on three variables

wafer1 thickness of the oxide layer for wafer1

wafer2 thickness of the oxide layer for wafer2

thickness average thickness of the oxide layer of the eight measurements obtained from each set of two wafers

Source

Yashchin, E. 1995. "Likelihood Ratio Methods for Monitoring Parameters of a Nested Random Effect Model." *Journal of the American Statistical Association*, 90, 729-738.

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
EDA(Chipavg$thickness)
t.test(Chipavg$thickness, mu = 1000)
boxplot(Chipavg$wafer1, Chipavg$wafer2, name = c("Wafer 1", "Wafer 2"))
shapiro.test(Chipavg$wafer1)
shapiro.test(Chipavg$wafer2)
t.test(Chipavg$wafer1, Chipavg$wafer2, var.equal = TRUE)
```

Chips	<i>Four measurements on a first wafer and four measurements on a second wafer selected from 30 lots</i>
-------	---

Description

Data for Exercise 10.9

Usage

Chips

Format

A data frame/tibble with 30 observations on eight variables

wafer11 first measurement of thickness of the oxide layer for wafer1

wafer12 second measurement of thickness of the oxide layer for wafer1

wafer13 third measurement of thickness of the oxide layer for wafer1

wafer14 fourth measurement of thickness of the oxide layer for wafer1

wafer21 first measurement of thickness of the oxide layer for wafer2

wafer22 second measurement of thickness of the oxide layer for wafer2

wafer23 third measurement of thickness of the oxide layer for wafer2

wafer24 fourth measurement of thickness of the oxide layer for wafer2

Source

Yashchin, E. 1995. "Likelihood Ratio Methods for Monitoring Parameters of a Nested Random Effect Model." *Journal of the American Statistical Association*, 90, 729-738.

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
with(data = Chips,  
      boxplot(wafer11, wafer12, wafer13, wafer14, wafer21,  
              wafer22, wafer23, wafer24, col = "pink")  
)
```

Cigar	<i>Milligrams of tar in 25 cigarettes selected randomly from 4 different brands</i>
-------	---

Description

Data for Example 10.4

Usage

Cigar

Format

A data frame/tibble with 100 observations on two variables

tar amount of tar (measured in milligrams)

brand a factor indicating cigarette brand with levels brandA, brandB, brandC, and brandD

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
boxplot(tar ~ brand, data = Cigar, col = "cyan", ylab = "mg tar")
anova(lm(tar ~ brand, data = Cigar))
```

Cigarett	<i>Effect of mother's smoking on birth weight of newborn</i>
----------	--

Description

Data for Exercise 2.27

Usage

Cigarett

Format

A data frame/tibble with 16 observations on two variables

cigarettes mothers' estimated average number of cigarettes smoked per day

weight children's birth weights (in pounds)

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
plot(weight ~ cigarettes, data = Cigarette)
model <- lm(weight ~ cigarettes, data = Cigarette)
abline(model, col = "red")
with(data = Cigarette,
      cor(weight, cigarettes)
)
rm(model)
```

CIsim

Confidence Interval Simulation Program

Description

This program simulates random samples from which it constructs confidence intervals for one of the parameters mean (μ), variance (σ^2), or proportion of successes (π).

Usage

```
CIsim(
  samples = 100,
  n = 30,
  mu = 0,
  sigma = 1,
  conf.level = 0.95,
  type = "Mean"
)
```

Arguments

<code>samples</code>	the number of samples desired.
<code>n</code>	the size of each sample.
<code>mu</code>	if constructing confidence intervals for the population mean or the population variance, <code>mu</code> is the population mean (i.e., <code>type</code> is one of either "Mean", or "Var"). If constructing confidence intervals for the population proportion of successes, the value entered for <code>mu</code> represents the population proportion of successes (π), and as such, must be a number between 0 and 1.
<code>sigma</code>	the population standard deviation. <code>sigma</code> is not required if confidence intervals are of type "Pi".
<code>conf.level</code>	confidence level for the graphed confidence intervals, restricted to lie between zero and one.

type character string, one of "Mean", "Var" or "Pi", or just the initial letter of each, indicating the type of confidence interval simulation to perform.

Details

Default is to construct confidence intervals for the population mean. Simulated confidence intervals for the population variance or population proportion of successes are possible by selecting the appropriate value in the type argument.

Value

Graph depicts simulated confidence intervals. The number of confidence intervals that do not contain the parameter of interest are counted and reported in the commands window.

Author(s)

Alan T. Arnholt

Examples

```
CIsim(100, 30, 100, 10)
# Simulates 100 samples of size 30 from
# a normal distribution with mean 100
# and standard deviation 10. From the
# 100 simulated samples, 95% confidence
# intervals for the Mean are constructed
# and depicted in the graph.

CIsim(100, 30, 100, 10, type="Var")
# Simulates 100 samples of size 30 from
# a normal distribution with mean 100
# and standard deviation 10. From the
# 100 simulated samples, 95% confidence
# intervals for the variance are constructed
# and depicted in the graph.

CIsim(100, 50, .5, type="Pi", conf.level=.90)
# Simulates 100 samples of size 50 from
# a binomial distribution where the population
# proportion of successes is 0.5. From the
# 100 simulated samples, 90% confidence
# intervals for Pi are constructed
# and depicted in the graph.
```

Citrus	<i>Percent of peak bone density of different aged children</i>
--------	--

Description

Data for Exercise 9.7

Usage

Citrus

Format

A data frame/tibble with nine observations on two variables

age age of children

percent percent peak bone density

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
model <- lm(percent ~ age, data = Citrus)
summary(model)
anova(model)
rm(model)
```

Clean	<i>Residual contaminant following the use of three different cleansing agents</i>
-------	---

Description

Data for Exercise 10.16

Usage

Clean

Format

A data frame/tibble with 45 observations on two variables

clean residual contaminants

agent a factor with levels A, B, and C

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
boxplot(clean ~ agent, col = c("red", "blue", "green"), data = Clean)
anova(lm(clean ~ agent, data = Clean))
```

Coaxial

Signal loss from three types of coaxial cable

Description

Data for Exercise 10.24 and 10.25

Usage

Coaxial

Format

A data frame/tibble with 45 observations on two variables

signal signal loss per 1000 feet

cable factor with three levels of coaxial cable typeA, typeB, and typeC

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
boxplot(signal ~ cable, data = Coaxial, col = c("red", "green", "yellow"))
kruskal.test(signal ~ cable, data = Coaxial)
```

Coffee

Productivity of workers with and without a coffee break

Description

Data for Exercise 7.55

Usage

Coffee

Format

A data frame/tibble with nine observations on three variables

without workers' productivity scores without a coffee break

with workers' productivity scores with a coffee break

differences with minus without

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
qqnorm(Coffee$differences)
qqline(Coffee$differences)
shapiro.test(Coffee$differences)
t.test(Coffee$with, Coffee$without, paired = TRUE, alternative = "greater")
wilcox.test(Coffee$with, Coffee$without, paired = TRUE,
  alterantive = "greater")
```

Coins

Yearly returns on 12 investments

Description

Data for Exercise 5.68

Usage

Coins

Format

A data frame/tibble with 12 observations on one variable

return yearly returns on each of 12 possible investments

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
qqnorm(Coins$return)
qqline(Coins$return)
```

Combinations

Combinations

Description

Computes all possible combinations of n objects taken k at a time.

Usage

```
Combinations(n, k)
```

Arguments

n a number.
 k a number less than or equal to n .

Value

Returns a matrix containing the possible combinations of n objects taken k at a time.

See Also

[SRS](#)

Examples

```
Combinations(5,2)
# The columns in the matrix list the values of the 10 possible
# combinations of 5 things taken 2 at a time.
```

Commute

Commuting times for selected cities in 1980 and 1990

Description

Data for Exercises 1.13, and 7.85

Usage

Commute

Format

A data frame/tibble with 39 observations on three variables

city a factor with levels Atlanta, Baltimore, Boston, Buffalo, Charlotte, Chicago, Cincinnati, Cleveland, Columbus, Dallas, Denver, Detroit, Hartford, Houston, Indianapolis, Kansas City, Los Angeles, Miami, Milwaukee, Minneapolis, New Orleans, New York, Norfolk, Orlando, Philadelphia, Phoenix, Pittsburgh, Portland, Providence, Rochester, Sacramento, Salt Lake City, San Antonio, San Diego, San Francisco, Seattle, St. Louis, Tampa, and Washington

year year

time commute times

Source

Federal Highway Administration.

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
stripplot(year ~ time, data = Commute, jitter = TRUE)
dotplot(year ~ time, data = Commute)
bwplot(year ~ time, data = Commute)
stripchart(time ~ year, data = Commute, method = "stack", pch = 1,
           cex = 2, col = c("red", "blue"),
           group.names = c("1980", "1990"),
           main = "", xlab = "minutes")
title(main = "Commute Time")
boxplot(time ~ year, data = Commute, names=c("1980", "1990"),
        horizontal = TRUE, las = 1)
```

Concept	<i>Tennessee self concept scale scores for a group of teenage boys</i>
Description	
Data for Exercise 1.68 and 1.82	
Usage	
Concept	
Format	
A data frame/tibble with 28 observations on one variable	
self Tennessee self concept scores	
References	
Kitchens, L. J. (2003) <i>Basic Statistics and Data Analysis</i> . Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.	
Examples	
<pre>summary(Concept\$self) sd(Concept\$self) diff(range(Concept\$self)) IQR(Concept\$self) summary(Concept\$self/10) IQR(Concept\$self/10) sd(Concept\$self/10) diff(range(Concept\$self/10))</pre>	
Concrete	<i>Compressive strength of concrete blocks made by two different methods</i>

Description

Data for Example 7.17

Usage

Concrete

Format

A data frame/tibble with 20 observations on two variables

strength comprehensive strength (in pounds per square inch)

method factor with levels new and old indicating the method used to construct a concrete block

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
wilcox.test(strength ~ method, data = Concrete, alternative = "greater")
```

Corn	<i>Comparison of the yields of a new variety and a standard variety of corn planted on 12 plots of land</i>
------	---

Description

Data for Exercise 7.77

Usage

Corn

Format

A data frame/tibble with 12 observations on three variables

new corn yield with new meathod

standard corn yield with standard method

differences new minus standard

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
boxplot(Corn$differences)
qqnorm(Corn$differences)
qqline(Corn$differences)
shapiro.test(Corn$differences)
t.test(Corn$differences, alternative = "greater")
```

Correlat

Exercise to illustrate correlation

Description

Data for Exercise 2.23

Usage

Correlat

Format

A data frame/tibble with 13 observations on two variables

x a numeric vector

y a numeric vector

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
plot(y ~ x, data = Correlat)
model <- lm(y ~ x, data = Correlat)
abline(model)
rm(model)
```

Counsel

Scores of 18 volunteers who participated in a counseling process

Description

Data for Exercise 6.96

Usage

Counsel

Format

A data frame/tibble with 18 observations on one variable

score standardized psychology scores after a counseling process

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
EDA(Counsel$score)
t.test(Counsel$score, mu = 70)
```

Cpi

Consumer price index from 1979 to 1998

Description

Data for Exercise 1.34

Usage

Cpi

Format

A data frame/tibble with 20 observations on two variables

year year

cpi consumer price index

Source

Bureau of Labor Statistics.

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
plot(cpi ~ year, data = Cpi, type = "l", lty = 2, lwd = 2, col = "red")
barplot(Cpi$cpi, col = "pink", las = 2, main = "Problem 1.34")
```

Crime

Violent crime rates for the states in 1983 and 1993

Description

Data for Exercises 1.90, 2.32, 3.64, and 5.113

Usage

Crime

Format

A data frame/tibble with 102 observations on three variables

state a factor with levels Alabama, Alaska, Arizona, Arkansas, California, Colorado, Connecticut, DC, Delaware, Florida, Georgia, Hawaii, Idaho, Illinois, Indiana, Iowa, Kansas, Kentucky, Louisiana, Maine, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Montana, Nebraska, Nevada, New Hampshire, New Jersey, New Mexico, New York, North Carolina, North Dakota, Ohio, Oklahoma, Oregon, Pennsylvania, Rhode Island, South Carolina, South Dakota, Tennessee, Texas, Utah, Vermont, Virginia, Washington, West Virginia, Wisconsin, and Wyoming

year a factor with levels 1983 and 1993

rate crime rate per 100,000 inhabitants

Source

U.S. Department of Justice, Bureau of Justice Statistics, *Sourcebook of Criminal Justice Statistics*, 1993.

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
boxplot(rate ~ year, data = Crime, col = "red")
```

Darwin*Charles Darwin's study of cross-fertilized and self-fertilized plants*

Description

Data for Exercise 7.62

Usage

Darwin

Format

A data frame/tibble with 15 observations on three variables

pot number of pot

cross height of plant (in inches) after a fixed period of time when cross-fertilized

self height of plant (in inches) after a fixed period of time when self-fertilized

Source

Darwin, C. (1876) *The Effect of Cross- and Self-Fertilization in the Vegetable Kingdom*, 2nd edition, London.

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
differ <- Darwin$cross - Darwin$self
qqnorm(differ)
qqline(differ)
shapiro.test(differ)
wilcox.test(Darwin$cross, Darwin$self, paired = TRUE)
rm(differ)
```

Dealers	<i>Automobile dealers classified according to type dealership and service rendered to customers</i>
---------	---

Description

Data for Example 2.22

Usage

Dealers

Format

A data frame/tibble with 122 observations on two variables

type a factor with levels Honda, Toyota, Mazda, Ford, Dodge, and Saturn

service a factor with levels Replaces unnecessarily and Follows manufacturer guidelines

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
xtabs(~type + service, data = Dealers)
T1 <- xtabs(~type + service, data = Dealers)
T1
addmargins(T1)
pt <- prop.table(T1, margin = 1)
pt
barplot(t(pt), col = c("red", "skyblue"), legend = colnames(T1))
rm(T1, pt)
```

Defectiv	<i>Number of defective items produced by 20 employees</i>
----------	---

Description

Data for Exercise 1.27

Usage

Defectiv

Format

A data frame/tibble with 20 observations on one variable

number number of defective items produced by the employees in a small business firm

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
T1 <- xtabs(~ number, data = Defectiv)
T1
barplot(T1, col = "pink", ylab = "Frequency",
xlab = "Defective Items Produced by Employees", main = "Problem 1.27")
rm(T1)
```

Degree

Percent of bachelor's degrees awarded women in 1970 versus 1990

Description

Data for Exercise 2.75

Usage

Degree

Format

A data frame/tibble with 1064 observations on two variables

field a factor with levels Health, Education, Foreign Language, Psychology, Fine Arts, Life Sciences, Business, Social Science, Physical Sciences, Engineering, and All Fields

awarded a factor with levels 1970 and 1990

Source

U.S. Department of Health and Human Services, National Center for Education Statistics.

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
T1 <- xtabs(~field + awarded, data = Degree)
T1
barplot(t(T1), beside = TRUE, col = c("red", "skyblue"), legend = colnames(T1))
rm(T1)
```

Delay

Delay times on 20 flights from four major air carriers

Description

Data for Exercise 10.55

Usage

Delay

Format

A data frame/tibble with 80 observations on two variables

delay the delay time (in minutes) for 80 randomly selected flights

carrier a factor with levels A, B, C, and D

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
boxplot(delay ~ carrier, data = Delay,
        main = "Exercise 10.55", ylab = "minutes",
        col = "pink")
kruskal.test(delay ~ carrier, data = Delay)
```

Depend	<i>Number of dependent children for 50 families</i>
--------	---

Description

Data for Exercise 1.26

Usage

Depend

Format

A data frame/tibble with 50 observations on one variable

number number of dependent children in a family

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
T1 <- xtabs(~ number, data = Depend)
T1
barplot(T1, col = "lightblue", main = "Problem 1.26",
xlab = "Number of Dependent Children", ylab = "Frequency")
rm(T1)
```

Detroit	<i>Educational levels of a sample of 40 auto workers in Detroit</i>
---------	---

Description

Data for Exercise 5.21

Usage

Detroit

Format

A data frame/tibble with 40 observations on one variable

educ the educational level (in years) of a sample of 40 auto workers in a plant in Detroit

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
EDA(Detroit$educ)
```

Develop	<i>Demographic characteristics of developmental students at 2-year colleges and 4-year colleges</i>
---------	---

Description

Data used for Exercise 8.50

Usage

Develop

Format

A data frame/tibble with 5656 observations on two variables

race a factor with levels African American, American Indian, Asian, Latino, and White

college a factor with levels Two-year and Four-year

Source

Research in Development Education (1994), V. 11, 2.

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
T1 <- xtabs(~race + college, data = Develop)
T1
chisq.test(T1)
rm(T1)
```

Devmath	<i>Test scores for students who failed developmental mathematics in the fall semester 1995</i>
---------	--

Description

Data for Exercise 6.47

Usage

Devmath

Format

A data frame/tibble with 40 observations on one variable

score first exam score

Source

Data provided by Dr. Anita Kitchens.

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
EDA(Devmath$score)
t.test(Devmath$score, mu = 80, alternative = "less")
```

Dice	<i>Outcomes and probabilities of the roll of a pair of fair dice</i>
------	--

Description

Data for Exercise 3.109

Usage

Dice

Format

A data frame/tibble with 11 observations on two variables

x possible outcomes for the sum of two dice

px probability for outcome x

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
roll1 <- sample(1:6, 20000, replace = TRUE)
roll2 <- sample(1:6, 20000, replace = TRUE)
outcome <- roll1 + roll2
T1 <- table(outcome)/length(outcome)
remove(roll1, roll2, outcome)
T1
round(t(Dice), 5)
rm(roll1, roll2, T1)
```

Diesel

Diesel fuel prices in 1999-2000 in nine regions of the country

Description

Data for Exercise 2.8

Usage

Diesel

Format

A data frame/tibble with 650 observations on three variables

date date when price was recorded

pricepergallon price per gallon (in dollars)

location a factor with levels California, CentralAtlantic, Coast, EastCoast, Gulf, LowerAtlantic, NatAvg, NorthEast, Rocky, and WesternMountain

Source

Energy Information Administration, National Energy Information Center: 1000 Independence Ave., SW, Washington, D.C., 20585.

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
par(las = 2)
boxplot(pricepergallon ~ location, data = Diesel)
boxplot(pricepergallon ~ location,
        data = droplevels(Diesel[Diesel$location == "EastCoast" |
                             Diesel$location == "Gulf" | Diesel$location == "NatAvg" |
                             Diesel$location == "Rocky" | Diesel$location == "California", ]),
        col = "pink", main = "Exercise 2.8")
par(las = 1)
## Not run:
library(ggplot2)
ggplot2::ggplot(data = Diesel, aes(x = date, y = pricepergallon,
                                   color = location)) +
  geom_point() +
  geom_smooth(se = FALSE) +
  theme_bw() +
  labs(y = "Price per Gallon (in dollars)")

## End(Not run)
```

Diplomat

Parking tickets issued to diplomats

Description

Data for Exercises 1.14 and 1.37

Usage

Diplomat

Format

A data frame/tibble with 10 observations on three variables

country a factor with levels Brazil, Bulgaria, Egypt, Indonesia, Israel, Nigeria, Russia, S. Korea, Ukraine, and Venezuela

number total number of tickets

rate number of tickets per vehicle per month

Source

Time, November 8, 1993. Figures are from January to June 1993.

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
par(las = 2, mfrow = c(2, 2))
stripchart(number ~ country, data = Diplomat, pch = 19,
           col= "red", vertical = TRUE)
stripchart(rate ~ country, data = Diplomat, pch = 19,
           col= "blue", vertical = TRUE)
with(data = Diplomat,
     barplot(number, names.arg = country, col = "red"))
with(data = Diplomat,
     barplot(rate, names.arg = country, col = "blue"))
par(las = 1, mfrow = c(1, 1))
## Not run:
library(ggplot2)
ggplot2::ggplot(data = Diplomat, aes(x = reorder(country, number),
                                     y = number)) +
  geom_bar(stat = "identity", fill = "pink", color = "black") +
  theme_bw() + labs(x = "", y = "Total Number of Tickets")
ggplot2::ggplot(data = Diplomat, aes(x = reorder(country, rate),
                                     y = rate)) +
  geom_bar(stat = "identity", fill = "pink", color = "black") +
  theme_bw() + labs(x = "", y = "Tickets per vehicle per month")

## End(Not run)
```

Disposal	<i>Toxic intensity for manufacturing plants producing herbicidal preparations</i>
----------	---

Description

Data for Exercise 1.127

Usage

Disposal

Format

A data frame/tibble with 29 observations on one variable
pounds pounds of toxic waste per \$1000 of shipments of its products

Source

Bureau of the Census, *Reducing Toxins*, Statistical Brief SB/95-3, February 1995.

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
stem(Disposal$pounds)
fivenum(Disposal$pounds)
EDA(Disposal$pounds)
```

Dogs	<i>Rankings of the favorite breeds of dogs</i>
------	--

Description

Data for Exercise 2.88

Usage

Dogs

Format

A data frame/tibble with 20 observations on three variables

breed a factor with levels Beagle, Boxer, Chihuahua, Chow, Dachshund, Dalmatian, Doberman, Huskie, Labrador, Pomeranian, Poodle, Retriever, Rotweiler, Schnauzer, Shepherd, Shetland, ShihTzu, Spaniel, Springer, and Yorkshire

ranking numeric ranking

year a factor with levels 1992, 1993, 1997, and 1998

Source

The World Almanac and Book of Facts, 2000.

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
cor(Dogs$ranking[Dogs$year == "1992"], Dogs$ranking[Dogs$year == "1993"])
cor(Dogs$ranking[Dogs$year == "1997"], Dogs$ranking[Dogs$year == "1998"])
## Not run:
library(ggplot2)
ggplot2::ggplot(data = Dogs, aes(x = reorder(breed, ranking), y = ranking)) +
  geom_bar(stat = "identity") +
  facet_grid(year ~. ) +
  theme(axis.text.x = element_text(angle = 85, vjust = 0.5))

## End(Not run)
```

Domestic

*Rates of domestic violence per 1,000 women by age groups***Description**

Data for Exercise 1.20

Usage

Domestic

Format

A data frame/tibble with five observations on two variables

age a factor with levels 12–19, 20–24, 25–34, 35–49, and 50–64

rate rate of domestic violence per 1000 women

Source

U.S. Department of Justice.

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
barplot(Domestic$rate, names.arg = Domestic$age)
## Not run:
library(ggplot2)
ggplot2::ggplot(data = Domestic, aes(x = age, y = rate)) +
  geom_bar(stat = "identity", fill = "purple", color = "black") +
  labs(x = "", y = "Domestic violence per 1000 women") +
  theme_bw()

## End(Not run)
```

Dopamine	<i>Dopamine b-hydroxylase activity of schizophrenic patients treated with an antipsychotic drug</i>
----------	---

Description

Data for Exercises 5.14 and 7.49

Usage

Dopamine

Format

A data frame/tibble with 25 observations on two variables

dbh dopamine b-hydroxylase activity (units are nmol/(ml)(h)/(mg) of protein)

group a factor with levels nonpsychotic and psychotic

Source

D.E. Sternberg, D.P. Van Kammen, and W.E. Bunney, "Schizophrenia: Dopamine b-Hydroxylase Activity and Treatment Response," *Science*, 216 (1982), 1423 - 1425.

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
boxplot(dbh ~ group, data = Dopamine, col = "orange")
t.test(dbh ~ group, data = Dopamine, var.equal = TRUE)
```

Dowjones	<i>Closing yearend Dow Jones Industrial averages from 1896 through 2000</i>
----------	---

Description

Data for Exercise 1.35

Usage

Dowjones

Format

A data frame/tibble with 105 observations on three variables

year date

close Dow Jones closing price

change percent change from previous year

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
plot(close ~ year, data = Dowjones, type = "l", main = "Exercise 1.35")
## Not run:
library(ggplot2)
ggplot2::ggplot(data = Dowjones, aes(x = year, y = close)) +
  geom_point(size = 0.5) +
  geom_line(color = "red") +
  theme_bw() +
  labs(y = "Dow Jones Closing Price")

## End(Not run)
```

Drink

Opinion on referendum by view on moral issue of selling alcoholic beverages

Description

Data for Exercise 8.53

Usage

Drink

Format

A data frame/tibble with 472 observations on two variables

drinking a factor with levels ok, tolerated, and immoral

referendum a factor with levels for, against, and undecided

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
T1 <- xtabs(~drinking + referendum, data = Drink)
T1
chisq.test(T1)
rm(T1)
```

Drug	<i>Number of trials to master a task for a group of 28 subjects assigned to a control and an experimental group</i>
------	---

Description

Data for Example 7.15

Usage

Drug

Format

A data frame/tibble with 28 observations on two variables

trials number of trials to master a task

group a factor with levels control and experimental

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
boxplot(trials ~ group, data = Drug,
        main = "Example 7.15", col = c("yellow", "red"))
wilcox.test(trials ~ group, data = Drug)
t.test(rank(trials) ~ group, data = Drug, var.equal = TRUE)
```

Dyslexia

Data on a group of college students diagnosed with dyslexia

Description

Data for Exercise 2.90

Usage

Dyslexia

Format

A data frame/tibble with eight observations on seven variables

words number of words read per minute

age age of participant

gender a factor with levels female and male

handed a factor with levels left and right

weight weight of participant (in pounds)

height height of participant (in inches)

children number of children in family

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
plot(height ~ weight, data = Dyslexia)
plot(words ~ factor(handed), data = Dyslexia,
      xlab = "hand", col = "lightblue")
```

Earthqk	<i>One hundred year record of worldwide seismic activity(1770-1869)</i>
---------	---

Description

Data for Exercise 6.97

Usage

Earthqk

Format

A data frame/tibble with 100 observations on two variables

year year seismic activity recorded

severity annual incidence of sever earthquakes

Source

Quenoille, M.H. (1952), *Associated Measurements*, Butterworth, London. p 279.

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
EDA(Earthqk$severity)
t.test(Earthqk$severity, mu = 100, alternative = "greater")
```

EDA	<i>Exploratory Data Anaalysis</i>
-----	-----------------------------------

Description

Function that produces a histogram, density plot, boxplot, and Q-Q plot.

Usage

```
EDA(x, trim = 0.05)
```

Arguments

<code>x</code>	numeric vector. NAs and Infs are allowed but will be removed.
<code>trim</code>	fraction (between 0 and 0.5, inclusive) of values to be trimmed from each end of the ordered data. If <code>trim = 0.5</code> , the result is the median.

Details

Will not return command window information on data sets containing more than 5000 observations. It will however still produce graphical output for data sets containing more than 5000 observations.

Value

Function returns various measures of center and location. The values returned for the Quartiles are based on the definitions provided in *BSDA*. The boxplot is based on the Quartiles returned in the commands window.

Note

Requires package **e1071**.

Author(s)

Alan T. Arnholt

Examples

```
EDA(rnorm(100))
# Produces four graphs for the 100 randomly
# generated standard normal variates.
```

Educat	<i>Crime rates versus the percent of the population without a high school degree</i>
--------	--

Description

Data for Exercise 2.41

Usage

Educat

Format

A data frame/tibble with 51 observations on three variables

state a factor with levels Alabama, Alaska, Arizona, Arkansas, California, Colorado, Connecticut, DC, Delaware, Florida, Georgia, Hawaii, Idaho, Illinois, Indiana, Iowa, Kansas, Kentucky, Louisiana, Maine, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Montana, Nebraska, Nevada, New Hampshire, New Jersey, New Mexico, New York, North Carolina, North Dakota, Ohio, Oklahoma, Oregon, Pennsylvania, Rhode Island, South Carolina, South Dakota, Tennessee, Texas, Utah, Vermont, Virginia, Washington, West Virginia, Wisconsin, and Wyoming

nodegree percent of the population without a high school degree

crime violent crimes per 100,000 population

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
plot(crime ~ nodegree, data = Educat,
     xlab = "Percent of population without high school degree",
     ylab = "Violent Crime Rate per 100,000")
```

Eggs

Number of eggs versus amounts of feed supplement

Description

Data for Exercise 9.22

Usage

Eggs

Format

A data frame/tibble with 12 observations on two variables

feed amount of feed supplement

eggs number of eggs per day for 100 chickens

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
plot(eggs ~ feed, data = Eggs)
model <- lm(eggs ~ feed, data = Eggs)
abline(model, col = "red")
summary(model)
rm(model)
```

Elderly

*Percent of the population over the age of 65***Description**

Data for Exercise 1.92 and 2.61

Usage

Elderly

Format

A data frame/tibble with 51 observations on three variables

state a factor with levels Alabama, Alaska, Arizona, Arkansas, California, Colorado, Connecticut, Delaware, District of Columbia, Florida, Georgia, Hawaii, Idaho, Illinois, Indiana, Iowa, Kansas, Kentucky, Louisiana, Maine, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Montana, Nebraska, Nevada, New Hampshire, New Jersey, New Mexico, New York, North Carolina, North Dakota, Ohio, Oklahoma, Oregon, Pennsylvania, Rhode Island, South Carolina, South Dakota, Tennessee, Texas, Utah, Vermont, Virginia, Washington, West Virginia, Wisconsin, and Wyoming

percent1985 percent of the population over the age of 65 in 1985

percent1998 percent of the population over the age of 65 in 1998

Source

U.S. Census Bureau Internet site, February 2000.

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```

with(data = Elderly,
  stripchart(x = list(percent1998, percent1985), method = "stack", pch = 19,
    col = c("red", "blue"), group.names = c("1998", "1985"))
  )
with(data = Elderly, cor(percent1998, percent1985))
## Not run:
library(ggplot2)
ggplot2::ggplot(data = Elderly, aes(x = percent1985, y = percent1998)) +
  geom_point() +
  theme_bw()

## End(Not run)

```

Energy

Amount of energy consumed by homes versus their sizes

Description

Data for Exercises 2.5, 2.24, and 2.55

Usage

Energy

Format

A data frame/tibble with 12 observations on two variables

size size of home (in square feet)

kilowatt kilowatt-hours per month

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```

plot(kilowatt ~ size, data = Energy)
with(data = Energy, cor(size, kilowatt))
model <- lm(kilowatt ~ size, data = Energy)
plot(Energy$size, resid(model), xlab = "size")

```

Engineer

Salaries after 10 years for graduates of three different universities

Description

Data for Example 10.7

Usage

Engineer

Format

A data frame/tibble with 51 observations on two variables

salary salary (in \$1000) 10 years after graduation

university a factor with levels A, B, and C

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
boxplot(salary ~ university, data = Engineer,
        main = "Example 10.7", col = "yellow")
kruskal.test(salary ~ university, data = Engineer)
anova(lm(salary ~ university, data = Engineer))
anova(lm(rank(salary) ~ university, data = Engineer))
```

Entrance

College entrance exam scores for 24 high school seniors

Description

Data for Example 1.8

Usage

Entrance

Format

A data frame/tibble with 24 observations on one variable

score college entrance exam score

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
stem(Entrance$score)
stem(Entrance$score, scale = 2)
```

Epaminicompact	<i>Fuel efficiency ratings for compact vehicles in 2001</i>
----------------	---

Description

Data for Exercise 1.65

Usage

Epaminicompact

Format

A data frame/tibble with 22 observations on ten variables

class a character variable with value MINICOMPACT CARS

manufacturer a character variable with values AUDI, BMW, JAGUAR, MERCEDES-BENZ, MITSUBISHI, and PORSCHE

carline a character variable with values 325CI CONVERTIBLE, 330CI CONVERTIBLE, 911 CARRERA 2/4, 911 TURBO, CLK320 (CABRIOLET), CLK430 (CABRIOLET), ECLIPSE SPYDER, JAGUAR XK8 CONVERTIBLE, JAGUAR XKR CONVERTIBLE, M3 CONVERTIBLE, TT COUPE, and TT COUPE QUATTRO

displ engine displacement (in liters)

cyl number of cylinders

trans a factor with levels Auto(L5), Auto(S4), Auto(S5), Manual(M5), and Manual(M6)

drv a factor with levels 4(four wheel drive), F(front wheel drive), and R(rear wheel drive)

cty city mpg

hwy highway mpg

cmb combined city and highway mpg

Source

EPA data.

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
summary(Epaminicompact$cty)
plot(hwy ~ cty, data = Epaminicompact)
```

Epatwoseater

Fuel efficiency ratings for two-seater vehicles in 2001

Description

Data for Exercise 5.8

Usage

Epatwoseater

Format

A data frame/tibble with 36 observations on ten variables

class a character variable with value TWO SEATERS

manufacturer a character variable with values ACURA, AUDI, BMW, CHEVROLET, DODGE, FERRARI, HONDA, LAMBORGHINI, MAZDA, MERCEDES-BENZ, PLYMOUTH, PORSCHE, and TOYOTA

carline a character variable with values BOXSTER, BOXSTER S, CORVETTE, DB132/144 DIABLO, FERRARI 360 MODENA/SPIDER, FERRARI 550 MARANELLO/BARCHETTA, INSIGHT, MR2 ,MX-5 MIATA, NSX, PROWLER, S2000, SL500, SL600, SLK230 KOMPRESSOR, SLK320, TT ROADSTER, TT ROADSTER QUATTRO, VIPER CONVERTIBLE, VIPER COUPE, Z3 COUPE, Z3 ROADSTER, and Z8

displ engine displacement (in liters)

cyl number of cylinders

trans a factor with levels Auto(L4), Auto(L5), Auto(S4), Auto(S5), Auto(S6), Manual(M5), and Manual(M6)

drv a factor with levels 4(four wheel drive) F(front wheel drive) R(rear wheel drive)

cty city mpg

hwy highway mpg

cmb combined city and highway mpg

@source Environmental Protection Agency.

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
summary(Epatwoseater$cty)
plot(hwy ~ cty, data = Epatwoseater)
boxplot(cty ~ drv, data = Epatwoseater, col = "lightgreen")
```

Executiv	<i>Ages of 25 executives</i>
----------	------------------------------

Description

Data for Exercise 1.104

Usage

Executiv

Format

A data frame/tibble with 25 observations on one variable

age a numeric vector

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
hist(Executiv$age, xlab = "Age of banking executives",
     breaks = 5, main = "", col = "gray")
```

Exercise	<i>Weight loss for 30 members of an exercise program</i>
----------	--

Description

Data for Exercise 1.44

Usage

Exercise

Format

A data frame/tibble with 30 observations on one variable

loss a numeric vector

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
stem(Exercise$loss)
```

Fabric	<i>Measures of softness of ten different clothing garments washed with and without a softener</i>
--------	---

Description

Data for Example 7.21

Usage

Fabric

Format

A data frame/tibble with 20 observations on three variables

garment a numeric vector

softner a character variable with values with and without

softness a numeric vector

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
## Not run:
library(tidyr)
tidyr::spread(Fabric, softner, softness) -> FabricWide
wilcox.test(Pair(with, without)~1, alternative = "greater", data = FabricWide)
T7 <- tidyr::spread(Fabric, softner, softness) %>%
mutate(di = with - without, adi = abs(di), rk = rank(adi),
       srk = sign(di)*rk)
T7
t.test(T7$srk, alternative = "greater")

## End(Not run)
```

Faithful

Waiting times between successive eruptions of the Old Faithful geyser

Description

Data for Exercise 5.12 and 5.111

Usage

Faithful

Format

A data frame/tibble with 299 observations on two variables

time a numeric vector

eruption a factor with levels 1 and 2

Source

A. Azzalini and A. Bowman, "A Look at Some Data on the Old Faithful Geyser," *Journal of the Royal Statistical Society, Series C*, 39 (1990), 357-366.

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
t.test(time ~ eruption, data = Faithful)
hist(Faithful$time, xlab = "wait time", main = "", freq = FALSE)
lines(density(Faithful$time))

## Not run:
library(ggplot2)
ggplot2::ggplot(data = Faithful, aes(x = time, y = ..density..)) +
  geom_histogram(binwidth = 5, fill = "pink", col = "black") +
  geom_density() +
  theme_bw() +
  labs(x = "wait time")

## End(Not run)
```

Family

Size of family versus cost per person per week for groceries

Description

Data for Exercise 2.89

Usage

Family

Format

A data frame/tibble with 20 observations on two variables

number number in family

cost cost per person (in dollars)

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
plot(cost ~ number, data = Family)
abline(lm(cost ~ number, data = Family), col = "red")
cor(Family$cost, Family$number)

## Not run:
library(ggplot2)
ggplot2::ggplot(data = Family, aes(x = number, y = cost)) +
  geom_point() +
  geom_smooth(method = "lm") +
```

```
theme_bw()

## End(Not run)
```

Ferraro1	<i>Choice of presidential ticket in 1984 by gender</i>
----------	--

Description

Data for Exercise 8.23

Usage

Ferraro1

Format

A data frame/tibble with 1000 observations on two variables

gender a factor with levels Men and Women

candidate a character vector of 1984 president and vice-president candidates

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
T1 <- xtabs(~gender + candidate, data = Ferraro1)
T1
chisq.test(T1)
rm(T1)
```

Ferraro2	<i>Choice of vice presidential candidate in 1984 by gender</i>
----------	--

Description

Data for Exercise 8.23

Usage

Ferraro2

Format

A data frame/tibble with 1000 observations on two variables

gender a factor with levels Men and Women

candidate a character vector of 1984 president and vice-president candidates

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
T1 <- xtabs(~gender + candidate, data = Ferraro2)
T1
chisq.test(T1)
rm(T1)
```

Fertility

Fertility rates of all 50 states and DC

Description

Data for Exercise 1.125

Usage

Fertility

Format

A data frame/tibble with 51 observations on two variables

state a character variable with values Alabama, Alaska, Arizona, Arkansas, California, Colorado, Connecticut, Delaware, District of Columbia, Florida, Georgia, Hawaii, Idaho, Illinois, Indiana, Iowa, Kansas, Kentucky, Louisiana, Maine, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Montana, Nebraska, Nevada, New Hampshire, New Jersey, New Mexico, New York, North Carolina, North Dakota, Ohio, Oklahoma, Oregon, Pennsylvania, Rhode Island, South Carolina, South Dakota, Tennessee, Texas, Utah, Vermont, Virginia, Washington, West Virginia, Wisconsin, and Wyoming

rate fertility rate (expected number of births during childbearing years)

Source

Population Reference Bureau.

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
stem(Fertility$rate)
fivenum(Fertility$rate)
EDA(Fertility$rate)
```

Firstchi

Ages of women at the birth of their first child

Description

Data for Exercise 5.11

Usage

Firstchi

Format

A data frame/tibble with 87 observations on one variable

age age of woman at birth of her first child

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
EDA(Firstchi$age)
```

Fish

Length and number of fish caught with small and large mesh codend

Description

Data for Exercises 5.83, 5.119, and 7.29

Usage

Fish

Format

A data frame/tibble with 1534 observations on two variables

codend a character variable with values smallmesh and largemesh

length length of the fish measured in centimeters

Source

R. Millar, “Estimating the Size - Selectivity of Fishing Gear by Conditioning on the Total Catch,” *Journal of the American Statistical Association*, 87 (1992), 962 - 968.

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
tapply(Fish$length, Fish$codend, median, na.rm = TRUE)
SIGN.test(Fish$length[Fish$codend == "smallmesh"], conf.level = 0.99)
## Not run:
dplyr::group_by(Fish, codend) %>%
  summarize(MEDIAN = median(length, na.rm = TRUE))

## End(Not run)
```

Fitness

Number of sit-ups before and after a physical fitness course

Description

Data for Exercise 7.71

Usage

Fitness

Format

A data frame/tibble with 18 observations on the three variables

subject a character variable indicating subject number

test a character variable with values After and Before

number a numeric vector recording the number of sit-ups performed in one minute

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
## Not run:
tidyr::spread(Fitness, test, number) -> FitnessWide
t.test(Pair(After, Before)~1, alternative = "greater", data = FitnessWide)

Wide <- tidyr::spread(Fitness, test, number) %>%
mutate(diff = After - Before)
Wide
qqnorm(Wide$diff)
qqline(Wide$diff)
t.test(Wide$diff, alternative = "greater")

## End(Not run)
```

Florida2000

Florida voter results in the 2000 presidential election

Description

Data for Statistical Insight Chapter 2

Usage

Florida2000

Format

A data frame/tibble with 67 observations on 12 variables

county a character variable with values ALACHUA, BAKER, BAY, BRADFORD, BREVARD, BROWARD, CALHOUN, CHARLOTTE, CITRUS, CLAY, COLLIER, COLUMBIA, DADE, DE SOTO, DIXIE, DUVAL, ESCAMBIA, FLAGLER, FRANKLIN, GADSDEN, GILCHRIST, GLADES, GULF, HAMILTON, HARDEE, HENDRY, HERNANDO, HIGHLANDS, HILLSBOROUGH, HOLMES, INDIAN RIVER, JACKSON, JEFFERSON, LAFAYETTE, LAKE, LEE, LEON, LEVY, LIBERTY, MADISON, MANATEE, MARION, MARTIN, MONROE, NASSAU, OKALOOSA, OKEECHOBEE, ORANGE, OSCEOLA, PALM BEACH, PASCO, PINELLAS, POLK, PUTNAM, SANTA ROSA, SARASOTA, SEMINOLE, ST. JOHNS, ST. LUCIE, SUMTER, SUWANNEE, TAYLOR, UNION, VOLUSIA, WAKULLA, WALTON, and WASHINGTON

gore number of votes

bush number of votes

buchanan number of votes

nader number of votes

browne number of votes

hagelin number of votes

harris number of votes

mcreynolds number of votes

moorehead number of votes

phillips number of votes

total number of votes

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
plot(buchanan ~ total, data = Florida2000,
     xlab = "Total votes cast (in thousands)",
     ylab = "Votes for Buchanan")
```

Fluid	<i>Breakdown times of an insulating fluid under various levels of voltage stress</i>
-------	--

Description

Data for Exercise 5.76

Usage

Fluid

Format

A data frame/tibble with 76 observations on two variables

kilovolts a character variable showing kilowats

time breakdown time (in minutes)

Source

E. Soofi, N. Ebrahimi, and M. Habibullah, 1995.

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
DF1 <- Fluid[Fluid$kilovolts == "34kV", ]
DF1
# OR
DF2 <- subset(Fluid, subset = kilovolts == "34kV")
DF2
stem(DF2$time)
SIGN.test(DF2$time)
## Not run:
library(dplyr)
DF3 <- dplyr::filter(Fluid, kilovolts == "34kV")
DF3

## End(Not run)
```

Food

Annual food expenditures for 40 single households in Ohio

Description

Data for Exercise 5.106

Usage

Food

Format

A data frame/tibble with 40 observations on one variable

expenditure a numeric vector recording annual food expenditure (in dollars) in the state of Ohio.

Source

Bureau of Labor Statistics.

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
EDA(Food$expenditure)
```

Framingh

Cholesterol values of 62 subjects in the Framingham Heart Study

Description

Data for Exercises 1.56, 1.75, 3.69, and 5.60

Usage

Framingh

Format

A data frame/tibble with 62 observations on one variable

cholest a numeric vector with cholesterol values

Source

R. D'Agostino, et al., (1990) "A Suggestion for Using Powerful and Informative Tests for Normality," *The American Statistician*, 44 316-321.

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
stem(Framingh$cholest)
boxplot(Framingh$cholest, horizontal = TRUE)
hist(Framingh$cholest, freq = FALSE)
lines(density(Framingh$cholest))
mean(Framingh$cholest > 200 & Framingh$cholest < 240)

## Not run:
library(ggplot2)
ggplot2::ggplot(data = Framingh, aes(x = factor(1), y = cholest)) +
  geom_boxplot() +                # boxplot
  labs(x = "") +                  # no x label
  theme_bw() +                    # black and white theme
  geom_jitter(width = 0.2) +      # jitter points
  coord_flip()                    # Create horizontal plot
ggplot2::ggplot(data = Framingh, aes(x = cholest, y = ..density..)) +
  geom_histogram(fill = "pink", binwidth = 15, color = "black") +
  geom_density() +
  theme_bw()

## End(Not run)
```

Freshman

Ages of a random sample of 30 college freshmen

Description

Data for Exercise 6.53

Usage

Freshman

Format

A data frame/tibble with 30 observations on one variable

age a numeric vector of ages

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
SIGN.test(Freshman$age, md = 19)
```

Funeral

Cost of funeral by region of country

Description

Data for Exercise 8.54

Usage

Funeral

Format

A data frame/tibble with 400 observations on two variables

region a factor with levels Central, East, South, and West

cost a factor with levels less than expected, about what expected, and more than expected

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
T1 <- xtabs(~region + cost, data = Funeral)
T1
chisq.test(T1)
rm(T1)
```

Galaxie

Velocities of 82 galaxies in the Corona Borealis region

Description

Data for Example 5.2

Usage

Galaxie

Format

A data frame/tibble with 82 observations on one variable

velocity velocity measured in kilometers per second

Source

K. Roeder, "Density Estimation with Confidence Sets Explained by Superclusters and Voids in the Galaxies," *Journal of the American Statistical Association*, 85 (1990), 617-624.

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
EDA(Galaxie$velocity)
```

Gallup

Results of a Gallup poll on possession of marijuana as a criminal offense conducted in 1980

Description

Data for Exercise 2.76

Usage

Gallup

Format

A data frame/tibble with 1,200 observations on two variables

demographics a factor with levels National, Gender: Male Gender: Female, Education: College, Education: High School, Education: Grade School, Age: 18–24, Age: 25–29, Age: 30–49, Age: 50–older, Religion: Protestant, and Religion: Catholic

opinion a factor with levels Criminal, Not Criminal, and No Opinion

Source

George H. Gallup *The Gallup Opinion Index Report No. 179* (Princeton, NJ: The Gallup Poll, July 1980), p. 15.

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
T1 <- xtabs(~demographics + opinion, data = Gallup)
T1
t(T1[c(2, 3), ])
barplot(t(T1[c(2, 3), ]))
barplot(t(T1[c(2, 3), ]), beside = TRUE)

## Not run:
library(dplyr)
library(ggplot2)
dplyr::filter(Gallup, demographics == "Gender: Male" | demographics == "Gender: Female") %>%
ggplot2::ggplot(aes(x = demographics, fill = opinion)) +
  geom_bar() +
  theme_bw() +
  labs(y = "Fraction")

## End(Not run)
```

Gasoline

Price of regular unleaded gasoline obtained from 25 service stations

Description

Data for Exercise 1.45

Usage

Gasoline

Format

A data frame/tibble with 25 observations on one variable

price price for one gallon of gasoline

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
stem(Gasoline$price)

## Not run:
library(ggplot2)
ggplot2::ggplot(data = Gasoline, aes(x = factor(1), y = price)) +
  geom_violin() +
  geom_jitter() +
  theme_bw()

## End(Not run)
```

German

Number of errors in copying a German passage before and after an experimental course in German

Description

Data for Exercise 7.60

Usage

German

Format

A data frame/tibble with ten observations on three variables

student a character variable indicating student number

when a character variable with values Before and After to indicate when the student received experimental instruction in German

errors the number of errors in copying a German passage

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
## Not run:
tidyr::spread(German, when, errors) -> GermanWide
t.test(Pair(After, Before) ~ 1, data = GermanWide)
wilcox.test(Pair(After, Before) ~ 1, data = GermanWide)
T8 <- tidyr::spread(German, when, errors) %>%
mutate(di = After - Before, adi = abs(di), rk = rank(adi), srk = sign(di)*rk)
T8
qqnorm(T8$di)
qqline(T8$di)
t.test(T8$srk)

## End(Not run)
```

Golf

Distances a golf ball can be driven by 20 professional golfers

Description

Data for Exercise 5.24

Usage

Golf

Format

A data frame/tibble with 20 observations on one variable

yards distance a golf ball is driven in yards

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
stem(Golf$yards)
qqnorm(Golf$yards)
qqline(Golf$yards)

## Not run:
library(ggplot2)
ggplot2::ggplot(data = Golf, aes(sample = yards)) +
  geom_qq() +
  theme_bw()

## End(Not run)
```

Governor

*Annual salaries for state governors in 1994 and 1999***Description**

Data for Exercise 5.112

Usage

Governor

Format

A data frame/tibble with 50 observations on three variables

state a character variable with values Alabama, Alaska, Arizona, Arkansas, California, Colorado, Connecticut, Delaware, Florida, Georgia, Hawaii, Idaho, Illinois, Indiana, Iowa, Kansas, Kentucky, Louisiana, Maine, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Montana, Nebraska, Nevada, New Hampshire, New Jersey, New Mexico, New York, North Carolina, North Dakota, Ohio, Oklahoma, Oregon, Pennsylvania, Rhode Island, South Carolina, South Dakota, Tennessee, Texas, Utah, Vermont, Virginia, Washington, West Virginia, Wisconsin, and Wyoming

year a factor indicating year

salary a numeric vector with the governor's salary (in dollars)

Source*The 2000 World Almanac and Book of Facts.***References**

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
boxplot(salary ~ year, data = Governor)

## Not run:
library(ggplot2)
ggplot2::ggplot(data = Governor, aes(x = salary)) +
  geom_density(fill = "pink") +
  facet_grid(year ~ .) +
  theme_bw()

## End(Not run)
```

Gpa

*High school GPA versus college GPA***Description**

Data for Example 2.13

Usage

Gpa

Format

A data frame/tibble with 10 observations on two variables

hsgpa high school gpa**collgpa** college gpa**References**

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
plot(collgpa ~ hsgpa, data = Gpa)
mod <- lm(collgpa ~ hsgpa, data = Gpa)
abline(mod)           # add line
yhat <- predict(mod)   # fitted values
e <- resid(mod)        # residuals
cbind(Gpa, yhat, e)    # Table 2.1
cor(Gpa$hsgpa, Gpa$collgpa)

## Not run:
library(ggplot2)
ggplot2::ggplot(data = Gpa, aes(x = hsgpa, y = collgpa)) +
  geom_point() +
  geom_smooth(method = "lm") +
  theme_bw()

## End(Not run)
```

Grades

*Test grades in a beginning statistics class***Description**

Data for Exercise 1.120

Usage

Grades

Format

A data frame with 29 observations on one variable

grades a numeric vector containing test grades**References**

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
hist(Grades$grades, main = "", xlab = "Test grades", right = FALSE)

## Not run:
library(ggplot2)
ggplot2::ggplot(data = Grades, aes(x = grades, y = ..density..)) +
  geom_histogram(fill = "pink", binwidth = 5, color = "black") +
  geom_density(lwd = 2, color = "red") +
  theme_bw()

## End(Not run)
```

Graduate

*Graduation rates for student athletes in the Southeastern Conf.***Description**

Data for Exercise 1.118

Usage

Graduate

Format

A data frame/tibble with 12 observations on three variables

school a character variable with values Alabama, Arkansas, Auburn, Florida, Georgia, Kentucky, Louisiana St, Mississippi, Mississippi St, South Carolina, Tennessee, and Vanderbilt

code a character variable with values Al, Ar, Au Fl, Ge, Ke, LSt, Mi, MSt, SC, Te, and Va

percent graduation rate

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
barplot(Graduate$percent, names.arg = Graduate$school,
        las = 2, cex.names = 0.7, col = "tomato")
```

Greenriv

Varve thickness from a sequence through an Eocene lake deposit in the Rocky Mountains

Description

Data for Exercise 6.57

Usage

Greenriv

Format

A data frame/tibble with 37 observations on one variable

thick varve thickness in millimeters

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
stem(Greenriv$thick)
SIGN.test(Greenriv$thick, md = 7.3, alternative = "greater")
```

Grnriv2	<i>Thickness of a varved section of the Green river oil shale deposit near a major lake in the Rocky Mountains</i>
---------	--

Description

Data for Exercises 6.45 and 6.98

Usage

Grnriv2

Format

A data frame/tibble with 101 observations on one variable

thick varve thickness (in millimeters)

Source

J. Davis, *Statistics and Data Analysis in Geology*, 2nd Ed., Jon Wiley and Sons, New York.

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
stem(Grnriv2$thick)
t.test(Grnriv2$thick, mu = 8, alternative = "less")
```

Groupabc	<i>Group data to illustrate analysis of variance</i>
----------	--

Description

Data for Exercise 10.42

Usage

Groupabc

Format

A data frame/tibble with 45 observations on two variables

group a factor with levels A, B, and C

response a numeric vector

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
boxplot(response ~ group, data = Groupabc,
        col = c("red", "blue", "green"))
anova(lm(response ~ group, data = Groupabc))
```

Groups

An illustration of analysis of variance

Description

Data for Exercise 10.4

Usage

Groups

Format

A data frame/tibble with 78 observations on two variables

group a factor with levels A, B, and C

response a numeric vector

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
boxplot(response ~ group, data = Groups, col = c("red", "blue", "green"))
anova(lm(response ~ group, data = Groups))
```

Gym*Children's age versus number of completed gymnastic activities*

Description

Data for Exercises 2.21 and 9.14

Usage

Gym

Format

A data frame/tibble with eight observations on three variables

age age of child

number number of gymnastic activities successfully completed

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
plot(number ~ age, data = Gym)
model <- lm(number ~ age, data = Gym)
abline(model, col = "red")
summary(model)
```

Habits*Study habits of students in two matched school districts*

Description

Data for Exercise 7.57

Usage

Habits

Format

A data frame/tibble with 11 observations on four variables

A study habit score

B study habit score

differ B minus A

signrks the signed-ranked-differences

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
shapiro.test(Habits$differ)
qqnorm(Habits$differ)
qqline(Habits$differ)
wilcox.test(Pair(B, A) ~ 1, data = Habits, alternative = "less")
t.test(Habits$signrks, alternative = "less")

## Not run:
library(ggplot2)
ggplot2::ggplot(data = Habits, aes(x = differ)) +
  geom_dotplot(fill = "blue") +
  theme_bw()

## End(Not run)
```

Haptoglo

Haptoglobin concentration in blood serum of 8 healthy adults

Description

Data for Example 6.9

Usage

Haptoglo

Format

A data frame/tibble with eight observations on one variable

concent haptoglobin concentration (in grams per liter)

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
shapiro.test(Haptoglo$concent)
t.test(Haptoglo$concent, mu = 2, alternative = "less")
```

Hardware

Daily receipts for a small hardware store for 31 working days

Description

Daily receipts for a small hardware store for 31 working days

Usage

Hardware

Format

A data frame with 31 observations on one variable

receipt a numeric vector of daily receipts (in dollars)

Source

J.C. Miller and J.N. Miller, (1988), *Statistics for Analytical Chemistry*, 2nd Ed. (New York: Halsted Press).

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
stem(Hardware$receipt)
```

Hardwood	<i>Tensile strength of Kraft paper for different percentages of hardwood in the batches of pulp</i>
----------	---

Description

Data for Example 2.18 and Exercise 9.34

Usage

Hardwood

Format

A data frame/tibble with 19 observations on two variables

tensile tensile strength of kraft paper (in pounds per square inch)

hardwood percent of hardwood in the batch of pulp that was used to produce the paper

Source

G. Joglekar, et al., "Lack-of-Fit Testing When Replicates Are Not Available," *The American Statistician*, 43(3), (1989), 135-143.

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
plot(tensile ~ hardwood, data = Hardwood)
model <- lm(tensile ~ hardwood, data = Hardwood)
abline(model, col = "red")
plot(model, which = 1)
```

Heat	<i>Primary heating sources of homes on indian reservations versus all households</i>
------	--

Description

Data for Exercise 1.29

Usage

Heat

Format

A data frame/tibble with 301 observations on two variables

fuel a factor with levels Utility gas, LP bottled gas, Electricity, Fuel oil, Wood, and Other

location a factor with levels American Indians on reservation, All U.S. households, and American Indians not on reservations

Source

Bureau of the Census, *Housing of the American Indians on Reservations*, Statistical Brief 95-11, April 1995.

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
T1 <- xtabs(~ fuel + location, data = Heat)
T1
barplot(t(T1), beside = TRUE, legend = TRUE)

## Not run:
library(ggplot2)
ggplot2::ggplot(data = Heat, aes(x = fuel, fill = location)) +
  geom_bar(position = "dodge") +
  labs(y = "percent") +
  theme_bw() +
  theme(axis.text.x = element_text(angle = 30, hjust = 1))

## End(Not run)
```

Heating

Fuel efficiency ratings for three types of oil heaters

Description

Data for Exercise 10.32

Usage

Heating

Format

A data frame/tibble with 90 observations on the two variables

type a factor with levels A, B, and C denoting the type of oil heater

efficiency heater efficiency rating

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
boxplot(efficiency ~ type, data = Heating,
        col = c("red", "blue", "green"))
kruskal.test(efficiency ~ type, data = Heating)
```

Hodgkin

Results of treatments for Hodgkin's disease

Description

Data for Exercise 2.77

Usage

Hodgkin

Format

A data frame/tibble with 538 observations on two variables

type a factor with levels LD, LP, MC, and NS

response a factor with levels Positive, Partial, and None

Source

I. Dunsmore, F. Daly, *Statistical Methods, Unit 9, Categorical Data*, Milton Keynes, The Open University, 18.

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
T1 <- xtabs(~type + response, data = Hodgkin)
T1
barplot(t(T1), legend = TRUE, beside = TRUE)

## Not run:
library(ggplot2)
ggplot2::ggplot(data = Hodgkin, aes(x = type, fill = response)) +
  geom_bar(position = "dodge") +
  theme_bw()

## End(Not run)
```

Homes	<i>Median prices of single-family homes in 65 metropolitan statistical areas</i>
-------	--

Description

Data for Statistical Insight Chapter 5

Usage

Homes

Format

A data frame/tibble with 65 observations on the four variables

city a character variable with values Akron OH, Albuquerque NM, Anaheim CA, Atlanta GA, Baltimore MD, Baton Rouge LA, Birmingham AL, Boston MA, Bradenton FL, Buffalo NY, Charleston SC, Chicago IL, Cincinnati OH, Cleveland OH, Columbia SC, Columbus OH, Corpus Christi TX, Dallas TX, Daytona Beach FL, Denver CO, Des Moines IA, Detroit MI, El Paso TX, Grand Rapids MI, Hartford CT, Honolulu HI, Houston TX, Indianapolis IN, Jacksonville FL, Kansas City MO, Knoxville TN, Las Vegas NV, Los Angeles CA, Louisville KY, Madison WI, Memphis TN, Miami FL, Milwaukee WI, Minneapolis MN, Mobile AL, Nashville TN, New Haven CT, New Orleans LA, New York NY, Oklahoma City OK, Omaha NE, Orlando FL, Philadelphia PA, Phoenix AZ, Pittsburgh PA, Portland OR, Providence RI, Sacramento CA, Salt Lake

City UT, San Antonio TX, San Diego CA, San Francisco CA, Seattle WA, Spokane WA, St Louis MO, Syracuse NY, Tampa FL, Toledo OH, Tulsa OK, and Washington DC

region a character variable with values Midwest, Northeast, South, and West

year a factor with levels 1994 and 2000

price median house price (in dollars)

Source

National Association of Realtors.

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
tapply(Homes$price, Homes$year, mean)
tapply(Homes$price, Homes$region, mean)
p2000 <- subset(Homes, year == "2000")
p1994 <- subset(Homes, year == "1994")
## Not run:
library(dplyr)
library(ggplot2)
dplyr::group_by(Homes, year, region) %>%
  summarize(AvgPrice = mean(price))
ggplot2::ggplot(data = Homes, aes(x = region, y = price)) +
  geom_boxplot() +
  theme_bw() +
  facet_grid(year ~ .)

## End(Not run)
```

Homework

Number of hours per week spent on homework for private and public high school students

Description

Data for Exercise 7.78

Usage

Homework

Format

A data frame with 30 observations on two variables

school type of school either private or public

time number of hours per week spent on homework

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
boxplot(time ~ school, data = Homework,
        ylab = "Hours per week spent on homework")
#
t.test(time ~ school, data = Homework)
```

Honda

Miles per gallon for a Honda Civic on 35 different occasions

Description

Data for Statistical Insight Chapter 6

Usage

Honda

Format

A data frame/tibble with 35 observations on one variable

mileage miles per gallon for a Honda Civic

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
t.test(Honda$mileage, mu = 40, alternative = "less")
```

Hostile	<i>Hostility levels of high school students from rural, suburban, and urban areas</i>
---------	---

Description

Data for Example 10.6

Usage

Hostile

Format

A data frame/tibble with 135 observations on two variables

location a factor with the location of the high school student (Rural, Suburban, or Urban)

hostility the score from the Hostility Level Test

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
boxplot(hostility ~ location, data = Hostile,  
        col = c("red", "blue", "green"))  
kruskal.test(hostility ~ location, data = Hostile)
```

Housing	<i>Median home prices for 1984 and 1993 in 37 markets across the U.S.</i>
---------	---

Description

Data for Exercise 5.82

Usage

Housing

Format

A data frame/tibble with 74 observations on three variables

city a character variable with values Albany, Anaheim, Atlanta, Baltimore, Birmingham, Boston, Chicago, Cincinnati, Cleveland, Columbus, Dallas, Denver, Detroit, Ft Lauderdale, Houston, Indianapolis, Kansas City, Los Angeles, Louisville, Memphis, Miami, Milwaukee, Minneapolis, Nashville, New York, Oklahoma City, Philadelphia, Providence, Rochester, Salt Lake City, San Antonio, San Diego, San Francisco, San Jose, St Louis, Tampa, and Washington

year a factor with levels 1984 and 1993

price median house price (in dollars)

Source

National Association of Realtors.

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
stripchart(price ~ year, data = Housing, method = "stack",
           pch = 1, col = c("red", "blue"))
## Not run:
library(ggplot2)
ggplot2::ggplot(data = Housing, aes(x = price, fill = year)) +
  geom_dotplot() +
  facet_grid(year ~ .) +
  theme_bw()

## End(Not run)
```

Hurrican

Number of storms, hurricanes and El Nino effects from 1950 through 1995

Description

Data for Exercises 1.38, 10.19, and Example 1.6

Usage

Hurrican

Format

A data frame/tibble with 46 observations on four variables

year a numeric vector indicating year

storms a numeric vector recording number of storms

hurricane a numeric vector recording number of hurricanes

elnino a factor with levels cold, neutral, and warm

Source

National Hurricane Center.

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
T1 <- xtabs(~hurricane, data = Hurricane)
T1
barplot(T1, col = "blue", main = "Problem 1.38",
        xlab = "Number of hurricanes",
        ylab = "Number of seasons")
boxplot(storms ~ elnino, data = Hurricane,
        col = c("blue", "yellow", "red"))
anova(lm(storms ~ elnino, data = Hurricane))
rm(T1)
```

Iceberg

Number of icebergs sighted each month south of Newfoundland and south of the Grand Banks in 1920

Description

Data for Exercise 2.46 and 2.60

Usage

Iceberg

Format

A data frame with 12 observations on three variables

month a character variable with abbreviated months of the year

Newfoundland number of icebergs sighted south of Newfoundland

Grand Banks number of icebergs sighted south of Grand Banks

Source

N. Shaw, *Manual of Meteorology*, Vol. 2 (London: Cambridge University Press 1942), 7; and F. Mosteller and J. Tukey, *Data Analysis and Regression* (Reading, MA: Addison - Wesley, 1977).

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
plot(Newfoundland ~ `Grand Banks`, data = Iceberg)
abline(lm(Newfoundland ~ `Grand Banks`, data = Iceberg), col = "blue")
```

Income	<i>Percent change in personal income from 1st to 2nd quarter in 2000</i>
--------	--

Description

Data for Exercise 1.33

Usage

Income

Format

A data frame/tibble with 51 observations on two variables

state a character variable with values Alabama, Alaska, Arizona, Arkansas, California, Colorado, Connecticut, Delaware, District of Columbia, Florida, Georgia, Hawaii, Idaho, Illinois, Indiana, Iowa, Kansas, Kentucky, Louisiana, Maine, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Montana, Nebraska, Nevada, New Hampshire, New Jersey, New Mexico, New York, North Carolina, North Dakota, Ohio, Oklahoma, Oregon, Pennsylvania, Rhode Island, South Carolina, South Dakota, Tennessee, Texas, Utah, Vermont, Virginia, Washington, West Virginia, Wisconsin, and Wyoming

percent_change percent change in income from first quarter to the second quarter of 2000

Source

US Department of Commerce.

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
Income$class <- cut(Income$percent_change,
                    breaks = c(-Inf, 0.5, 1.0, 1.5, 2.0, Inf))
T1 <- xtabs(~class, data = Income)
T1
barplot(T1, col = "pink")
## Not run:
library(ggplot2)
DF <- as.data.frame(T1)
DF
ggplot2::ggplot(data = DF, aes(x = class, y = Freq)) +
  geom_bar(stat = "identity", fill = "purple") +
  theme_bw()

## End(Not run)
```

Independent

Illustrates a comparison problem for long-tailed distributions

Description

Data for Exercise 7.41

Usage

Independent

Format

A data frame/tibble with 46 observations on two variables

score a numeric vector

group a factor with levels A and B

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
qqnorm(Independent$score[Independent$group=="A"])
qqline(Independent$score[Independent$group=="A"])
qqnorm(Independent$score[Independent$group=="B"])
qqline(Independent$score[Independent$group=="B"])
boxplot(score ~ group, data = Independent, col = "blue")
wilcox.test(score ~ group, data = Independent)
```

Indian	<i>Educational attainment versus per capita income and poverty rate for American Indians living on reservations</i>
--------	---

Description

Data for Exercise 2.95

Usage

Indian

Format

A data frame/tibble with ten observations on four variables

reservation a character variable with values Blackfeet, Fort Apache, Gila River, Hopi, Navajo, Papago, Pine Ridge, Rosebud, San Carlos, and Zuni Pueblo

percent high school percent who have graduated from high school

per capita income per capita income (in dollars)

poverty rate percent poverty

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
par(mfrow = c(1, 2))
plot(`per capita income` ~ `percent high school`, data = Indian,
     xlab = "Percent high school graduates", ylab = "Per capita income")
plot(`poverty rate` ~ `percent high school`, data = Indian,
     xlab = "Percent high school graduates", ylab = "Percent poverty")
par(mfrow = c(1, 1))
```

Indiapol

Average miles per hour for the winners of the Indianapolis 500 race

Description

Data for Exercise 1.128

Usage

Indiapol

Format

A data frame/tibble with 39 observations on two variables

year the year of the race

speed the winners average speed (in mph)

Source

The World Almanac and Book of Facts, 2000, p. 1004.

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
plot(speed ~ year, data = Indiapol, type = "b")
```

Indy500

Qualifying miles per hour and number of previous starts for drivers in 79th Indianapolis 500 race

Description

Data for Exercises 7.11 and 7.36

Usage

Indy500

Format

A data frame/tibble with 33 observations on four variables

driver a character variable with values andretti, bachelart, boesel, brayton, c.guerrero, cheever, fabi, fernandez, ferran, fittipaldi, fox, goodyear, gordon, gugelmin, herta, james, johansson, jones, lazier, luyendyk, matsuda, matsushita, pruet, r.guerrero, rahal, ribeiro, salazar, sharp, sullivan, tracy, vasser, villeneuve, and zampedri

qualif qualifying speed (in mph)

starts number of Indianapolis 500 starts

group a numeric vector where 1 indicates the driver has 4 or fewer Indianapolis 500 starts and a 2 for drivers with 5 or more Indianapolis 500 starts

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
stripchart(qualif ~ group, data = Indy500, method = "stack",
           pch = 19, col = c("red", "blue"))
boxplot(qualif ~ group, data = Indy500)
t.test(qualif ~ group, data = Indy500)
## Not run:
library(ggplot2)
ggplot2::ggplot(data = Indy500, aes(sample = qualif)) +
  geom_qq() +
  facet_grid(group ~ .) +
  theme_bw()

## End(Not run)
```

Inflatio

Private pay increase of salaried employees versus inflation rate

Description

Data for Exercises 2.12 and 2.29

Usage

Inflatio

Format

A data frame/tibble with 24 observations on four variables

year a numeric vector of years

pay average hourly wage for salaried employees (in dollars)

increase percent increase in hourly wage over previous year

inflation percent inflation rate

Source

Bureau of Labor Statistics.

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
plot(increase ~ inflation, data = Inflation)
cor(Inflation$increase, Inflation$inflation, use = "complete.obs")
```

Inletoil

Inlet oil temperature through a valve

Description

Data for Exercises 5.91 and 6.48

Usage

Inletoil

Format

A data frame/tibble with 12 observations on one variable

temp inlet oil temperature (Fahrenheit)

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
hist(Inletoil$temp, breaks = 3)
qqnorm(Inletoil$temp)
qqline(Inletoil$temp)
t.test(Inletoil$temp)
t.test(Inletoil$temp, mu = 98, alternative = "less")
```

Inmate	<i>Type of drug offense by race</i>
--------	-------------------------------------

Description

Data for Statistical Insight Chapter 8

Usage

Inmate

Format

A data frame/tibble with 28,047 observations on two variables

race a factor with levels white, black, and hispanic

drug a factor with levels heroin, crack, cocaine, and marijuana

Source

C. Wolf Harlow (1994), *Comparing Federal and State Prison Inmates*, NCJ-145864, U.S. Department of Justice, Bureau of Justice Statistics.

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
T1 <- xtabs(~race + drug, data = Inmate)
T1
chisq.test(T1)
rm(T1)
```

Inspect

*Percent of vehicles passing inspection by type inspection station***Description**

Data for Exercise 8.59

Usage

Inspect

Format

A data frame/tibble with 174 observations on two variables

station a factor with levels auto inspection, auto repair, car care center, gas station, new car dealer, and tire store

passed a factor with levels less than 70%, between 70% and 84%, and more than 85%

Source

The Charlotte Observer, December 13, 1992.

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
T1 <- xtabs(~ station + passed, data = Inspect)
T1
barplot(T1, beside = TRUE, legend = TRUE)
chisq.test(T1)
rm(T1)

## Not run:
library(ggplot2)
ggplot2::ggplot(data = Inspect, aes(x = passed, fill = station)) +
  geom_bar(position = "dodge") +
  theme_bw()

## End(Not run)
```

Insulate

Heat loss through a new insulating medium

Description

Data for Exercise 9.50

Usage

Insulate

Format

A data frame/tibble with ten observations on two variables

temp outside temperature (in degrees Celcius)

loss heat loss (in BTUs)

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
plot(loss ~ temp, data = Insulate)
model <- lm(loss ~ temp, data = Insulate)
abline(model, col = "blue")
summary(model)

## Not run:
library(ggplot2)
ggplot2::ggplot(data = Insulate, aes(x = temp, y = loss)) +
  geom_point() +
  geom_smooth(method = "lm", se = FALSE) +
  theme_bw()

## End(Not run)
```

Iqgpa	<i>GPA versus IQ for 12 individuals</i>
-------	---

Description

Data for Exercises 9.51 and 9.52

Usage

Iqgpa

Format

A data frame/tibble with 12 observations on two variables

iq IQ scores

gpa Grade point average

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
plot(gpa ~ iq, data = Iqgpa, col = "blue", pch = 19)
model <- lm(gpa ~ iq, data = Iqgpa)
summary(model)
rm(model)
```

Irises	<i>R.A. Fishers famous data on Irises</i>
--------	---

Description

Data for Examples 1.15 and 5.19

Usage

Irises

Format

A data frame/tibble with 150 observations on five variables

sepal_length sepal length (in cm)

sepal_width sepal width (in cm)

petal_length petal length (in cm)

petal_width petal width (in cm)

species a factor with levels *setosa*, *versicolor*, and *virginica*

Source

Fisher, R. A. (1936) The use of multiple measurements in taxonomic problems. *Annals of Eugenics*, 7, Part II, 179-188.

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
tapply(Iris$sepal_length, Iris$species, mean)
t.test(Iris$sepal_length[Iris$species == "setosa"], conf.level = 0.99)
hist(Iris$sepal_length[Iris$species == "setosa"],
     main = "Sepal length for\n Iris Setosa",
     xlab = "Length (in cm)")
boxplot(sepal_length ~ species, data = Iris)
```

Jdpower

Number of problems reported per 100 cars in 1994 versus 1995s

Description

Data for Exercise 2.14, 2.17, 2.31, 2.33, and 2.40

Usage

Jdpower

Format

A data frame/tibble with 29 observations on three variables

car a factor with levels Acura, BMW, Buick, Cadillac, Chevrolet, Dodge Eagle, Ford, Geo, Honda, Hyundai, Infiniti, Jaguar, Lexus, Lincoln, Mazda, Mercedes-Benz, Mercury, Mitsubishi, Nissan, Oldsmobile, Plymouth, Pontiac, Saab, Saturn, and Subaru, Toyota Volkswagen, Volvo

1994 number of problems per 100 cars in 1994

1995 number of problems per 100 cars in 1995

Source

USA Today, May 25, 1995.

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
model <- lm(`1995` ~ `1994`, data = Jdpower)
summary(model)
plot(`1995` ~ `1994`, data = Jdpower)
abline(model, col = "red")
rm(model)
```

Jobsat	<i>Job satisfaction and stress level for 9 school teachers</i>
--------	--

Description

Data for Exercise 9.60

Usage

Jobsat

Format

A data frame/tibble with nine observations on two variables

wspt Wilson Stress Profile score for teachers

satisfaction job satisfaction score

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
plot(satisfaction ~ wspt, data = Jobsat)
model <- lm(satisfaction ~ wspt, data = Jobsat)
abline(model, col = "blue")
summary(model)
rm(model)
```

Kidsmoke

Smoking habits of boys and girls ages 12 to 18

Description

Data for Exercise 4.85

Usage

Kidsmoke

Format

A data frame/tibble with 1000 observations on two variables

gender character vector with values female and male

smoke a character vector with values no and yes

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
T1 <- xtabs(~smoke + gender, data = Kidsmoke)
T1
prop.table(T1)
prop.table(T1, 1)
prop.table(T1, 2)
```

Kilowatt	<i>Rates per kilowatt-hour for each of the 50 states and DC</i>
----------	---

Description

Data for Example 5.9

Usage

Kilowatt

Format

A data frame/tibble with 51 observations on two variables

state a factor with levels Alabama Alaska, Arizona, Arkansas California, Colorado, Connecticut, Delaware, District of Columbia, Florida,Georgia, Hawaii, Idaho, Illinois, Indiana, Iowa Kansas Kentucky, Louisiana, Maine, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Montana Nebraska, Nevada, New Hampshire, New Jersey, New Mexico, New York, North Carolina, North Dakota, Ohio, Oklahoma, Oregon, Pennsylvania, Rhode Island, South Carolina, South Dakota, Tennessee, Texas, Utah, Vermont, Virginia Washington, West Virginia, Wisconsin, and Wyoming

rate a numeric vector indicating rates for kilowatt per hour

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

EDA(Kilowatt\$rate)

Kinder	<i>Reading scores for first grade children who attended kindergarten versus those who did not</i>
--------	---

Description

Data for Exercise 7.68

Usage

Kinder

Format

A data frame/tibble with eight observations on three variables

pair a numeric indicator of pair

kinder reading score of kids who went to kindergarten

nokinder reading score of kids who did not go to kindergarten

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
boxplot(Kinder$kindergarten, Kinder$notkindergarten)
diff <- Kinder$kindergarten - Kinder$notkindergarten
qqnorm(diff)
qqline(diff)
shapiro.test(diff)
t.test(diff)
rm(diff)
```

Laminect

Median costs of laminectomies at hospitals across North Carolina in 1992

Description

Data for Exercise 10.18

Usage

Laminect

Format

A data frame/tibble with 138 observations on two variables

area a character vector indicating the area of the hospital with Rural, Regional, and Metropol

cost a numeric vector indicating cost of a laminectomy

Source

Consumer's Guide to Hospitalization Charges in North Carolina Hospitals (August 1994), North Carolina Medical Database Commission, Department of Insurance.

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
boxplot(cost ~ area, data = Laminect, col = topo.colors(3))
anova(lm(cost ~ area, data = Laminect))
```

Lead	<i>Lead levels in children's blood whose parents worked in a battery factory</i>
------	--

Description

Data for Example 1.17

Usage

Lead

Format

A data frame/tibble with 66 observations on the two variables

group a character vector with values exposed and control

lead a numeric vector indicating the level of lead in children's blood (in micrograms/dl)

Source

Morton, D. et al. (1982), "Lead Absorption in Children of Employees in a Lead-Related Industry," *American Journal of Epidemiology*, 155, 549-555.

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
boxplot(lead ~ group, data = Lead, col = topo.colors(2))
```

Leader

Leadership exam scores by age for employees on an industrial plant

Description

Data for Exercise 7.31

Usage

Leader

Format

A data frame/tibble with 34 observations on two variables

age a character vector indicating age with values under35 and over35

score score on a leadership exam

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
boxplot(score ~ age, data = Leader, col = c("gray", "green"))
t.test(score ~ age, data = Leader)
```

Lethal

Survival time of mice injected with an experimental lethal drug

Description

Data for Example 6.12

Usage

Lethal

Format

A data frame/tibble with 30 observations on one variable

survival a numeric vector indicating time survived after injection (in seconds)

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
SIGN.test(Lethal$survival, md = 45, alternative = "less")
```

Life	<i>Life expectancy of men and women in U.S.</i>
------	---

Description

Data for Exercise 1.31

Usage

Life

Format

A data frame/tibble with eight observations on three variables

year a numeric vector indicating year

men life expectancy for men (in years)

women life expectancy for women (in years)

Source

National Center for Health Statistics.

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
plot(men ~ year, type = "l", ylim = c(min(men, women), max(men, women)),
     col = "blue", main = "Life Expectancy vs Year", ylab = "Age",
     xlab = "Year", data = Life)
lines(women ~ year, col = "red", data = Life)
text(1955, 65, "Men", col = "blue")
text(1955, 70, "Women", col = "red")
```

Lifespan	<i>Life span of electronic components used in a spacecraft versus heat</i>
----------	--

Description

Data for Exercise 2.4, 2.37, and 2.49

Usage

Lifespan

Format

A data frame/tibble with six observations two variables

heat temperature (in Celcius)

life lifespan of component (in hours)

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
plot(life ~ heat, data = Lifespan)
model <- lm(life ~ heat, data = Lifespan)
abline(model, col = "red")
resid(model)
sum((resid(model))^2)
anova(model)
rm(model)
```

Ligntmonth	<i>Relationship between damage reports and deaths caused by lightning</i>
------------	---

Description

Data for Exercise 2.6

Usage

Ligntmonth

Format

A data frame/tibble with 12 observations on four variables

month a factor with levels 1/01/2000, 10/01/2000, 11/01/2000, 12/01/2000, 2/01/2000, 3/01/2000, 4/01/2000, 5/01/2000, 6/01/2000, 7/01/2000, 8/01/2000, and 9/01/2000

deaths number of deaths due to lightning strikes

injuries number of injuries due to lightning strikes

damage damage due to lightning strikes (in dollars)

Source

Lighting Fatalities, Injuries and Damage Reports in the United States, 1959-1994, NOAA Technical Memorandum NWS SR-193, Dept. of Commerce.

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
plot(deaths ~ damage, data = Lightmonth)
model = lm(deaths ~ damage, data = Lightmonth)
abline(model, col = "red")
rm(model)
```

Lodge

Measured traffic at three prospective locations for a motor lodge

Description

Data for Exercise 10.33

Usage

Lodge

Format

A data frame/tibble with 45 observations on six variables

traffic a numeric vector indicating the amount of vehicles that passed a site in 1 hour

site a numeric vector with values 1, 2, and 3

ranks ranks for variable traffic

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
boxplot(traffic ~ site, data = Lodge, col = cm.colors(3))
anova(lm(traffic ~ factor(site), data = Lodge))
```

Longtail

Long-tailed distributions to illustrate Kruskal Wallis test

Description

Data for Exercise 10.45

Usage

Longtail

Format

A data frame/tibble with 60 observations on three variables

score a numeric vector

group a numeric vector with values 1, 2, and 3

ranks ranks for variable score

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
boxplot(score ~ group, data = Longtail, col = heat.colors(3))
kruskal.test(score ~ factor(group), data = Longtail)
anova(lm(score ~ factor(group), data = Longtail))
```

Lowabil

Reading skills of 24 matched low ability students

Description

Data for Example 7.18

Usage

Lowabil

Format

A data frame/tibble with 12 observations on three variables

pair a numeric indicator of pair

experiment score of the child with the experimental method

control score of the child with the standard method

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
diff = Lowabil$experiment - Lowabil$control
qqnorm(diff)
qqline(diff)
shapiro.test(diff)
t.test(diff)
rm(diff)
```

Magnesiu

Magnesium concentration and distances between samples

Description

Data for Exercise 9.9

Usage

Magnesiu

Format

A data frame/tibble with 20 observations on two variables

distance distance between samples

magnesium concentration of magnesium

Source

Davis, J. (1986), *Statistics and Data Analysis in Geology*, 2d. Ed., John Wiley and Sons, New York, p. 146.

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
plot(magnesium ~ distance, data = Magnesium)
model = lm(magnesium ~ distance, data = Magnesium)
abline(model, col = "red")
summary(model)
rm(model)
```

Malpract

Amounts awarded in 17 malpractice cases

Description

Data for Exercise 5.73

Usage

Malpract

Format

A data frame/tibble with 17 observations on one variable

award malpractice reward (in \$1000)

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
SIGN.test(Malpract$award, conf.level = 0.90)
```

Manager	<i>Advertised salaries offered general managers of major corporations in 1995</i>
---------	---

Description

Data for Exercise 5.81

Usage

Manager

Format

A data frame/tibble with 26 observations on one variable

salary random sample of advertised annual salaries of top executives (in dollars)

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
stem(Manager$salary)
SIGN.test(Manager$salary)
```

Marked	<i>Percent of marked cars in 65 police departments in Florida</i>
--------	---

Description

Data for Exercise 6.100

Usage

Marked

Format

A data frame/tibble with 65 observations on one variable

percent percentage of marked cars in 65 Florida police departments

Source

Law Enforcement Management and Administrative Statistics, 1993, Bureau of Justice Statistics, NCJ-148825, September 1995, p. 147-148.

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
EDA(Marked$percent)
SIGN.test(Marked$percent, md = 60, alternative = "greater")
t.test(Marked$percent, mu = 60, alternative = "greater")
```

Math

Standardized math test scores for 30 students

Description

Data for Exercise 1.69

Usage

Math

Format

A data frame/tibble with 30 observations on one variable

score scores on a standardized test for 30 tenth graders

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```

stem(Math$score)
hist(Math$score, main = "Math Scores", xlab = "score", freq = FALSE)
lines(density(Math$score), col = "red")
CharlieZ <- (62 - mean(Math$score))/sd(Math$score)
CharlieZ
scale(Math$score)[which(Math$score == 62)]

```

Mathcomp

Standardized math competency for a group of entering freshmen at a small community college

Description

Data for Exercise 5.26

Usage

Mathcomp

Format

A data frame/tibble with 31 observations one variable

score scores of 31 entering freshmen at a community college on a national standardized test

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```

stem(Mathcomp$score)
EDA(Mathcomp$score)

```

Mathpro

Math proficiency and SAT scores by states

Description

Data for Exercise 9.24, Example 9.1, and Example 9.6

Usage

Mathpro

Format

A data frame/tibble with 51 observations on four variables

state a factor with levels Conn, D.C., Del, Ga, Hawaii, Ind, Maine, Mass, Md, N.C., N.H., N.J., N.Y., Ore, Pa, R.I., S.C., Va, and Vt

sat_math SAT math scores for high school seniors

profic math proficiency scores for eighth graders

group a numeric vector

Source

National Assessment of Educational Progress and The College Board.

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
model <- lm(sat_math ~ profic, data = Mathpro)
plot(sat_math ~ profic, data = Mathpro, ylab = "SAT", xlab = "proficiency")
abline(model, col = "red")
summary(model)
rm(model)
```

Maze	<i>Error scores for four groups of experimental animals running a maze</i>
------	--

Description

Data for Exercise 10.13

Usage

Maze

Format

A data frame/tibble with 32 observations on two variables

score error scores for animals running through a maze under different conditions

condition a factor with levels CondA, CondB, CondC, and CondD

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
boxplot(score ~ condition, data = Maze, col = rainbow(4))
anova(lm(score ~ condition, data = Maze))
```

Median	<i>Illustrates test of equality of medians with the Kruskal Wallis test</i>
--------	---

Description

Data for Exercise 10.52

Usage

Median

Format

A data frame/tibble with 45 observations on two variables

sample a vector with values Sample1, Sample 2, and Sample 3

value a numeric vector

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
boxplot(value ~ sample, data = Median, col = rainbow(3))
anova(lm(value ~ sample, data = Median))
kruskal.test(value ~ factor(sample), data = Median)
```

Mental	<i>Median mental ages of 16 girls</i>
--------	---------------------------------------

Description

Data for Exercise 6.52

Usage

Mental

Format

A data frame/tibble with 16 observations on one variable

age mental age of 16 girls

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
SIGN.test(Mental$age, md = 100)
```

Mercury	<i>Concentration of mercury in 25 lake trout</i>
---------	--

Description

Data for Example 1.9

Usage

Mercury

Format

A data frame/tibble with 25 observations on one variable

mercury a numeric vector measuring mercury (in parts per million)

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
stem(Mercury$mercury)
```

Metrent	<i>Monthly rental costs in metro areas with 1 million or more persons</i>
---------	---

Description

Data for Exercise 5.117

Usage

Metrent

Format

A data frame/tibble with 46 observations on one variable

rent monthly rent in dollars

Source

U.S. Bureau of the Census, *Housing in the Metropolitan Areas, Statistical Brief SB/94/19*, September 1994.

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
boxplot(Metrent$rent, col = "magenta")
t.test(Metrent$rent, conf.level = 0.99)$conf
```

Miller	<i>Miller personality test scores for a group of college students applying for graduate school</i>
--------	--

Description

Data for Example 5.7

Usage

Miller

Format

A data frame/tibble with 25 observations on one variable

miller scores on the Miller Personality test

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
stem(Miller$miller)
fivenum(Miller$miller)
boxplot(Miller$miller)
qqnorm(Miller$miller,col = "blue")
qqline(Miller$miller, col = "red")
```

Miller1	<i>Twenty scores on the Miller personality test</i>
---------	---

Description

Data for Exercise 1.41

Usage

Miller1

Format

A data frame/tibble with 20 observations on one variable

millier scores on the Miller personality test

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
stem(Miller1$millier)
stem(Miller1$millier, scale = 2)
```

Moisture	<i>Moisture content and depth of core sample for marine muds in eastern Louisiana</i>
----------	---

Description

Data for Exercise 9.32

Usage

Moisture

Format

A data frame/tibble with 16 observations on four variables

depth a numeric vector

moisture g of water per 100 g of dried sediment

lnmoist a numeric vector

depthsq a numeric vector

Source

Davis, J. C. (1986), *Statistics and Data Analysis in Geology*, 2d. ed., John Wiley and Sons, New York, pp. 177, 185.

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
plot(moisture ~ depth, data = Moisture)
model <- lm(moisture ~ depth, data = Moisture)
abline(model, col = "red")
plot(resid(model) ~ depth, data = Moisture)
rm(model)
```

Monoxide

Carbon monoxide emitted by smoke stacks of a manufacturer and a competitor

Description

Data for Exercise 7.45

Usage

Monoxide

Format

A data frame/tibble with ten observations on two variables

company a vector with values manufacturer and competitor

emission carbon monoxide emitted

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```

boxplot(emission ~ company, data = Monoxide, col = topo.colors(2))
t.test(emission ~ company, data = Monoxide)
wilcox.test(emission ~ company, data = Monoxide)
## Not run:
library(ggplot2)
ggplot2::ggplot(data = Monoxide, aes(x = company, y = emission)) +
  geom_boxplot() +
  theme_bw()

## End(Not run)

```

Movie

Moral attitude scale on 15 subjects before and after viewing a movie

Description

Data for Exercise 7.53

Usage

Movie

Format

A data frame/tibble with 12 observations on three variables

before moral aptitude before viewing the movie

after moral aptitude after viewing the movie

differ a numeric vector

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```

qqnorm(Movie$differ)
qqline(Movie$differ)
shapiro.test(Movie$differ)
t.test(Movie$differ, conf.level = 0.99)
wilcox.test(Movie$differ)

```

Music	<i>Improvement scores for identical twins taught music recognition by two techniques</i>
-------	--

Description

Data for Exercise 7.59

Usage

Music

Format

A data frame/tibble with 12 observations on three variables

method1 a numeric vector measuring the improvement scores on a music recognition test

method2 a numeric vector measuring the improvement scores on a music recognition test

differ method1 - method2

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
qqnorm(Music$differ)
qqline(Music$differ)
shapiro.test(Music$differ)
t.test(Music$differ)
## Not run:
library(ggplot2)
ggplot2::ggplot(data = Music, aes(x = differ)) +
  geom_dotplot() +
  theme_bw()

## End(Not run)
```

Name

*Estimated value of a brand name product and the company's revenue***Description**

Data for Exercises 2.28, 9.19, and Example 2.8

Usage

Name

Format

A data frame/tibble with 42 observations on three variables

brand a factor with levels Band-Aid, Barbie, Birds Eye, Budweiser, Camel, Campbell, Carlsberg, Coca-Cola, Colgate, Del Monte, Fisher-Price, Gordon's, Green Giant, Guinness, Haagen-Dazs, Heineken, Heinz, Hennessy, Hermes, Hershey, Ivory, Jell-o, Johnnie Walker, Kellogg, Kleenex, Kraft, Louis Vuitton, Marlboro, Nescafe, Nestle, Nivea, Oil of Olay, Pampers, Pepsi-Cola, Planters, Quaker, Sara Lee, Schweppes, Smirnoff, Tampax, Winston, and Wrigley's

value value in billions of dollars

revenue revenue in billions of dollars

Source

Financial World.

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
plot(value ~ revenue, data = Name)
model <- lm(value ~ revenue, data = Name)
abline(model, col = "red")
cor(Name$value, Name$revenue)
summary(model)
rm(model)
```

Nascar

Efficiency of pit crews for three major NASCAR teams

Description

Data for Exercise 10.53

Usage

Nascar

Format

A data frame/tibble with 36 observations on six variables

time duration of pit stop (in seconds)

team a numeric vector representing team 1, 2, or 3

ranks a numeric vector ranking each pit stop in order of speed

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
boxplot(time ~ team, data = Nascar, col = rainbow(3))
model <- lm(time ~ factor(team), data = Nascar)
summary(model)
anova(model)
rm(model)
```

Nervous

Reaction effects of 4 drugs on 25 subjects with a nervous disorder

Description

Data for Example 10.3

Usage

Nervous

Format

A data frame/tibble with 25 observations on two variables

react a numeric vector representing reaction time

drug a numeric vector indicating each of the 4 drugs

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
boxplot(react ~ drug, data = Nervous, col = rainbow(4))
model <- aov(react ~ factor(drug), data = Nervous)
summary(model)
TukeyHSD(model)
plot(TukeyHSD(model), las = 1)
```

Newsstand

Daily profits for 20 newsstands

Description

Data for Exercise 1.43

Usage

Newsstand

Format

A data frame/tibble with 20 observations on one variable

profit profit of each newsstand (in dollars)

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
stem(Newsstand$profit)
stem(Newsstand$profit, scale = 3)
```

Nfldraf2	<i>Rating, time in 40-yard dash, and weight of top defensive linemen in the 1994 NFL draft</i>
----------	--

Description

Data for Exercise 9.63

Usage

Nfldraf2

Format

A data frame/tibble with 47 observations on three variables

rating rating of each player on a scale out of 10

forty forty yard dash time (in seconds)

weight weight of each player (in pounds)

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
plot(rating ~ forty, data = Nfldraf2)
summary(lm(rating ~ forty, data = Nfldraf2))
```

Nfldraft	<i>Rating, time in 40-yard dash, and weight of top offensive linemen in the 1994 NFL draft</i>
----------	--

Description

Data for Exercises 9.10 and 9.16

Usage

Nfldraft

Format

A data frame/tibble with 29 observations on three variables

rating rating of each player on a scale out of 10

forty forty yard dash time (in seconds)

weight weight of each player (in pounds)

Source

USA Today, April 20, 1994.

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
plot(rating ~ forty, data = Nfldraft)
cor(Nfldraft$rating, Nfldraft$forty)
summary(lm(rating ~ forty, data = Nfldraft))
```

Nicotine

Nicotine content versus sales for eight major brands of cigarettes

Description

Data for Exercise 9.21

Usage

Nicotine

Format

A data frame/tibble with eight observations on two variables

nicotine nicotine content (in milligrams)

sales sales figures (in \$100,000)

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
model <- lm(sales ~ nicotine, data = Nicotine)
plot(sales ~ nicotine, data = Nicotine)
abline(model, col = "red")
summary(model)
predict(model, newdata = data.frame(nicotine = 1),
        interval = "confidence", level = 0.99)
```

normarea

Normal Area

Description

Function that computes and draws the area between two user specified values in a user specified normal distribution with a given mean and standard deviation

Usage

```
normarea(lower = -Inf, upper = Inf, m, sig)
```

Arguments

lower	the lower value
upper	the upper value
m	the mean for the population
sig	the standard deviation of the population

Author(s)

Alan T. Arnholt

Examples

```
normarea(70, 130, 100, 15)
# Finds and P(70 < X < 130) given X is N(100,15).
```

nsize	<i>Required Sample Size</i>
-------	-----------------------------

Description

Function to determine required sample size to be within a given margin of error.

Usage

```
nsize(b, sigma = NULL, p = 0.5, conf.level = 0.95, type = "mu")
```

Arguments

b	the desired bound.
sigma	population standard deviation. Not required if using type "pi".
p	estimate for the population proportion of successes. Not required if using type "mu".
conf.level	confidence level for the problem, restricted to lie between zero and one.
type	character string, one of "mu" or "pi", or just the initial letter of each, indicating the appropriate parameter. Default value is "mu".

Details

Answer is based on a normal approximation when using type "pi".

Value

Returns required sample size.

Author(s)

Alan T. Arnholt

Examples

```
nsize(b=.03, p=708/1200, conf.level=.90, type="pi")
# Returns the required sample size (n) to estimate the population
# proportion of successes with a 0.9 confidence interval
# so that the margin of error is no more than 0.03 when the
# estimate of the population propotion of successes is 708/1200.
# This is problem 5.38 on page 257 of Kitchen's BSDA.

nsize(b=.15, sigma=.31, conf.level=.90, type="mu")
# Returns the required sample size (n) to estimate the population
# mean with a 0.9 confidence interval so that the margin
# of error is no more than 0.15. This is Example 5.17 on page
# 261 of Kitchen's BSDA.
```

ntester*Normality Tester*

Description

Q-Q plots of randomly generated normal data of the same size as the tested data are generated and plotted on the perimeter of the graph while a Q-Q plot of the actual data is depicted in the center of the graph.

Usage

```
ntester(actual.data)
```

Arguments

<code>actual.data</code>	a numeric vector. Missing and infinite values are allowed, but are ignored in the calculation. The length of <code>actual.data</code> must be less than 5000 after dropping nonfinite values.
--------------------------	---

Details

Q-Q plots of randomly generated normal data of the same size as the tested data are generated and plotted on the perimeter of the graph sheet while a Q-Q plot of the actual data is depicted in the center of the graph. The p-values are calculated from the Shapiro-Wilk W-statistic. Function will only work on numeric vectors containing less than or equal to 5000 observations.

Author(s)

Alan T. Arnholt

References

Shapiro, S.S. and Wilk, M.B. (1965). An analysis of variance test for normality (complete samples). *Biometrika* **52** : 591-611.

Examples

```
ntester(rexp(50,1))
# Q-Q plot of random exponential data in center plot
# surrounded by 8 Q-Q plots of randomly generated
# standard normal data of size 50.
```

Orange

Price of oranges versus size of the harvest

Description

Data for Exercise 9.61

Usage

Orange

Format

A data frame/tibble with six observations on two variables

harvest harvest in millions of boxes

price average price charged by California growers for a 75-pound box of navel oranges

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
plot(price ~ harvest, data = Orange)
model <- lm(price ~ harvest, data = Orange)
abline(model, col = "red")
summary(model)
rm(model)
```

Orioles

Salaries of members of the Baltimore Orioles baseball team

Description

Data for Example 1.3

Usage

Orioles

Format

A data frame/tibble with 27 observations on three variables

first name a factor with levels Albert, Arthur, B. J., Brady, Cal, Charles, dl-Delino, dl-Scott, Doug, Harold, Heathcliff, Jeff, Jesse, Juan, Lenny, Mike, Rich, Ricky, Scott, Sidney, Will, and Willis

last name a factor with levels Amaral, Anderson, Baines, Belle, Bones, Bordick, Clark, Conine, Deshields, Erickson, Feters, Garcia, Guzman, Johns, Johnson, Kamieniecki, Mussina, Orosco, Otanez, Ponson, Reboulet, Rhodes, Ripken Jr., Slocumb, Surhoff, Timlin, and Webster

1999salary a numeric vector containing each player's salary (in dollars)

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
stripchart(Orioles$`1999salary`, method = "stack", pch = 19)
## Not run:
library(ggplot2)
ggplot2::ggplot(data = Orioles, aes(x = `1999salary`)) +
  geom_dotplot(dotsize = 0.5) +
  labs(x = "1999 Salary") +
  theme_bw()

## End(Not run)
```

Oxytocin	<i>Arterial blood pressure of 11 subjects before and after receiving oxytocin</i>
----------	---

Description

Data for Exercise 7.86

Usage

Oxytocin

Format

A data frame/tibble with 11 observations on three variables

subject a numeric vector indicating each subject

before mean arterial blood pressure of subject before receiving oxytocin

after mean arterial blood pressure of subject after receiving oxytocin

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
diff = Oxytocin$after - Oxytocin$before
qqnorm(diff)
qqline(diff)
shapiro.test(diff)
t.test(diff)
rm(diff)
```

Parented

Education backgrounds of parents of entering freshmen at a state university

Description

Data for Exercise 1.32

Usage

Parented

Format

A data frame/tibble with 200 observations on two variables

education a factor with levels 4yr college degree, Doctoral degree, Grad degree, H.S grad or less, Some college, and Some grad school

parent a factor with levels mother and father

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
T1 <- xtabs(~education + parent, data = Parented)
T1
barplot(t(T1), beside = TRUE, legend = TRUE, col = c("blue", "red"))
rm(T1)
## Not run:
library(ggplot2)
ggplot2::ggplot(data = Parented, aes(x = education, fill = parent)) +
  geom_bar(position = "dodge") +
```

```

theme_bw() +
  theme(axis.text.x = element_text(angle = 85, vjust = 0.5)) +
  scale_fill_manual(values = c("pink", "blue")) +
  labs(x = "", y = "")

## End(Not run)

```

Patrol	<i>Years of experience and number of tickets given by patrolpersons in New York City</i>
--------	--

Description

Data for Example 9.3

Usage

Patrol

Format

A data frame/tibble with ten observations on three variables

tickets number of tickets written per week

years patrolperson's experience (in years)

log_tickets natural log of tickets

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```

model <- lm(tickets ~ years, data = Patrol)
summary(model)
confint(model, level = 0.98)

```

Pearson

Karl Pearson's data on heights of brothers and sisters

Description

Data for Exercise 2.20

Usage

Pearson

Format

A data frame/tibble with 11 observations on three variables

family number indicating family of brother and sister pair

brother height of brother (in inches)

sister height of sister (in inches)

Source

Pearson, K. and Lee, A. (1902-3), On the Laws of Inheritance in Man, *Biometrika*, 2, 357.

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
plot(brother ~ sister, data = Pearson, col = "lightblue")
cor(Pearson$brother, Pearson$sister)
```

Phone

Length of long-distance phone calls for a small business firm

Description

Data for Exercise 6.95

Usage

Phone

Format

A data frame/tibble with 20 observations on one variable

time duration of long distance phone call (in minutes)

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
qqnorm(Phone$time)
qqline(Phone$time)
shapiro.test(Phone$time)
SIGN.test(Phone$time, md = 5, alternative = "greater")
```

Poison

Number of poisonings reported to 16 poison control centers

Description

Data for Exercise 1.113

Usage

Poison

Format

A data frame/tibble with 226,361 observations on one variable

type a factor with levels Alcohol, Cleaning agent, Cosmetics, Drugs, Insecticides, and Plants

Source

Centers for Disease Control, Atlanta, Georgia.

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```

T1 <- xtabs(~type, data = Poison)
T1
par(mar = c(5.1 + 2, 4.1, 4.1, 2.1))
barplot(sort(T1, decreasing = TRUE), las = 2, col = rainbow(6))
par(mar = c(5.1, 4.1, 4.1, 2.1))
rm(T1)
## Not run:
library(ggplot2)
ggplot2::ggplot(data = Poison, aes(x = type, fill = type)) +
  geom_bar() +
  theme_bw() +
  theme(axis.text.x = element_text(angle = 85, vjust = 0.5)) +
  guides(fill = FALSE)

## End(Not run)

```

Politic

Political party and gender in a voting district

Description

Data for Example 8.3

Usage

Politic

Format

A data frame/tibble with 250 observations on two variables

party a factor with levels republican, democrat, and other

gender a factor with levels female and male

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```

T1 <- xtabs(~party + gender, data = Politic)
T1
chisq.test(T1)
rm(T1)

```

Pollutio	<i>Air pollution index for 15 randomly selected days for a major western city</i>
----------	---

Description

Data for Exercise 5.59

Usage

Pollutio

Format

A data frame/tibble with 15 observations on one variable

inde air pollution index

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
stem(Pollutio$inde)
t.test(Pollutio$inde, conf.level = 0.98)$conf
```

Porosity	<i>Porosity measurements on 20 samples of Tensleep Sandstone, Pennsylvanian from Bighorn Basin in Wyoming</i>
----------	---

Description

Data for Exercise 5.86

Usage

Porosity

Format

A data frame/tibble with 20 observations on one variable

porosity porosity measurement (percent)

Source

Davis, J. C. (1986), *Statistics and Data Analysis in Geology*, 2nd edition, pages 63-65.

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
stem(Porosity$porosity)
fivenum(Porosity$porosity)
boxplot(Porosity$porosity, col = "lightgreen")
```

Poverty

Percent poverty and crime rate for selected cities

Description

Data for Exercise 9.11 and 9.17

Usage

Poverty

Format

A data frame/tibble with 20 observations on four variables

city a factor with levels Atlanta, Buffalo, Cincinnati, Cleveland, Dayton, O, Detroit, Flint, Mich, Fresno, C, Gary, Ind, Hartford, C, Laredo, Macon, Ga, Miami, Milwaukee, New Orleans, Newark, NJ, Rochester, NY, Shreveport, St. Louis, and Waco, Tx

poverty percent of children living in poverty

crime crime rate (per 1000 people)

population population of city

Source

Children's Defense Fund and the Bureau of Justice Statistics.

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
plot(poverty ~ crime, data = Poverty)
model <- lm(poverty ~ crime, data = Poverty)
abline(model, col = "red")
summary(model)
rm(model)
```

Precinct

Robbery rates versus percent low income in eight precincts

Description

Data for Exercise 2.2 and 2.38

Usage

Precinct

Format

A data frame/tibble with eight observations on two variables

rate robbery rate (per 1000 people)

income percent with low income

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
plot(rate ~ income, data = Precinct)
model <- (lm(rate ~ income, data = Precinct))
abline(model, col = "red")
rm(model)
```

Prejudic

Racial prejudice measured on a sample of 25 high school students

Description

Data for Exercise 5.10 and 5.22

Usage

Prejudic

Format

A data frame with 25 observations on one variable

prejud racial prejudice score

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
stem(Prejudic$prejud)
EDA(Prejudic$prejud)
```

Presiden

Ages at inauguration and death of U.S. presidents

Description

Data for Exercise 1.126

Usage

Presiden

Format

A data frame/tibble with 43 observations on five variables

first_initial a factor with levels A., B., C., D., F., G., G. W., H., J., L., M., R., T., U., W., and Z.

last_name a factor with levels Adams, Arthur, Buchanan, Bush, Carter, Cleveland, Clinton, Coolidge, Eisenhower, Fillmore, Ford, Garfield, Grant, Harding, Harrison, Hayes, Hoover, Jackson, Jefferson, Johnson, Kennedy, Lincoln, Madison, McKinley, Monroe, Nixon, Pierce, Polk, Reagan, Roosevelt, Taft, Taylor, Truman, Tyler, VanBuren, Washington, and Wilson

birth_state a factor with levels ARK, CAL, CONN, GA, IA, ILL, KY, MASS, MO, NC, NEB, NH, NJ, NY, OH, PA, SC, TEX, VA, and VT

inaugural_age President’s age at inauguration

death_age President’s age at death

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
pie(xtabs(~birth_state, data = Presiden))
stem(Presiden$inaugural_age)
stem(Presiden$death_age)
par(mar = c(5.1, 4.1 + 3, 4.1, 2.1))
stripchart(x=list(Presiden$inaugural_age, Presiden$death_age),
           method = "stack", col = c("green","brown"), pch = 19, las = 1)
par(mar = c(5.1, 4.1, 4.1, 2.1))
```

Press	<i>Degree of confidence in the press versus education level for 20 randomly selected persons</i>
-------	--

Description

Data for Exercise 9.55

Usage

Press

Format

A data frame/tibble with 20 observations on two variables

education_yrs years of education

confidence degree of confidence in the press (the higher the score, the more confidence)

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
plot(confidence ~ education_yrs, data = Press)
model <- lm(confidence ~ education_yrs, data = Press)
abline(model, col = "purple")
summary(model)
rm(model)
```

Prognost

Klopfer's prognostic rating scale for subjects receiving behavior modification therapy

Description

Data for Exercise 6.61

Usage

Prognost

Format

A data frame/tibble with 15 observations on one variable

kprs_score Klopfer's Prognostic Rating Scale score

Source

Newmark, C., et al. (1973), Predictive Validity of the Rorschach Prognostic Rating Scale with Behavior Modification Techniques, *Journal of Clinical Psychology*, 29, 246-248.

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
EDA(Prognost$kprs_score)
t.test(Prognost$kprs_score, mu = 9)
```

Program	<i>Effects of four different methods of programmed learning for statistics students</i>
---------	---

Description

Data for Exercise 10.17

Usage

Program

Format

A data frame/tibble with 44 observations on two variables

method a character variable with values method1, method2, method3, and method4

score standardized test score

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
boxplot(score ~ method, col = c("red", "blue", "green", "yellow"), data = Program)
anova(lm(score ~ method, data = Program))
TukeyHSD(aov(score ~ method, data = Program))
par(mar = c(5.1, 4.1 + 4, 4.1, 2.1))
plot(TukeyHSD(aov(score ~ method, data = Program)), las = 1)
par(mar = c(5.1, 4.1, 4.1, 2.1))
```

Psat	<i>PSAT scores versus SAT scores</i>
------	--------------------------------------

Description

Data for Exercise 2.50

Usage

Psat

Format

A data frame/tibble with seven observations on the two variables

psat PSAT score

sat SAT score

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
model <- lm(sat ~ psat, data = Psat)
par(mfrow = c(1, 2))
plot(Psat$psat, resid(model))
plot(model, which = 1)
rm(model)
par(mfrow = c(1, 1))
```

Psych

Correct responses for 24 students in a psychology experiment

Description

Data for Exercise 1.42

Usage

Psych

Format

A data frame/tibble with 23 observations on one variable

score number of correct responses in a psychology experiment

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
stem(Psych$score)
EDA(Psych$score)
```

Puerto	<i>Weekly incomes of a random sample of 50 Puerto Rican families in Miami</i>
--------	---

Description

Data for Exercise 5.22 and 5.65

Usage

Puerto

Format

A data frame/tibble with 50 observations on one variable

income weekly family income (in dollars)

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
stem(Puerto$income)
boxplot(Puerto$income, col = "purple")
t.test(Puerto$income, conf.level = .90)$conf
```

Quail	<i>Plasma LDL levels in two groups of quail</i>
-------	---

Description

Data for Exercise 1.53, 1.77, 1.88, 5.66, and 7.50

Usage

Quail

Format

A data frame/tibble with 40 observations on two variables

group a character variable with values placebo and treatment

level low-density lipoprotein (LDL) cholesterol level

Source

J. McKean, and T. Vidmar (1994), "A Comparison of Two Rank-Based Methods for the Analysis of Linear Models," *The American Statistician*, 48, 220-229.

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
boxplot(level ~ group, data = Quail, horizontal = TRUE, xlab = "LDL Level",
        col = c("yellow", "lightblue"))
```

Quality

Quality control test scores on two manufacturing processes

Description

Data for Exercise 7.81

Usage

Quality

Format

A data frame/tibble with 15 observations on two variables

process a character variable with values Process1 and Process2

score results of a quality control test

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
boxplot(score ~ process, data = Quality, col = "lightgreen")
t.test(score ~ process, data = Quality)
```

Rains	<i>Rainfall in an area of west central Kansas and four surrounding counties</i>
-------	---

Description

Data for Exercise 9.8

Usage

Rains

Format

A data frame/tibble with 35 observations on five variables

rain rainfall (in inches)

x1 rainfall (in inches)

x2 rainfall (in inches)

x3 rainfall (in inches)

x4 rainfall (in inches)

Source

R. Picard, K. Berk (1990), Data Splitting, *The American Statistician*, 44, (2), 140-147.

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
cor(Rains)
model <- lm(rain ~ x2, data = Rains)
summary(model)
```

Randd	<i>Research and development expenditures and sales of a large company</i>
-------	---

Description

Data for Exercise 9.36 and Example 9.8

Usage

Randd

Format

A data frame/tibble with 12 observations on two variables

rd research and development expenditures (in million dollars)

sales sales (in million dollars)

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
plot(sales ~ rd, data = Randd)
model <- lm(sales ~ rd, data = Randd)
abline(model, col = "purple")
summary(model)
plot(model, which = 1)
rm(model)
```

Rat	<i>Survival times of 20 rats exposed to high levels of radiation</i>
-----	--

Description

Data for Exercise 1.52, 1.76, 5.62, and 6.44

Usage

Rat

Format

A data frame/tibble with 20 observations on one variable

survival_time survival time in weeks for rats exposed to a high level of radiation

Source

J. Lawless, *Statistical Models and Methods for Lifetime Data* (New York: Wiley, 1982).

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
hist(Rat$survival_time)
qqnorm(Rat$survival_time)
qqline(Rat$survival_time)
summary(Rat$survival_time)
t.test(Rat$survival_time)
t.test(Rat$survival_time, mu = 100, alternative = "greater")
```

Ratings

Grade point averages versus teacher's ratings

Description

Data for Example 2.6

Usage

Ratings

Format

A data frame/tibble with 250 observations on two variables

rating character variable with students' ratings of instructor (A-F)

gpa students' grade point average

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
boxplot(gpa ~ rating, data = Ratings, xlab = "Student rating of instructor",
        ylab = "Student GPA")
## Not run:
library(ggplot2)
ggplot2::ggplot(data = Ratings, aes(x = rating, y = gpa, fill = rating)) +
  geom_boxplot() +
  theme_bw() +
  theme(legend.position = "none") +
  labs(x = "Student rating of instructor", y = "Student GPA")

## End(Not run)
```

Reaction	<i>Threshold reaction time for persons subjected to emotional stress</i>
----------	--

Description

Data for Example 6.11

Usage

Reaction

Format

A data frame/tibble with 12 observations on one variable

time threshold reaction time (in seconds) for persons subjected to emotional stress

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
stem(Reaction$time)
SIGN.test(Reaction$time, md = 15, alternative = "less")
```

Reading

Standardized reading scores for 30 fifth graders

Description

Data for Exercise 1.72 and 2.10

Usage

Reading

Format

A data frame/tibble with 30 observations on four variables

score standardized reading test score

sorted sorted values of score

trimmed trimmed values of sorted

winsoriz winsorized values of score

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
hist(Reading$score, main = "Exercise 1.72",  
     col = "lightgreen", xlab = "Standardized reading score")  
summary(Reading$score)  
sd(Reading$score)
```

Readiq

Reading scores versus IQ scores

Description

Data for Exercises 2.10 and 2.53

Usage

Readiq

Format

A data frame/tibble with 14 observations on two variables

reading reading achievement score

iq IQ score

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
plot(reading ~ iq, data = Readiq)
model <- lm(reading ~ iq, data = Readiq)
abline(model, col = "purple")
predict(model, newdata = data.frame(iq = c(100, 120)))
residuals(model)[c(6, 7)]
rm(model)
```

Referend

Opinion on referendum by view on freedom of the press

Description

Data for Exercise 8.20

Usage

Referend

Format

A data frame with 237 observations on two variables

choice a factor with levels A, B, and C

response a factor with levels for, against, and undecided

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
T1 <- xtabs(~choice + response, data = Referend)
T1
chisq.test(T1)
chisq.test(T1)$expected
```

Region

Pollution index taken in three regions of the country

Description

Data for Exercise 10.26

Usage

Region

Format

A data frame/tibble with 48 observations on three variables

pollution pollution index

region region of a county (west, central, and east)

ranks ranked values of pollution

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
boxplot(pollution ~ region, data = Region, col = "gray")
anova(lm(pollution ~ region, data = Region))
```

Register	<i>Maintenance cost versus age of cash registers in a department store</i>
----------	--

Description

Data for Exercise 2.3, 2.39, and 2.54

Usage

Register

Format

A data frame/tibble with nine observations on two variables

age age of cash register (in years)

cost maintenance cost of cash register (in dollars)

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
plot(cost ~ age, data = Register)
model <- lm(cost ~ age, data = Register)
abline(model, col = "red")
predict(model, newdata = data.frame(age = c(5, 10)))
plot(model, which = 1)
rm(model)
```

Rehab	<i>Rehabilitative potential of 20 prison inmates as judged by two psychiatrists</i>
-------	---

Description

Data for Exercise 7.61

Usage

Rehab

Format

A data frame/tibble with 20 observations on four variables

inmate inmate identification number

psych1 rating from first psychiatrist on the inmates rehabilitative potential

psych2 rating from second psychiatrist on the inmates rehabilitative potential

differ psych1 - psych2

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
boxplot(Rehab$differ)
qqnorm(Rehab$differ)
qqline(Rehab$differ)
t.test(Rehab$differ)
```

Remedial	<i>Math placement test score for 35 freshmen females and 42 freshmen males</i>
----------	--

Description

Data for Exercise 7.43

Usage

Remedial

Format

A data frame/tibble with 84 observations on two variables

gender a character variable with values female and male

score math placement score

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```

boxplot(score ~ gender, data = Remedial,
col = c("purple", "blue"))
t.test(score ~ gender, data = Remedial, conf.level = 0.98)
t.test(score ~ gender, data = Remedial, conf.level = 0.98)$conf
wilcox.test(score ~ gender, data = Remedial,
            conf.int = TRUE, conf.level = 0.98)

```

Rentals

Weekly rentals for 45 apartments

Description

Data for Exercise 1.122

Usage

Rentals

Format

A data frame/tibble with 45 observations on one variable

rent weekly apartment rental price (in dollars)

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```

stem(Rentals$rent)
sum(Rentals$rent < mean(Rentals$rent) - 3*sd(Rentals$rent) |
    Rentals$rent > mean(Rentals$rent) + 3*sd(Rentals$rent))

```

Repair

Recorded times for repairing 22 automobiles involved in wrecks

Description

Data for Exercise 5.77

Usage

Repair

Format

A data frame/tibble with 22 observations on one variable

time time to repair a wrecked in car (in hours)

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
stem(Repair$time)
SIGN.test(Repair$time, conf.level = 0.98)
```

Retail

Length of employment versus gross sales for 10 employees of a large retail store

Description

Data for Exercise 9.59

Usage

Retail

Format

A data frame/tibble with 10 observations on two variables

months length of employment (in months)

sales employee gross sales (in dollars)

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
plot(sales ~ months, data = Retail)
model <- lm(sales ~ months, data = Retail)
abline(model, col = "blue")
summary(model)
```

Ronbrown1	<i>Oceanography data obtained at site 1 by scientist aboard the ship Ron Brown</i>
-----------	--

Description

Data for Exercise 2.9

Usage

Ronbrown1

Format

A data frame/tibble with 75 observations on two variables

depth ocean depth (in meters)

temperature ocean temperature (in Celsius)

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
plot(temperature ~ depth, data = Ronbrown1, ylab = "Temperature")
```

Ronbrown2	<i>Oceanography data obtained at site 2 by scientist aboard the ship Ron Brown</i>
-----------	--

Description

Data for Exercise 2.56 and Example 2.4

Usage

Ronbrown2

Format

A data frame/tibble with 150 observations on three variables

depth ocean depth (in meters)

temperature ocean temperature (in Celcius)

salinity ocean salinity level

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
plot(salinity ~ depth, data = Ronbrown2)
model <- lm(salinity ~ depth, data = Ronbrown2)
summary(model)
plot(model, which = 1)
rm(model)
```

Rural	<i>Social adjustment scores for a rural group and a city group of children</i>
-------	--

Description

Data for Example 7.16

Usage

Rural

Format

A data frame/tibble with 33 observations on two variables

score child's social adjustment score

area character variable with values `city` and `rural`

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
boxplot(score ~ area, data = Rural)
wilcox.test(score ~ area, data = Rural)
## Not run:
library(dplyr)
Rural <- dplyr::mutate(Rural, r = rank(score))
Rural
t.test(r ~ area, data = Rural)

## End(Not run)
```

Salary

Starting salaries for 25 new PhD psychologist

Description

Data for Exercise 3.66

Usage

Salary

Format

A data frame/tibble with 25 observations on one variable

salary starting salary for Ph.D. psychologists (in dollars)

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
qqnorm(Salary$salary, pch = 19, col = "purple")
qqline(Salary$salary, col = "blue")
```

Salinity

*Surface-water salinity measurements from Whitewater Bay, Florida***Description**

Data for Exercise 5.27 and 5.64

Usage

Salinity

Format

A data frame/tibble with 48 observations on one variable

salinity surface-water salinity value**Source**J. Davis, *Statistics and Data Analysis in Geology*, 2nd ed. (New York: John Wiley, 1986).**References**Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.**Examples**

```
stem(Salinity$salinity)
qqnorm(Salinity$salinity, pch = 19, col = "purple")
qqline(Salinity$salinity, col = "blue")
t.test(Salinity$salinity, conf.level = 0.99)
t.test(Salinity$salinity, conf.level = 0.99)$conf
```

Sat

*SAT scores, percent taking exam and state funding per student by state for 1994, 1995 and 1999***Description**

Data for Statistical Insight Chapter 9

Usage

Sat

Format

A data frame/tibble with 102 observations on seven variables

state U.S. state

verbal verbal SAT score

math math SAT score

total combined verbal and math SAT score

percent percent of high school seniors taking the SAT

expend state expenditure per student (in dollars)

year year

Source

The 2000 World Almanac and Book of Facts, Funk and Wagnalls Corporation, New Jersey.

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
Sat94 <- Sat[Sat$year == 1994, ]
Sat94
Sat99 <- subset(Sat, year == 1999)
Sat99
stem(Sat99$total)
plot(total ~ percent, data = Sat99)
model <- lm(total ~ percent, data = Sat99)
abline(model, col = "blue")
summary(model)
rm(model)
```

Saving

Problem asset ration for savings and loan companies in California, New York, and Texas

Description

Data for Exercise 10.34 and 10.49

Usage

Saving

Format

A data frame/tibble with 65 observations on two variables

par problem-asset-ratio for Savings & Loans that were listed as being financially troubled in 1992

state U.S. state

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
boxplot(par ~ state, data = Saving, col = "red")
boxplot(par ~ state, data = Saving, log = "y", col = "red")
model <- aov(par ~ state, data = Saving)
summary(model)
plot(TukeyHSD(model))
kruskal.test(par ~ factor(state), data = Saving)
```

Scales

Readings obtained from a 100 pound weight placed on four brands of bathroom scales

Description

Data for Exercise 1.89

Usage

Scales

Format

A data frame/tibble with 20 observations on two variables

brand variable indicating brand of bathroom scale (A, B, C, or D)

reading recorded value (in pounds) of a 100 pound weight

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
boxplot(reading ~ brand, data = Scales, col = rainbow(4),
ylab = "Weight (lbs)")
## Not run:
library(ggplot2)
ggplot2::ggplot(data = Scales, aes(x = brand, y = reading, fill = brand)) +
  geom_boxplot() +
  labs(y = "weight (lbs)") +
  theme_bw() +
  theme(legend.position = "none")

## End(Not run)
```

Schizop2

Exam scores for 17 patients to assess the learning ability of schizophrenics after taking a specified dose of a tranquilizer

Description

Data for Exercise 6.99

Usage

Schizop2

Format

A data frame/tibble with 17 observations on one variable

score schizophrenics score on a second standardized exam

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
hist(Schizop2$score, xlab = "score on standardized test after a tranquilizer",
main = "Exercise 6.99", breaks = 10, col = "orange")
EDA(Schizop2$score)
SIGN.test(Schizop2$score, md = 22, alternative = "greater")
```

Schizoph	<i>Standardized exam scores for 13 patients to investigate the learning ability of schizophrenics after a specified dose of a tranquilizer</i>
----------	--

Description

Data for Example 6.10

Usage

Schizoph

Format

A data frame/tibble with 13 observations on one variable

score schizophrenics score on a standardized exam one hour after receiving a specified dose of a tranquilizer.

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
hist(Schizoph$score, xlab = "score on standardized test",
     main = "Example 6.10", breaks = 10, col = "orange")
EDA(Schizoph$score)
t.test(Schizoph$score, mu = 20)
```

Seatbelt	<i>Injury level versus seatbelt usage</i>
----------	---

Description

Data for Exercise 8.24

Usage

Seatbelt

Format

A data frame/tibble with 86,759 observations on two variables

seatbelt a factor with levels No and Yes

injuries a factor with levels None, Minimal, Minor, or Major indicating the extent of the drivers injuries

Source

Jobson, J. (1982), *Applied Multivariate Data Analysis*, Springer-Verlag, New York, p. 18.

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
T1 <- xtabs(~seatbelt + injuries, data = Seatbelt)
T1
chisq.test(T1)
rm(T1)
```

Selfdefe	<i>Self-confidence scores for 9 women before and after instructions on self-defense</i>
----------	---

Description

Data for Example 7.19

Usage

Selfdefe

Format

A data frame/tibble with nine observations on three variables

woman number identifying the woman

before before the course self-confidence score

after after the course self-confidence score

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
Selfdefe$differ <- Selfdefe$after - Selfdefe$before
Selfdefe
t.test(Selfdefe$differ, alternative = "greater")
```

Senior	<i>Reaction times of 30 senior citizens applying for drivers license renewals</i>
--------	---

Description

Data for Exercise 1.83 and 3.67

Usage

```
Senior
```

Format

A data frame/tibble with 31 observations on one variable

reaction reaction time for senior citizens applying for a driver's license renewal

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
stem(Senior$reaction)
fivenum(Senior$reaction)
boxplot(Senior$reaction, main = "Problem 1.83, part d",
        horizontal = TRUE, col = "purple")
```

Sentence	<i>Sentences of 41 prisoners convicted of a homicide offense</i>
----------	--

Description

Data for Exercise 1.123

Usage

Sentence

Format

A data frame/tibble with 41 observations on one variable

months sentence length (in months) for prisoners convicted of homicide

Source

U.S. Department of Justice, Bureau of Justice Statistics, *Prison Sentences and Time Served for Violence*, NCJ-153858, April 1995.

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
stem(Sentence$months)
ll <- mean(Sentence$months)-2*sd(Sentence$months)
ul <- mean(Sentence$months)+2*sd(Sentence$months)
limits <- c(ll, ul)
limits
rm(ul, ll, limits)
```

Shkdrug	<i>Effects of a drug and electroshock therapy on the ability to solve simple tasks</i>
---------	--

Description

Data for Exercises 10.11 and 10.12

Usage

Shkdrug

Format

A data frame/tibble with 64 observations on two variables

treatment type of treatment Drug/NoS, Drug/Shk, NoDg/NoS, or NoDrug/S

response number of tasks completed in a 10-minute period

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
boxplot(response ~ treatment, data = Shkdrug, col = "gray")
model <- lm(response ~ treatment, data = Shkdrug)
anova(model)
rm(model)
```

Shock

Effect of experimental shock on time to complete difficult task

Description

Data for Exercise 10.50

Usage

Shock

Format

A data frame/tibble with 27 observations on two variables

group grouping variable with values of Group1 (no shock), Group2 (medium shock), and Group3 (severe shock)

attempts number of attempts to complete a task

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
boxplot(attempts ~ group, data = Shock, col = "violet")
model <- lm(attempts ~ group, data = Shock)
anova(model)
rm(model)
```

Shoplift*Sales receipts versus shoplifting losses for a department store*

Description

Data for Exercise 9.58

Usage

Shoplift

Format

A data frame/tibble with eight observations on two variables

sales sales (in 1000 dollars)

loss loss (in 100 dollars)

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
plot(loss ~ sales, data = Shoplift)
model <- lm(loss ~ sales, data = Shoplift)
summary(model)
rm(model)
```

Short

James Short's measurements of the parallax of the sun

Description

Data for Exercise 6.65

Usage

Short

Format

A data frame/tibble with 158 observations on two variables

sample sample number

parallax parallax measurements (seconds of a degree)

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
hist(Short$parallax, main = "Problem 6.65",
     xlab = "", col = "orange")
SIGN.test(Short$parallax, md = 8.798)
t.test(Short$parallax, mu = 8.798)
```

Shuttle

Number of people riding shuttle versus number of automobiles in the downtown area

Description

Data for Exercise 9.20

Usage

Shuttle

Format

A data frame/tibble with 15 observations on two variables

users number of shuttle riders

autos number of automobiles in the downtown area

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
plot(autos ~ users, data = Shuttle)
model <- lm(autos ~ users, data = Shuttle)
summary(model)
rm(model)
```

SIGN.test

Sign Test

Description

This function will test a hypothesis based on the sign test and reports linearly interpolated confidence intervals for one sample problems.

Usage

```
SIGN.test(
  x,
  y = NULL,
  md = 0,
  alternative = "two.sided",
  conf.level = 0.95,
  ...
)
```

Arguments

x	numeric vector; NAs and Infs are allowed but will be removed.
y	optional numeric vector; NAs and Infs are allowed but will be removed.
md	a single number representing the value of the population median specified by the null hypothesis

<code>alternative</code>	is a character string, one of "greater", "less", or "two.sided", or the initial letter of each, indicating the specification of the alternative hypothesis. For one-sample tests, <code>alternative</code> refers to the true median of the parent population in relation to the hypothesized value of the median.
<code>conf.level</code>	confidence level for the returned confidence interval, restricted to lie between zero and one
<code>...</code>	further arguments to be passed to or from methods

Details

Computes a "Dependent-samples Sign-Test" if both `x` and `y` are provided. If only `x` is provided, computes the "Sign-Test".

Value

A list of class `hstest_S`, containing the following components:

<code>statistic</code>	the S-statistic (the number of positive differences between the data and the hypothesized median), with names attribute "S".
<code>p.value</code>	the p-value for the test
<code>conf.int</code>	is a confidence interval (vector of length 2) for the true median based on linear interpolation. The confidence level is recorded in the attribute <code>conf.level</code> . When the <code>alternative</code> is not "two.sided", the confidence interval will be half-infinite, to reflect the interpretation of a confidence interval as the set of all values <code>k</code> for which one would not reject the null hypothesis that the true mean or difference in means is <code>k</code> . Here infinity will be represented by <code>Inf</code> .
<code>estimate</code>	is a vector of length 1, giving the sample median; this estimates the corresponding population parameter. Component <code>estimate</code> has a names attribute describing its elements.
<code>null.value</code>	is the value of the median specified by the null hypothesis. This equals the input argument <code>md</code> . Component <code>null.value</code> has a names attribute describing its elements.
<code>alternative</code>	records the value of the input argument <code>alternative</code> : "greater", "less", or "two.sided"
<code>data.name</code>	a character string (vector of length 1) containing the actual name of the input vector <code>x</code>
<code>Confidence.Intervals</code>	a 3 by 3 matrix containing the lower achieved confidence interval, the interpolated confidence interval, and the upper achieved confidence interval

Null Hypothesis

For the one-sample sign-test, the null hypothesis is that the median of the population from which `x` is drawn is `md`. For the two-sample dependent case, the null hypothesis is that the median for the differences of the populations from which `x` and `y` are drawn is `md`. The alternative hypothesis indicates the direction of divergence of the population median for `x` from `md` (i.e., "greater", "less", "two.sided".)

Note

The reported confidence interval is based on linear interpolation. The lower and upper confidence levels are exact.

Author(s)

Alan T. Arnholt

References

- Gibbons, J.D. and Chakraborti, S. (1992). *Nonparametric Statistical Inference*. Marcel Dekker Inc., New York.
- Kitchens, L.J.(2003). *Basic Statistics and Data Analysis*. Duxbury.
- Conover, W. J. (1980). *Practical Nonparametric Statistics, 2nd ed.* Wiley, New York.
- Lehmann, E. L. (1975). *Nonparametrics: Statistical Methods Based on Ranks*. Holden and Day, San Francisco.

See Also

[z.test](#), [zsum.test](#), [tsum.test](#)

Examples

```
x <- c(7.8, 6.6, 6.5, 7.4, 7.3, 7., 6.4, 7.1, 6.7, 7.6, 6.8)
SIGN.test(x, md = 6.5)
# Computes two-sided sign-test for the null hypothesis
# that the population median for 'x' is 6.5. The alternative
# hypothesis is that the median is not 6.5. An interpolated 95%
# confidence interval for the population median will be computed.

reaction <- c(14.3, 13.7, 15.4, 14.7, 12.4, 13.1, 9.2, 14.2,
              14.4, 15.8, 11.3, 15.0)
SIGN.test(reaction, md = 15, alternative = "less")
# Data from Example 6.11 page 330 of Kitchens BSDA.
# Computes one-sided sign-test for the null hypothesis
# that the population median is 15. The alternative
# hypothesis is that the median is less than 15.
# An interpolated upper 95% upper bound for the population
# median will be computed.
```

Simpson	<i>Grade point averages of men and women participating in various sports-an illustration of Simpson's paradox</i>
---------	---

Description

Data for Example 1.18

Usage

Simpson

Format

A data frame/tibble with 100 observations on three variables

gpa grade point average

sport sport played (basketball, soccer, or track)

gender athlete sex (male, female)

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
boxplot(gpa ~ gender, data = Simpson, col = "violet")
boxplot(gpa ~ sport, data = Simpson, col = "lightgreen")
## Not run:
library(ggplot2)
ggplot2::ggplot(data = Simpson, aes(x = gender, y = gpa, fill = gender)) +
  geom_boxplot() +
  facet_grid(.~sport) +
  theme_bw()

## End(Not run)
```

Situp	<i>Maximum number of situps by participants in an exercise class</i>
-------	--

Description

Data for Exercise 1.47

Usage

Situp

Format

A data frame/tibble with 20 observations on one variable

number maximum number of situps completed in an exercise class after 1 month in the program

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
stem(Situp$number)
hist(Situp$number, breaks = seq(0, 70, 10), right = FALSE)
hist(Situp$number, breaks = seq(0, 70, 10), right = FALSE,
     freq = FALSE, col = "pink", main = "Problem 1.47",
     xlab = "Maximum number of situps")
lines(density(Situp$number), col = "red")
```

Skewed	<i>Illustrates the Wilcoxon Rank Sum test</i>
--------	---

Description

Data for Exercise 7.65

Usage

Skewed

Format

A data frame/tibble with 21 observations on two variables

C1 values from a sample of size 16 from a particular population

C2 values from a sample of size 14 from a particular population

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
boxplot(Skewed$C1, Skewed$C2, col = c("pink", "lightblue"))
wilcox.test(Skewed$C1, Skewed$C2)
```

Skin	<i>Survival times of closely and poorly matched skin grafts on burn patients</i>
------	--

Description

Data for Exercise 5.20

Usage

Skin

Format

A data frame/tibble with 11 observations on four variables

patient patient identification number

close graft survival time in days for a closely matched skin graft on the same burn patient

poor graft survival time in days for a poorly matched skin graft on the same burn patient

differ difference between close and poor (in days)

Source

R. F. Woolon and P. A. Lachenbruch, "Rank Tests for Censored Matched Pairs," *Biometrika*, 67(1980), 597-606.

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
stem(Skin$differ)
boxplot(Skin$differ, col = "pink")
summary(Skin$differ)
```

Slc	<i>Sodium-lithium countertransport activity on 190 individuals from six large English kindred</i>
-----	---

Description

Data for Exercise 5.116

Usage

Slc

Format

A data frame/tibble with 190 observations on one variable

slc Red blood cell sodium-lithium countertransport

Source

Roeder, K., (1994), "A Graphical Technique for Determining the Number of Components in a Mixture of Normals," *Journal of the American Statistical Association*, 89, 497-495.

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
EDA(Slc$slc)
hist(Slc$slc, freq = FALSE, xlab = "sodium lithium countertransport",
     main = "", col = "lightblue")
lines(density(Slc$slc), col = "purple")
```

Smokyph	<i>Water pH levels of 75 water samples taken in the Great Smoky Mountains</i>
---------	---

Description

Data for Exercises 6.40, 6.59, 7.10, and 7.35

Usage

Smokyph

Format

A data frame/tibble with 75 observations on three variables

waterph water sample pH level

code character variable with values low (elevation below 0.6 miles), and high (elevation above 0.6 miles)

elev elevation in miles

Source

Schmoyer, R. L. (1994), Permutation Tests for Correlation in Regression Errors, *Journal of the American Statistical Association*, 89, 1507-1516.

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
summary(Smokyph$waterph)
tapply(Smokyph$waterph, Smokyph$code, mean)
stripchart(waterph ~ code, data = Smokyph, method = "stack",
           pch = 19, col = c("red", "blue"))
t.test(Smokyph$waterph, mu = 7)
SIGN.test(Smokyph$waterph, md = 7)
t.test(waterph ~ code, data = Smokyph, alternative = "less")
t.test(waterph ~ code, data = Smokyph, conf.level = 0.90)

## Not run:
library(ggplot2)
ggplot2::ggplot(data = Smokyph, aes(x = waterph, fill = code)) +
  geom_dotplot() +
  facet_grid(code ~ .) +
  guides(fill = FALSE)

## End(Not run)
```

Snore

Snoring versus heart disease

Description

Data for Exercise 8.21

Usage

Snore

Format

A data frame/tibble with 2,484 observations on two variables

snore factor with levels nonsnorer, occasional snorer, nearly every night, and snores every night

heartdisease factor indicating whether the individual has heart disease (no or yes)

Source

Norton, P. and Dunn, E. (1985), Snoring as a Risk Factor for Disease, *British Medical Journal*, 291, 630-632.

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
T1 <- xtabs(~ heartdisease + snore, data = Snore)
T1
chisq.test(T1)
rm(T1)
```

Snow

Concentration of microparticles in snowfields of Greenland and Antarctica

Description

Data for Exercise 7.87

Usage

Snow

Format

A data frame/tibble with 34 observations on two variables

concent concentration of microparticles from melted snow (in parts per billion)

site location of snow sample (Antarctica or Greenland)

Source

Davis, J., *Statistics and Data Analysis in Geology*, John Wiley, New York.

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
boxplot(concent ~ site, data = Snow, col = c("lightblue", "lightgreen"))
```

Soccer

Weights of 25 soccer players

Description

Data for Exercise 1.46

Usage

Soccer

Format

A data frame/tibble with 25 observations on one variable

weight soccer players weight (in pounds)

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
stem(Soccer$weight, scale = 2)
hist(Soccer$weight, breaks = seq(110, 210, 10), col = "orange",
     main = "Problem 1.46 \n Weights of Soccer Players",
     xlab = "weight (lbs)", right = FALSE)
```

 Social

Median income level for 25 social workers from North Carolina

Description

Data for Exercise 6.63

Usage

Social

Format

A data frame/tibble with 25 observations on one variable

income annual income (in dollars) of North Carolina social workers with less than five years experience.

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
SIGN.test(Social$income, md = 27500, alternative = "less")
```

 Sophomor

Grade point averages, SAT scores and final grade in college algebra for 20 sophomores

Description

Data for Exercise 2.42

Usage

Sophomor

Format

A data frame/tibble with 20 observations on four variables

student identification number

gpa grade point average

sat SAT math score

exam final exam grade in college algebra

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
cor(Sophomor)
plot(exam ~ gpa, data = Sophomor)
## Not run:
library(ggplot2)
ggplot2::ggplot(data = Sophomor, aes(x = gpa, y = exam)) +
  geom_point()
ggplot2::ggplot(data = Sophomor, aes(x = sat, y = exam)) +
  geom_point()

## End(Not run)
```

South

Murder rates for 30 cities in the South

Description

Data for Exercise 1.84

Usage

South

Format

A data frame/tibble with 31 observations on one variable

rate murder rate per 100,000 people

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
boxplot(South$rate, col = "gray", ylab = "Murder rate per 100,000 people")
```

Speed

Speed reading scores before and after a course on speed reading

Description

Data for Exercise 7.58

Usage

Speed

Format

A data frame/tibble with 15 observations on four variables

before reading comprehension score before taking a speed-reading course

after reading comprehension score after taking a speed-reading course

differ after - before (comprehension reading scores)

signranks signed ranked differences

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
t.test(Speed$differ, alternative = "greater")
t.test(Speed$signranks, alternative = "greater")
wilcox.test(Pair(Speed$after, Speed$before) ~ 1, data = Speed, alternative = "greater")
```

Spellers

Standardized spelling test scores for two fourth grade classes

Description

Data for Exercise 7.82

Usage

Spellers

Format

A data frame/tibble with ten observations on two variables

teacher character variable with values Fourth and Colleague

score score on a standardized spelling test

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
boxplot(score ~ teacher, data = Spellers, col = "pink")
t.test(score ~ teacher, data = Spellers)
```

Spelling	<i>Spelling scores for 9 eighth graders before and after a 2-week course of instruction</i>
----------	---

Description

Data for Exercise 7.56

Usage

Spelling

Format

A data frame/tibble with nine observations on three variables

before spelling score before a 2-week course of instruction

after spelling score after a 2-week course of instruction

differ after - before (spelling score)

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
qqnorm(Spelling$differ)
qqline(Spelling$differ)
shapiro.test(Spelling$differ)
t.test(Spelling$differ)
```

Sports	<i>Favorite sport by gender</i>
--------	---------------------------------

Description

Data for Exercise 8.32

Usage

Sports

Format

A data frame/tibble with 200 observations on two variables

gender a factor with levels male and female

sport a factor with levels football, basketball, baseball, and tennis

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
T1 <- xtabs(~gender + sport, data = Sports)
T1
chisq.test(T1)
rm(T1)
```

Spouse	<i>Convictions in spouse murder cases by gender</i>
--------	---

Description

Data for Exercise 8.33

Usage

Spouse

Format

A data frame/tibble with 540 observations on two variables

result a factor with levels not prosecuted, pleaded guilty, convicted, and acquitted

spouse a factor with levels husband and wife

Source

Bureau of Justice Statistics (September 1995), *Spouse Murder Defendants in Large Urban Counties*, Executive Summary, NCJ-156831.

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
T1 <- xtabs(~result + spouse, data = Spouse)
T1
chisq.test(T1)
rm(T1)
```

SRS

Simple Random Sampling

Description

Computes all possible samples from a given population using simple random sampling.

Usage

```
SRS(POPvalues, n)
```

Arguments

POPvalues vector containing the population values.
n the sample size.

Value

Returns a matrix containing the possible simple random samples of size n taken from a population POPvalues.

Author(s)

Alan T. Arnholt

See Also

[Combinations](#)

Examples

```
SRS(c(5,8,3),2)
# The rows in the matrix list the values for the 3 possible
# simple random samples of size 2 from the population of 5,8, and 3.
```

Stable	<i>Times of a 2-year old stallion on a one mile run</i>
--------	---

Description

Data for Exercise 6.93

Usage

Stable

Format

A data frame/tibble with nine observations on one variable

time time (in seconds) for horse to run 1 mile

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
SIGN.test(Stable$time, md = 98.5, alternative = "greater")
```

Stamp	<i>Thicknesses of 1872 Hidalgo stamps issued in Mexico</i>
-------	--

Description

Data for Statistical Insight Chapter 1 and Exercise 5.110

Usage

Stamp

Format

A data frame/tibble with 485 observations on one variable

thickness stamp thickness (in mm)

Source

Izenman, A., Sommer, C. (1988), Philatelic Mixtures and Multimodal Densities, *Journal of the American Statistical Association*, 83, 941-953.

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
hist(Stamp$thickness, freq = FALSE, col = "lightblue",
     main = "", xlab = "stamp thickness (mm)")
lines(density(Stamp$thickness), col = "blue")
t.test(Stamp$thickness, conf.level = 0.99)
```

Statclas

Grades for two introductory statistics classes

Description

Data for Exercise 7.30

Usage

Statclas

Format

A data frame/tibble with 72 observations on two variables

class class meeting time (9am or 2pm)

score grade for an introductory statistics class

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
str(Statclas)
boxplot(score ~ class, data = Statclas, col = "red")
t.test(score ~ class, data = Statclas)
```

Statelaw	<i>Operating expenditures per resident for each of the state law enforcement agencies</i>
----------	---

Description

Data for Exercise 6.62

Usage

Statelaw

Format

A data frame/tibble with 50 observations on two variables

state U.S. state

cost dollars spent per resident on law enforcement

Source

Bureau of Justice Statistics, *Law Enforcement Management and Administrative Statistics, 1993*, NCJ-148825, September 1995, page 84.

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
EDA(Statelaw$cost)
SIGN.test(Statelaw$cost, md = 8, alternative = "less")
```

Statisti	<i>Test scores for two beginning statistics classes</i>
----------	---

Description

Data for Exercises 1.70 and 1.87

Usage

Statisti

Format

A data frame/tibble with 62 observations on two variables

class character variable with values Class1 and Class2

score test score for an introductory statistics test

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
boxplot(score ~ class, data = Statisti, col = "violet")
tapply(Statisti$score, Statisti$class, summary, na.rm = TRUE)
## Not run:
library(dplyr)
dplyr::group_by(Statisti, class) %>%
  summarize(Mean = mean(score, na.rm = TRUE),
            Median = median(score, na.rm = TRUE),
            SD = sd(score, na.rm = TRUE),
            RS = IQR(score, na.rm = TRUE))

## End(Not run)
```

Step	<i>STEP science test scores for a class of ability-grouped students</i>
------	---

Description

Data for Exercise 6.79

Usage

Step

Format

A data frame/tibble with 12 observations on one variable

score State test of educational progress (STEP) science test score

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
EDA(Step$score)
t.test(Step$score, mu = 80, alternative = "less")
wilcox.test(Step$score, mu = 80, alternative = "less")
```

Stress	<i>Short-term memory test scores on 12 subjects before and after a stressful situation</i>
--------	--

Description

Data for Example 7.20

Usage

Stress

Format

A data frame/tibble with 12 observations on two variables

prestress short term memory score before being exposed to a stressful situation

poststress short term memory score after being exposed to a stressful situation

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
diff <- Stress$prestress - Stress$poststress
qqnorm(diff)
qqline(diff)
t.test(diff)
## Not run:
wilcox.test(Pair(Stress$prestress, Stress$poststress)~1, data = Stress)

## End(Not run)
```

Study

Number of hours studied per week by a sample of 50 freshmen

Description

Data for Exercise 5.25

Usage

Study

Format

A data frame/tibble with 50 observations on one variable

hours number of hours a week freshmen reported studying for their courses

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
stem(Study$hours)
hist(Study$hours, col = "violet")
summary(Study$hours)
```

Submarin	<i>Number of German submarines sunk by U.S. Navy in World War II</i>
----------	--

Description

Data for Exercises 2.16, 2.45, and 2.59

Usage

Submarin

Format

A data frame/tibble with 16 observations on three variables

month month

reported number of submarines reported sunk by U.S. Navy

actual number of submarines actually sunk by U.S. Navy

Source

F. Mosteller, S. Fienberg, and R. Rourke, *Beginning Statistics with Data Analysis* (Reading, MA: Addison-Wesley, 1983).

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
model <- lm(actual ~ reported, data = Submarin)
summary(model)
plot(actual ~ reported, data = Submarin)
abline(model, col = "red")
rm(model)
```

Subway*Time it takes a subway to travel from the airport to downtown*

Description

Data for Exercise 5.19

Usage

Subway

Format

A data frame/tibble with 30 observations on one variable

time time (in minutes) it takes a subway to travel from the airport to downtown

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
hist(Subway$time, main = "Exercise 5.19",  
     xlab = "Time (in minutes)", col = "purple")  
summary(Subway$time)
```

Sunspot*Wolfer sunspot numbers from 1700 through 2000*

Description

Data for Example 1.7

Usage

Sunspot

Format

A data frame/tibble with 301 observations on two variables

year year

sunspots average number of sunspots for the year

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
plot(sunspots ~ year, data = Sunspot, type = "l")
## Not run:
library(ggplot2)
lattice::xyplot(sunspots ~ year, data = Sunspot,
               main = "Yearly sunspots", type = "l")
lattice::xyplot(sunspots ~ year, data = Sunspot, type = "l",
               main = "Yearly sunspots", aspect = "xy")
ggplot2::ggplot(data = Sunspot, aes(x = year, y = sunspots)) +
  geom_line() +
  theme_bw()

## End(Not run)
```

Superbowl

Margin of victory in Superbowls I to XXXV

Description

Data for Exercise 1.54

Usage

Superbowl

Format

A data frame/tibble with 35 observations on five variables

winning_team name of Superbowl winning team

winner_score winning score for the Superbowl

losing_team name of Superbowl losing team

loser_score score of losing team a numeric vector

victory_margin winner_score - loser_score

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
stem(Superbowl$viictory_margin)
```

Supercar

Top speeds attained by five makes of supercars

Description

Data for Statistical Insight Chapter 10

Usage

Supercar

Format

A data frame/tibble with 30 observations on two variables

speed top speed (in miles per hour) of car without redlining

car name of sports car

Source

Car and Drvier (July 1995).

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
boxplot(speed ~ car, data = Supercar, col = rainbow(6),  
        ylab = "Speed (mph)")  
summary(aov(speed ~ car, data = Supercar))  
anova(lm(speed ~ car, data = Supercar))
```

Tablrock*Ozone concentrations at Mt. Mitchell, North Carolina*

Description

Data for Exercise 5.63

Usage

Tablrock

Format

A data frame/tibble with 719 observations on the following 17 variables.

day date

hour time of day

ozone ozone concentration

tmp temperature (in Celcius)

vdc a numeric vector

wd a numeric vector

ws a numeric vector

amb a numeric vector

dew a numeric vector

so2 a numeric vector

no a numeric vector

no2 a numeric vector

nox a numeric vector

co a numeric vector

co2 a numeric vector

gas a numeric vector

air a numeric vector

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
summary(Tablrock$ozone)
boxplot(Tablrock$ozone)
qqnorm(Tablrock$ozone)
qqline(Tablrock$ozone)
par(mar = c(5.1 - 1, 4.1 + 2, 4.1 - 2, 2.1))
boxplot(ozone ~ day, data = Tablrock,
        horizontal = TRUE, las = 1, cex.axis = 0.7)
par(mar = c(5.1, 4.1, 4.1, 2.1))
## Not run:
library(ggplot2)
ggplot2::ggplot(data = Tablrock, aes(sample = ozone)) +
  geom_qq() +
  theme_bw()
ggplot2::ggplot(data = Tablrock, aes(x = as.factor(day), y = ozone)) +
  geom_boxplot(fill = "pink") +
  coord_flip() +
  labs(x = "") +
  theme_bw()

## End(Not run)
```

Teacher

*Average teacher's salaries across the states in the 70s 80s and 90s***Description**

Data for Exercise 5.114

Usage

Teacher

Format

A data frame/tibble with 51 observations on three variables

state U.S. state**year** academic year**salary** average salary (in dollars)**Source**

National Education Association.

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```

par(mfrow = c(3, 1))
hist(Teacher$salary[Teacher$year == "1973-74"],
     main = "Teacher salary 1973-74", xlab = "salary",
     xlim = range(Teacher$salary, na.rm = TRUE))
hist(Teacher$salary[Teacher$year == "1983-84"],
     main = "Teacher salary 1983-84", xlab = "salary",
     xlim = range(Teacher$salary, na.rm = TRUE))
hist(Teacher$salary[Teacher$year == "1993-94"],
     main = "Teacher salary 1993-94", xlab = "salary",
     xlim = range(Teacher$salary, na.rm = TRUE))
par(mfrow = c(1, 1))
## Not run:
library(ggplot2)
ggplot2::ggplot(data = Teacher, aes(x = salary)) +
  geom_histogram(fill = "purple", color = "black") +
  facet_grid(year ~ .) +
  theme_bw()

## End(Not run)

```

Tennes

Tennessee self concept scores for 20 gifted high school students

Description

Data for Exercise 6.56

Usage

Tennes

Format

A data frame/tibble with 20 observations on one variable

score Tennessee Self-Concept Scale score

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
hist(Tenness$score, freq= FALSE, main = "", col = "green",
xlab = "Tennessee Self-Concept Scale score")
lines(density(Tenness$score))
## Not run:
library(ggplot2)
ggplot2::ggplot(data = Tenness, aes(x = score, y = ..density..)) +
  geom_histogram(binwidth = 2, fill = "purple", color = "black") +
  geom_density(color = "red", fill = "pink", alpha = 0.3) +
  theme_bw()

## End(Not run)
```

Tensile	<i>Tensile strength of plastic bags from two production runs</i>
---------	--

Description

Data for Example 7.11

Usage

Tensile

Format

A data frame/tibble with 72 observations on two variables

- tensile** plastic bag tensile strength (pounds per square inch)
- run** factor with run number (1 or 2)

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
boxplot(tensile ~ run, data = Tensile,
col = c("purple", "cyan"))
t.test(tensile ~ run, data = Tensile)
```

Test1	<i>Grades on the first test in a statistics class</i>
-------	---

Description

Data for Exercise 5.80

Usage

Test1

Format

A data frame/tibble with 25 observations on one variable

score score on first statistics exam

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
stem(Test1$score)
boxplot(Test1$score, col = "purple")
```

Thermal	<i>Heat loss of thermal pane windows versus outside temperature</i>
---------	---

Description

Data for Example 9.5

Usage

Thermal

Format

A data frame/tibble with 12 observations on the two variables

temp temperature (degrees Celcius)

loss heat loss (BTUs)

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
model <- lm(loss ~ temp, data = Thermal)
summary(model)
plot(loss ~ temp, data = Thermal)
abline(model, col = "red")
rm(model)
```

Tiaa

1999-2000 closing prices for TIAA-CREF stocks

Description

Data for your enjoyment

Usage

Tiaa

Format

A data frame/tibble with 365 observations on four variables

crefstk closing price (in dollars)

crefgwt closing price (in dollars)

tiaa closing price (in dollars)

date day of the year

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
data(Tiaa)
```

Ticket	<i>Time to complete an airline ticket reservation</i>
--------	---

Description

Data for Exercise 5.18

Usage

Ticket

Format

A data frame/tibble with 20 observations on one variable

time time (in seconds) to check out a reservation

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
EDA(Ticket$time)
```

Toaster	<i>Consumer Reports (Oct 94) rating of toaster ovens versus the cost</i>
---------	--

Description

Data for Exercise 9.36

Usage

Toaster

Format

A data frame/tibble with 17 observations on three variables

toaster name of toaster

score Consumer Reports score

cost price of toaster (in dollars)

Source

Consumer Reports (October 1994).

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
plot(cost ~ score, data = Toaster)
model <- lm(cost ~ score, data = Toaster)
summary(model)
names(summary(model))
summary(model)$r.squared
plot(model, which = 1)
```

Tonsils

Size of tonsils collected from 1,398 children

Description

Data for Exercise 2.78

Usage

Tonsils

Format

A data frame/tibble with 1,398 observations on two variables

size a factor with levels Normal, Large, and Very Large

status a factor with levels Carrier and Non-carrier

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
T1 <- xtabs(~size + status, data = Tonsils)
T1
prop.table(T1, 1)
prop.table(T1, 1)[2, 1]
barplot(t(T1), legend = TRUE, beside = TRUE, col = c("red", "green"))
## Not run:
library(dplyr)
library(ggplot2)
NDF <- dplyr::count(Tonsils, size, status)
ggplot2::ggplot(data = NDF, aes(x = size, y = n, fill = status)) +
  geom_bar(stat = "identity", position = "dodge") +
  scale_fill_manual(values = c("red", "green")) +
  theme_bw()

## End(Not run)
```

Tort

The number of torts, average number of months to process a tort, and county population from the court files of the nation's largest counties

Description

Data for Exercise 5.13

Usage

Tort

Format

A data frame/tibble with 45 observations on five variables

county U.S. county

months average number of months to process a tort

population population of the county

torts number of torts

rate rate per 10,000 residents

Source

U.S. Department of Justice, *Tort Cases in Large Counties*, Bureau of Justice Statistics Special Report, April 1995.

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
EDA(Tort$months)
```

Toxic

Hazardous waste sites near minority communities

Description

Data for Exercises 1.55, 5.08, 5.109, 8.58, and 10.35

Usage

Toxic

Format

A data frame/tibble with 51 observations on five variables

state U.S. state

region U.S. region

sites number of commercial hazardous waste sites

minority percent of minorities living in communities with commercial hazardous waste sites

percent a numeric vector

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
hist(Toxic$sites, col = "red")
hist(Toxic$minority, col = "blue")
qqnorm(Toxic$minority)
qqline(Toxic$minority)
boxplot(sites ~ region, data = Toxic, col = "lightgreen")
tapply(Toxic$sites, Toxic$region, median)
kruskal.test(sites ~ factor(region), data = Toxic)
```

Track

National Olympic records for women in several races

Description

Data for Exercises 2.97, 5.115, and 9.62

Usage

Track

Format

A data frame with 55 observations on eight variables

country athlete's country

100m time in seconds for 100 m

200m time in seconds for 200 m

400m time in seconds for 400 m

800m time in minutes for 800 m

1500m time in minutes for 1500 m

3000m time in minutes for 3000 m

marathon time in minutes for marathon

Source

Dawkins, B. (1989), "Multivariate Analysis of National Track Records," *The American Statistician*, 43(2), 110-115.

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
plot(`200m` ~ `100m`, data = Track)
plot(`400m` ~ `100m`, data = Track)
plot(`400m` ~ `200m`, data = Track)
cor(Track[, 2:8])
```

Track15

*Olympic winning times for the men's 1500-meter run***Description**

Data for Exercise 1.36

Usage

Track15

Format

A data frame/tibble with 26 observations on two variables

year Olympic year**time** Olympic winning time (in seconds) for the 1500-meter run**Source***The World Almanac and Book of Facts*, 2000.**References**Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.**Examples**

```
plot(time~ year, data = Track15, type = "b", pch = 19,
      ylab = "1500m time in seconds", col = "green")
```

Treatments

*Illustrates analysis of variance for three treatment groups***Description**

Data for Exercise 10.44

Usage

Treatments

Format

A data frame/tibble with 24 observations on two variables

score score from an experiment

group factor with levels 1, 2, and 3

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
boxplot(score ~ group, data = Treatments, col = "violet")
summary(aov(score ~ group, data = Treatments))
summary(lm(score ~ group, data = Treatments))
anova(lm(score ~ group, data = Treatments))
```

Trees

Number of trees in 20 grids

Description

Data for Exercise 1.50

Usage

Trees

Format

A data frame/tibble with 20 observations on one variable

number number of trees in a grid

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
stem(Trees$number)
hist(Trees$number, main = "Exercise 1.50", xlab = "number",
     col = "brown")
```

Trucks	<i>Miles per gallon for standard 4-wheel drive trucks manufactured by Chevrolet, Dodge and Ford</i>
--------	---

Description

Data for Example 10.2

Usage

Trucks

Format

A data frame/tibble with 15 observations on two variables

mpg miles per gallon

truck a factor with levels chevy, dodge, and ford

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
boxplot(mpg ~ truck, data = Trucks, horizontal = TRUE, las = 1)
summary(aov(mpg ~ truck, data = Trucks))
```

tsum.test	<i>Summarized t-test</i>
-----------	--------------------------

Description

Performs a one-sample, two-sample, or a Welch modified two-sample t-test based on user supplied summary information. Output is identical to that produced with `t.test`.

Usage

```
tsum.test(
  mean.x,
  s.x = NULL,
  n.x = NULL,
  mean.y = NULL,
  s.y = NULL,
  n.y = NULL,
  alternative = "two.sided",
  mu = 0,
  var.equal = FALSE,
  conf.level = 0.95
)
```

Arguments

<code>mean.x</code>	a single number representing the sample mean of x
<code>s.x</code>	a single number representing the sample standard deviation for x
<code>n.x</code>	a single number representing the sample size for x
<code>mean.y</code>	a single number representing the sample mean of y
<code>s.y</code>	a single number representing the sample standard deviation for y
<code>n.y</code>	a single number representing the sample size for y
<code>alternative</code>	is a character string, one of "greater", "less" or "two.sided", or just the initial letter of each, indicating the specification of the alternative hypothesis. For one-sample tests, <code>alternative</code> refers to the true mean of the parent population in relation to the hypothesized value <code>mu</code> . For the standard two-sample tests, <code>alternative</code> refers to the difference between the true population mean for x and that for y, in relation to <code>mu</code> . For the one-sample and paired t-tests, <code>alternative</code> refers to the true mean of the parent population in relation to the hypothesized value <code>mu</code> . For the standard and Welch modified two-sample t-tests, <code>alternative</code> refers to the difference between the true population mean for x and that for y, in relation to <code>mu</code> .
<code>mu</code>	is a single number representing the value of the mean or difference in means specified by the null hypothesis.
<code>var.equal</code>	logical flag: if TRUE, the variances of the parent populations of x and y are assumed equal. Argument <code>var.equal</code> should be supplied only for the two-sample tests.
<code>conf.level</code>	is the confidence level for the returned confidence interval; it must lie between zero and one.

Details

If `y` is `NULL`, a one-sample t-test is carried out with `x`. If `y` is not `NULL`, either a standard or Welch modified two-sample t-test is performed, depending on whether `var.equal` is `TRUE` or `FALSE`.

Value

A list of class `htest`, containing the following components:

<code>statistic</code>	the t-statistic, with names attribute <code>"t"</code>
<code>parameters</code>	is the degrees of freedom of the t-distribution associated with <code>statistic</code> . Component <code>parameters</code> has names attribute <code>"df"</code> .
<code>p.value</code>	the p-value for the test.
<code>conf.int</code>	is a confidence interval (vector of length 2) for the true mean or difference in means. The confidence level is recorded in the attribute <code>conf.level</code> . When <code>alternative</code> is not <code>"two.sided"</code> , the confidence interval will be half-infinite, to reflect the interpretation of a confidence interval as the set of all values <code>k</code> for which one would not reject the null hypothesis that the true mean or difference in means is <code>k</code> . Here infinity will be represented by <code>Inf</code> .
<code>estimate</code>	vector of length 1 or 2, giving the sample mean(s) or mean of differences; these estimate the corresponding population parameters. Component <code>estimate</code> has a names attribute describing its elements.
<code>null.value</code>	the value of the mean or difference in means specified by the null hypothesis. This equals the input argument <code>mu</code> . Component <code>null.value</code> has a names attribute describing its elements.
<code>alternative</code>	records the value of the input argument <code>alternative</code> : <code>"greater"</code> , <code>"less"</code> or <code>"two.sided"</code> .
<code>data.name</code>	a character string (vector of length 1) containing the names <code>x</code> and <code>y</code> for the two summarized samples.

Null Hypothesis

For the one-sample t-test, the null hypothesis is that the mean of the population from which `x` is drawn is `mu`. For the standard and Welch modified two-sample t-tests, the null hypothesis is that the population mean for `x` less that for `y` is `mu`.

The alternative hypothesis in each case indicates the direction of divergence of the population mean for `x` (or difference of means for `x` and `y`) from `mu` (i.e., `"greater"`, `"less"`, or `"two.sided"`).

Author(s)

Alan T. Arnholt

References

- Kitchens, L.J. (2003). *Basic Statistics and Data Analysis*. Duxbury.
- Hogg, R. V. and Craig, A. T. (1970). *Introduction to Mathematical Statistics, 3rd ed.* Toronto, Canada: Macmillan.

Mood, A. M., Graybill, F. A. and Boes, D. C. (1974). *Introduction to the Theory of Statistics*, 3rd ed. New York: McGraw-Hill.

Snedecor, G. W. and Cochran, W. G. (1980). *Statistical Methods*, 7th ed. Ames, Iowa: Iowa State University Press.

See Also

[z.test](#), [zsum.test](#)

Examples

```
tsum.test(mean.x=5.6, s.x=2.1, n.x=16, mu=4.9, alternative="greater")
# Problem 6.31 on page 324 of BSDA states: The chamber of commerce
# of a particular city claims that the mean carbon dioxide
# level of air pollution is no greater than 4.9 ppm. A random
# sample of 16 readings resulted in a sample mean of 5.6 ppm,
# and s=2.1 ppm. One-sided one-sample t-test. The null
# hypothesis is that the population mean for 'x' is 4.9.
# The alternative hypothesis states that it is greater than 4.9.

x <- rnorm(12)
tsum.test(mean(x), sd(x), n.x=12)
# Two-sided one-sample t-test. The null hypothesis is that
# the population mean for 'x' is zero. The alternative
# hypothesis states that it is either greater or less
# than zero. A confidence interval for the population mean
# will be computed. Note: above returns same answer as:
t.test(x)

x <- c(7.8, 6.6, 6.5, 7.4, 7.3, 7.0, 6.4, 7.1, 6.7, 7.6, 6.8)
y <- c(4.5, 5.4, 6.1, 6.1, 5.4, 5.0, 4.1, 5.5)
tsum.test(mean(x), s.x=sd(x), n.x=11, mean(y), s.y=sd(y), n.y=8, mu=2)
# Two-sided standard two-sample t-test. The null hypothesis
# is that the population mean for 'x' less that for 'y' is 2.
# The alternative hypothesis is that this difference is not 2.
# A confidence interval for the true difference will be computed.
# Note: above returns same answer as:
t.test(x, y)

tsum.test(mean(x), s.x=sd(x), n.x=11, mean(y), s.y=sd(y), n.y=8, conf.level=0.90)
# Two-sided standard two-sample t-test. The null hypothesis
# is that the population mean for 'x' less that for 'y' is zero.
# The alternative hypothesis is that this difference is not
# zero. A 90% confidence interval for the true difference will
# be computed. Note: above returns same answer as:
t.test(x, y, conf.level=0.90)
```

Tv	<i>Percent of students that watch more than 6 hours of TV per day versus national math test scores</i>
----	--

Description

Data for Examples 2.1 and 2.7

Usage

Tv

Format

A data frame/tibble with 53 observations on three variables

state U.S. state

percent percent of students who watch more than six hours of TV a day

test state average on national math test

Source

Educational Testing Services.

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
plot(test ~ percent, data = Tv, col = "blue")
cor(Tv$test, Tv$percent)
```

Twin	<i>Intelligence test scores for identical twins in which one twin is given a drug</i>
------	---

Description

Data for Exercise 7.54

Usage

Twin

Format

A data frame/tibble with nine observations on three variables

twinA score on intelligence test without drug

twinB score on intelligence test after taking drug

differ $\text{twinA} - \text{twinB}$

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
qqnorm(Twin$differ)
qqline(Twin$differ)
shapiro.test(Twin$differ)
t.test(Twin$differ)
```

Undergrad

Data set describing a sample of undergraduate students

Description

Data for Exercise 1.15

Usage

Undergrad

Format

A data frame/tibble with 100 observations on six variables

gender character variable with values Female and Male

major college major

class college year group classification

gpa grade point average

sat Scholastic Assessment Test score

drops number of courses dropped

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
stripchart(gpa ~ class, data = Undergrad, method = "stack",
col = c("blue", "red", "green", "lightblue"),
pch = 19, main = "GPA versus Class")
stripchart(gpa ~ gender, data = Undergrad, method = "stack",
col = c("red", "blue"), pch = 19,
main = "GPA versus Gender")
stripchart(sat ~ drops, data = Undergrad, method = "stack",
col = c("blue", "red", "green", "lightblue"),
pch = 19, main = "SAT versus Drops")
stripchart(drops ~ gender, data = Undergrad, method = "stack",
col = c("red", "blue"), pch = 19, main = "Drops versus Gender")

## Not run:
library(ggplot2)
ggplot2::ggplot(data = Undergrad, aes(x = sat, y = drops, fill = factor(drops))) +
  facet_grid(drops ~.) +
  geom_dotplot() +
  guides(fill = FALSE)

## End(Not run)
```

Vacation	<i>Number of days of paid holidays and vacation leave for sample of 35 textile workers</i>
----------	--

Description

Data for Exercise 6.46 and 6.98

Usage

Vacation

Format

A data frame/tibble with 35 observations on one variable

number number of days of paid holidays and vacation leave taken

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
boxplot(Vacation$number, col = "violet")
hist(Vacation$number, main = "Exercise 6.46", col = "blue",
      xlab = "number of days of paid holidays and vacation leave taken")
t.test(Vacation$number, mu = 24)
```

Vaccine

Reported serious reactions due to vaccines in 11 southern states

Description

Data for Exercise 1.111

Usage

Vaccine

Format

A data frame/tibble with 11 observations on two variables

state U.S. state

number number of reported serious reactions per million doses of a vaccine

Source

Center for Disease Control, Atlanta, Georgia.

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
stem(Vaccine$number, scale = 2)
fn <- fivenum(Vaccine$number)
fn
iqr <- IQR(Vaccine$number)
iqr
```

Vehicle	<i>Fatality ratings for foreign and domestic vehicles</i>
---------	---

Description

Data for Exercise 8.34

Usage

Vehicle

Format

A data frame/tibble with 151 observations on two variables

make a factor with levels domestic and foreign

rating a factor with levels Much better than average, Above average, Average, Below average, and Much worse than average

Source

Insurance Institute for Highway Safety and the Highway Loss Data Institute, 1995.

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
T1 <- xtabs(~make + rating, data = Vehicle)
T1
chisq.test(T1)
```

Verbal	<i>Verbal test scores and number of library books checked out for 15 eighth graders</i>
--------	---

Description

Data for Exercise 9.30

Usage

Verbal

Format

A data frame/tibble with 15 observations on two variables

number number of library books checked out

verbal verbal test score

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
plot(verbal ~ number, data = Verbal)
abline(lm(verbal ~ number, data = Verbal), col = "red")
summary(lm(verbal ~ number, data = Verbal))
```

Victoria	<i>Number of sunspots versus mean annual level of Lake Victoria Nyanza from 1902 to 1921</i>
----------	--

Description

Data for Exercise 2.98

Usage

Victoria

Format

A data frame/tibble with 20 observations on three variables

year year

level mean annual level of Lake Victoria Nyanza

sunspot number of sunspots

Source

N. Shaw, *Manual of Meteorology*, Vol. 1 (London: Cambridge University Press, 1942), p. 284; and F. Mosteller and J. W. Tukey, *Data Analysis and Regression* (Reading, MA: Addison-Wesley, 1977).

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
plot(level ~ sunspot, data = Victoria)
model <- lm(level ~ sunspot, data = Victoria)
summary(model)
rm(model)
```

Viscosit

Viscosity measurements of a substance on two different days

Description

Data for Exercise 7.44

Usage

Viscosit

Format

A data frame/tibble with 11 observations on two variables

first viscosity measurement for a certain substance on day one

second viscosity measurement for a certain substance on day two

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
boxplot(Viscosit$first, Viscosit$second, col = "blue")
t.test(Viscosit$first, Viscosit$second, var.equal = TRUE)
```

Visual	<i>Visual acuity of a group of subjects tested under a specified dose of a drug</i>
--------	---

Description

Data for Exercise 5.6

Usage

Visual

Format

A data frame/tibble with 18 observations on one variable

visual visual acuity measurement

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
stem(Visual$visual)
boxplot(Visual$visual, col = "purple")
```

Vocab	<i>Reading scores before and after vocabulary training for 14 employees who did not complete high school</i>
-------	--

Description

Data for Exercise 7.80

Usage

Vocab

Format

A data frame/tibble with 14 observations on two variables

first reading test score before formal vocabulary training

second reading test score after formal vocabulary training

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
t.test(Pair(Vocab$first, Vocab$second) ~ 1)
```

Wastewat	<i>Volume of injected waste water from Rocky Mountain Arsenal and number of earthquakes near Denver</i>
----------	---

Description

Data for Exercise 9.18

Usage

Wastewat

Format

A data frame/tibble with 44 observations on two variables

gallons injected water (in million gallons)

number number of earthquakes detected in Denver

Source

Davis, J. C. (1986), *Statistics and Data Analysis in Geology*, 2 ed., John Wiley and Sons, New York, p. 228, and Bardwell, G. E. (1970), Some Statistical Features of the Relationship between Rocky Mountain Arsenal Waste Disposal and Frequency of Earthquakes, *Geological Society of America, Engineering Geology Case Histories*, 8, 33-337.

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
plot(number ~ gallons, data = Wastewat)
model <- lm(number ~ gallons, data = Wastewat)
summary(model)
anova(model)
plot(model, which = 2)
```

Weather94*Weather casualties in 1994*

Description

Data for Exercise 1.30

Usage

Weather94

Format

A data frame/tibble with 388 observations on one variable

type factor with levels Extreme Temp, Flash Flood, Fog, High Wind, Hurricane, Lighting, Other, River Flood, Thunderstorm, Tornado, and Winter Weather

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
T1 <- xtabs(~type, data = Weather94)
T1
par(mar = c(5.1 + 2, 4.1 - 1, 4.1 - 2, 2.1))
barplot(sort(T1, decreasing = TRUE), las = 2, col = rainbow(11))
par(mar = c(5.1, 4.1, 4.1, 2.1))
## Not run:
library(ggplot2)
T2 <- as.data.frame(T1)
T2
ggplot2::ggplot(data = T2, aes(x = reorder(type, Freq), y = Freq)) +
  geom_bar(stat = "identity", fill = "purple") +
  theme_bw() +
  theme(axis.text.x = element_text(angle = 55, vjust = 0.5)) +
  labs(x = "", y = "count")

## End(Not run)
```

Wheat	<i>Price of a bushel of wheat versus the national weekly earnings of production workers</i>
-------	---

Description

Data for Exercise 2.11

Usage

Wheat

Format

A data frame/tibble with 19 observations on three variables

year year

earnings national weekly earnings (in dollars) for production workers

price price for a bushel of wheat (in dollars)

Source

The World Almanac and Book of Facts, 2000.

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
par(mfrow = c(1, 2))
plot(earnings ~ year, data = Wheat)
plot(price ~ year, data = Wheat)
par(mfrow = c(1, 1))
```

Windmill

Direct current produced by different wind velocities

Description

Data for Exercise 9.34

Usage

Windmill

Format

A data frame/tibble with 25 observations on two variables

velocity wind velocity (miles per hour)

output power generated (DC volts)

Source

Joglekar, et al. (1989), Lack of Fit Testing when Replicates Are Not Available, *The American Statistician*, 43,(3), 135-143.

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
summary(lm(output ~ velocity, data = Windmill))
anova(lm(output ~ velocity, data = Windmill))
```

Window

Wind leakage for storm windows exposed to a 50 mph wind

Description

Data for Exercise 6.54

Usage

Window

Format

A data frame/tibble with nine observations on two variables

window window number

leakage percent leakage from a 50 mph wind

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
SIGN.test(Window$leakage, md = 0.125, alternative = "greater")
```

Wins	<i>Baseball team wins versus seven independent variables for National league teams in 1990</i>
------	--

Description

Data for Exercise 9.23

Usage

Wins

Format

A data frame with 12 observations on nine variables

team name of team

wins number of wins

batavg batting average

rbi runs batted in

stole bases stole

strkout number of strikeouts

caught number of times caught stealing

errors number of errors

era earned run average

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
plot(wins ~ era, data = Wins)
## Not run:
library(ggplot2)
ggplot2::ggplot(data = Wins, aes(x = era, y = wins)) +
  geom_point() +
  geom_smooth(method = "lm", se = FALSE) +
  theme_bw()

## End(Not run)
```

Wool

Strength tests of two types of wool fabric

Description

Data for Exercise 7.42

Usage

Wool

Format

A data frame/tibble with 20 observations on two variables

type type of wool (Type I, Type 2)

strength strength of wool

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
boxplot(strength ~ type, data = Wool, col = c("blue", "purple"))
t.test(strength ~ type, data = Wool, var.equal = TRUE)
```

Yearsunspot

Monthly sunspot activity from 1974 to 2000

Description

Data for Exercise 2.7

Usage

Yearsunspot

Format

A data frame/tibble with 252 observations on two variables

number average number of sunspots

year date

Source

NASA/Marshall Space Flight Center, Huntsville, AL 35812.

References

Kitchens, L. J. (2003) *Basic Statistics and Data Analysis*. Pacific Grove, CA: Brooks/Cole, a division of Thomson Learning.

Examples

```
plot(number ~ year, data = Yearsunspot)
```

z.test

Z-test

Description

This function is based on the standard normal distribution and creates confidence intervals and tests hypotheses for both one and two sample problems.

Usage

```
z.test(
  x,
  y = NULL,
  alternative = "two.sided",
  mu = 0,
  sigma.x = NULL,
  sigma.y = NULL,
  conf.level = 0.95
)
```

Arguments

<code>x</code>	numeric vector; NAs and Infs are allowed but will be removed.
<code>y</code>	numeric vector; NAs and Infs are allowed but will be removed.
<code>alternative</code>	character string, one of "greater", "less" or "two.sided", or the initial letter of each, indicating the specification of the alternative hypothesis. For one-sample tests, <code>alternative</code> refers to the true mean of the parent population in relation to the hypothesized value <code>mu</code> . For the standard two-sample tests, <code>alternative</code> refers to the difference between the true population mean for <code>x</code> and that for <code>y</code> , in relation to <code>mu</code> .
<code>mu</code>	a single number representing the value of the mean or difference in means specified by the null hypothesis
<code>sigma.x</code>	a single number representing the population standard deviation for <code>x</code>
<code>sigma.y</code>	a single number representing the population standard deviation for <code>y</code>
<code>conf.level</code>	confidence level for the returned confidence interval, restricted to lie between zero and one

Details

If `y` is `NULL`, a one-sample z-test is carried out with `x`. If `y` is not `NULL`, a standard two-sample z-test is performed.

Value

A list of class `htest`, containing the following components:

<code>statistic</code>	the z-statistic, with names attribute "z"
<code>p.value</code>	the p-value for the test
<code>conf.int</code>	is a confidence interval (vector of length 2) for the true mean or difference in means. The confidence level is recorded in the attribute <code>conf.level</code> . When <code>alternative</code> is not "two.sided", the confidence interval will be half-infinite, to reflect the interpretation of a confidence interval as the set of all values <code>k</code> for which one would not reject the null hypothesis that the true mean or difference in means is <code>k</code> . Here infinity will be represented by <code>Inf</code> .

estimate	vector of length 1 or 2, giving the sample mean(s) or mean of differences; these estimate the corresponding population parameters. Component estimate has a names attribute describing its elements.
null.value	is the value of the mean or difference in means specified by the null hypothesis. This equals the input argument mu. Component null.value has a names attribute describing its elements.
alternative	records the value of the input argument alternative: "greater", "less" or "two.sided".
data.name	a character string (vector of length 1) containing the actual names of the input vectors x and y

Null Hypothesis

For the one-sample z-test, the null hypothesis is that the mean of the population from which x is drawn is mu. For the standard two-sample z-tests, the null hypothesis is that the population mean for x less than for y is mu.

The alternative hypothesis in each case indicates the direction of divergence of the population mean for x (or difference of means for x and y) from mu (i.e., "greater", "less", "two.sided").

Author(s)

Alan T. Arnholt

References

- Kitchens, L.J. (2003). *Basic Statistics and Data Analysis*. Duxbury.
- Hogg, R. V. and Craig, A. T. (1970). *Introduction to Mathematical Statistics*, 3rd ed. Toronto, Canada: Macmillan.
- Mood, A. M., Graybill, F. A. and Boes, D. C. (1974). *Introduction to the Theory of Statistics*, 3rd ed. New York: McGraw-Hill.
- Snedecor, G. W. and Cochran, W. G. (1980). *Statistical Methods*, 7th ed. Ames, Iowa: Iowa State University Press.

See Also

[zsum.test](#), [tsum.test](#)

Examples

```
x <- rnorm(12)
z.test(x,sigma.x=1)
# Two-sided one-sample z-test where the assumed value for
# sigma.x is one. The null hypothesis is that the population
# mean for 'x' is zero. The alternative hypothesis states
# that it is either greater or less than zero. A confidence
# interval for the population mean will be computed.

x <- c(7.8, 6.6, 6.5, 7.4, 7.3, 7., 6.4, 7.1, 6.7, 7.6, 6.8)
```

```

y <- c(4.5, 5.4, 6.1, 6.1, 5.4, 5., 4.1, 5.5)
z.test(x, sigma.x=0.5, y, sigma.y=0.5, mu=2)
# Two-sided standard two-sample z-test where both sigma.x
# and sigma.y are both assumed to equal 0.5. The null hypothesis
# is that the population mean for 'x' less that for 'y' is 2.
# The alternative hypothesis is that this difference is not 2.
# A confidence interval for the true difference will be computed.

z.test(x, sigma.x=0.5, y, sigma.y=0.5, conf.level=0.90)
# Two-sided standard two-sample z-test where both sigma.x and
# sigma.y are both assumed to equal 0.5. The null hypothesis
# is that the population mean for 'x' less that for 'y' is zero.
# The alternative hypothesis is that this difference is not
# zero. A 90% confidence interval for the true difference will
# be computed.
rm(x, y)

```

zsum.test

Summarized z-test

Description

This function is based on the standard normal distribution and creates confidence intervals and tests hypotheses for both one and two sample problems based on summarized information the user passes to the function. Output is identical to that produced with `z.test`.

Usage

```

zsum.test(
  mean.x,
  sigma.x = NULL,
  n.x = NULL,
  mean.y = NULL,
  sigma.y = NULL,
  n.y = NULL,
  alternative = "two.sided",
  mu = 0,
  conf.level = 0.95
)

```

Arguments

<code>mean.x</code>	a single number representing the sample mean of x
<code>sigma.x</code>	a single number representing the population standard deviation for x
<code>n.x</code>	a single number representing the sample size for x
<code>mean.y</code>	a single number representing the sample mean of y

<code>sigma.y</code>	a single number representing the population standard deviation for y
<code>n.y</code>	a single number representing the sample size for y
<code>alternative</code>	is a character string, one of "greater", "less" or "two.sided", or the initial letter of each, indicating the specification of the alternative hypothesis. For one-sample tests, alternative refers to the true mean of the parent population in relation to the hypothesized value mu. For the standard two-sample tests, alternative refers to the difference between the true population mean for x and that for y, in relation to mu.
<code>mu</code>	a single number representing the value of the mean or difference in means specified by the null hypothesis
<code>conf.level</code>	confidence level for the returned confidence interval, restricted to lie between zero and one

Details

If y is NULL, a one-sample z-test is carried out with x. If y is not NULL, a standard two-sample z-test is performed.

Value

A list of class `htest`, containing the following components:

<code>statistic</code>	the z-statistic, with names attribute z.
<code>p.value</code>	the p-value for the test
<code>conf.int</code>	is a confidence interval (vector of length 2) for the true mean or difference in means. The confidence level is recorded in the attribute <code>conf.level</code> . When alternative is not "two.sided", the confidence interval will be half-infinite, to reflect the interpretation of a confidence interval as the set of all values k for which one would not reject the null hypothesis that the true mean or difference in means is k. Here, infinity will be represented by <code>Inf</code> .
<code>estimate</code>	vector of length 1 or 2, giving the sample mean(s) or mean of differences; these estimate the corresponding population parameters. Component <code>estimate</code> has a names attribute describing its elements.
<code>null.value</code>	the value of the mean or difference in means specified by the null hypothesis. This equals the input argument mu. Component <code>null.value</code> has a names attribute describing its elements.
<code>alternative</code>	records the value of the input argument alternative: "greater", "less" or "two.sided".
<code>data.name</code>	a character string (vector of length 1) containing the names x and y for the two summarized samples

Null Hypothesis

For the one-sample z-test, the null hypothesis is that the mean of the population from which x is drawn is mu. For the standard two-sample z-tests, the null hypothesis is that the population mean for x less than that for y is mu.

The alternative hypothesis in each case indicates the direction of divergence of the population mean for x (or difference of means of x and y) from mu (i.e., "greater", "less", "two.sided").

Author(s)

Alan T. Arnholt

References

- Kitchens, L. J. (2003). *Basic Statistics and Data Analysis*. Duxbury.
- Hogg, R. V. and Craig, A. T. (1970). *Introduction to Mathematical Statistics, 3rd ed.* Toronto, Canada: Macmillan.
- Mood, A. M., Graybill, F. A. and Boes, D. C. (1974). *Introduction to the Theory of Statistics, 3rd ed.* New York: McGraw-Hill.
- Snedecor, G. W. and Cochran, W. G. (1980). *Statistical Methods, 7th ed.* Ames, Iowa: Iowa State University Press.

See Also

[z.test](#), [tsum.test](#)

Examples

```
zsum.test(mean.x=56/30,sigma.x=2, n.x=30, alternative="greater", mu=1.8)
# Example 9.7 part a. from PASWR.
x <- rnorm(12)
zsum.test(mean(x),sigma.x=1,n.x=12)
# Two-sided one-sample z-test where the assumed value for
# sigma.x is one. The null hypothesis is that the population
# mean for 'x' is zero. The alternative hypothesis states
# that it is either greater or less than zero. A confidence
# interval for the population mean will be computed.
# Note: returns same answer as:
z.test(x,sigma.x=1)
#
x <- c(7.8, 6.6, 6.5, 7.4, 7.3, 7.0, 6.4, 7.1, 6.7, 7.6, 6.8)
y <- c(4.5, 5.4, 6.1, 6.1, 5.4, 5.0, 4.1, 5.5)
zsum.test(mean(x), sigma.x=0.5, n.x=11 ,mean(y), sigma.y=0.5, n.y=8, mu=2)
# Two-sided standard two-sample z-test where both sigma.x
# and sigma.y are both assumed to equal 0.5. The null hypothesis
# is that the population mean for 'x' less that for 'y' is 2.
# The alternative hypothesis is that this difference is not 2.
# A confidence interval for the true difference will be computed.
# Note: returns same answer as:
z.test(x, sigma.x=0.5, y, sigma.y=0.5)
#
zsum.test(mean(x), sigma.x=0.5, n.x=11, mean(y), sigma.y=0.5, n.y=8,
conf.level=0.90)
# Two-sided standard two-sample z-test where both sigma.x and
# sigma.y are both assumed to equal 0.5. The null hypothesis
# is that the population mean for 'x' less that for 'y' is zero.
# The alternative hypothesis is that this difference is not
# zero. A 90% confidence interval for the true difference will
# be computed. Note: returns same answer as:
```



```
z.test(x, sigma.x=0.5, y, sigma.y=0.5, conf.level=0.90)  
rm(x, y)
```

Index

* datasets

Abbey, 9
Abc, 10
Abilene, 10
Ability, 11
Abortion, 12
Absent, 13
Achieve, 14
Adsales, 14
Aggress, 15
Aid, 16
Aids, 17
Airdisasters, 18
Airline, 19
Alcohol, 20
Allergy, 20
Anesthet, 21
Anxiety, 22
Apolipop, 22
Append, 23
Appendec, 24
Aptitude, 25
Archaeo, 25
Arthriti, 26
Artifici, 27
Asprin, 28
Asthmati, 28
Attorney, 29
Autogear, 30
Backtoback, 31
Bbsalaries, 31
Bigten, 32
Biology, 33
Birth, 33
Blackedu, 34
Blood, 35
Board, 36
Bones, 37
Books, 37
Bookstor, 38
Brain, 39
Bumpers, 40
Bus, 41
Bypass, 41
Cabinets, 42
Cancer, 43
Carbon, 44
Cat, 45
Censored, 45
Challeng, 46
Chemist, 47
Chesapea, 48
Chevy, 48
Chicken, 49
Chipavg, 50
Chips, 51
Cigar, 52
Cigarette, 52
Citrus, 55
Clean, 55
Coaxial, 56
Coffee, 57
Coins, 57
Commute, 59
Concept, 60
Concrete, 60
Corn, 61
Correlat, 62
Counsel, 62
Cpi, 63
Crime, 64
Darwin, 65
Dealers, 66
Defectiv, 66
Degree, 67
Delay, 68
Depend, 69
Detroit, 69

Develop, [70](#)
Devmath, [71](#)
Dice, [71](#)
Diesel, [72](#)
Diplomat, [73](#)
Disposal, [74](#)
Dogs, [75](#)
Domestic, [76](#)
Dopamine, [77](#)
Dowjones, [77](#)
Drink, [78](#)
Drug, [79](#)
Dyslexia, [80](#)
Earthqk, [81](#)
Educat, [82](#)
Eggs, [83](#)
Elderly, [84](#)
Energy, [85](#)
Engineer, [86](#)
Entrance, [86](#)
Epaminicompat, [87](#)
Epatwoseater, [88](#)
Executiv, [89](#)
Exercise, [90](#)
Fabric, [90](#)
Faithful, [91](#)
Family, [92](#)
Ferraro1, [93](#)
Ferraro2, [93](#)
Fertility, [94](#)
Firstchi, [95](#)
Fish, [96](#)
Fitness, [97](#)
Florida2000, [98](#)
Fluid, [99](#)
Food, [100](#)
Framingh, [100](#)
Freshman, [101](#)
Funeral, [102](#)
Galaxie, [103](#)
Gallup, [103](#)
Gasoline, [104](#)
German, [105](#)
Golf, [106](#)
Governor, [107](#)
Gpa, [108](#)
Grades, [109](#)
Graduate, [109](#)
Greenriv, [110](#)
Grnriv2, [111](#)
Groupabc, [111](#)
Groups, [112](#)
Gym, [113](#)
Habits, [113](#)
Haptoglo, [114](#)
Hardware, [115](#)
Hardwood, [116](#)
Heat, [117](#)
Heating, [118](#)
Hodgkin, [118](#)
Homes, [119](#)
Homework, [120](#)
Honda, [121](#)
Hostile, [122](#)
Housing, [122](#)
Hurrican, [123](#)
Iceberg, [124](#)
Income, [125](#)
Independent, [126](#)
Indian, [127](#)
Indiapol, [128](#)
Indy500, [128](#)
Inflatio, [129](#)
Inletoil, [130](#)
Inmate, [131](#)
Inspect, [132](#)
Insulate, [133](#)
Iqgpa, [134](#)
Irises, [134](#)
Jdpower, [135](#)
Jobsat, [136](#)
Kidsmoke, [137](#)
Kilowatt, [138](#)
Kinder, [138](#)
Laminect, [139](#)
Lead, [140](#)
Leader, [141](#)
Lethal, [141](#)
Life, [142](#)
Lifespan, [143](#)
Ligntmonth, [143](#)
Lodge, [144](#)
Longtail, [145](#)
Lowabil, [146](#)
Magnesiu, [146](#)
Malpract, [147](#)

- Manager, 148
Marked, 148
Math, 149
Mathcomp, 150
Mathpro, 151
Maze, 152
Median, 152
Mental, 153
Mercury, 154
Metrent, 154
Miller, 155
Miller1, 156
Moisture, 156
Monoxide, 157
Movie, 158
Music, 159
Name, 160
Nascar, 161
Nervous, 161
Newsstand, 162
Nfldraf2, 163
Nfldraft, 163
Nicotine, 164
Orange, 168
Orioles, 168
Oxytocin, 169
Parented, 170
Patrol, 171
Pearson, 172
Phone, 172
Poison, 173
Politic, 174
Pollutio, 175
Porosity, 175
Poverty, 176
Precinct, 177
Prejudic, 178
Presiden, 178
Press, 179
Prognost, 180
Program, 181
Psat, 181
Psych, 182
Puerto, 183
Quail, 183
Quality, 184
Rainks, 185
Randd, 186
Rat, 186
Ratings, 187
Reaction, 188
Reading, 189
Readiq, 189
Referend, 190
Region, 191
Register, 192
Rehab, 192
Remedial, 193
Rentals, 194
Repair, 195
Retail, 195
Ronbrow1, 196
Ronbrow2, 197
Rural, 197
Salary, 198
Salinity, 199
Sat, 199
Saving, 200
Scales, 201
Schizop2, 202
Schizoph, 203
Seatbelt, 203
Selfdefe, 204
Senior, 205
Sentence, 206
Shkdrug, 206
Shock, 207
Shoplift, 208
Short, 209
Shuttle, 209
Simpson, 213
Situp, 214
Skewed, 214
Skin, 215
Slc, 216
Smokyph, 216
Snore, 217
Snow, 218
Soccer, 219
Social, 220
Sophomor, 220
South, 221
Speed, 222
Spellers, 222
Spelling, 223
Sports, 224

- Spouse, [224](#)
- Stable, [226](#)
- Stamp, [226](#)
- Statclas, [227](#)
- Statelaw, [228](#)
- Statisti, [229](#)
- Step, [229](#)
- Stress, [230](#)
- Study, [231](#)
- Submarin, [232](#)
- Subway, [233](#)
- Sunspot, [233](#)
- Superbowl, [234](#)
- Supercar, [235](#)
- Tablrock, [236](#)
- Teacher, [237](#)
- Tenness, [238](#)
- Tensile, [239](#)
- Test1, [240](#)
- Thermal, [240](#)
- Tiaa, [241](#)
- Ticket, [242](#)
- Toaster, [242](#)
- Tonsils, [243](#)
- Tort, [244](#)
- Toxic, [245](#)
- Track, [246](#)
- Track15, [247](#)
- Treatments, [247](#)
- Trees, [248](#)
- Trucks, [249](#)
- Tv, [253](#)
- Twin, [253](#)
- Undergrad, [254](#)
- Vacation, [255](#)
- Vaccine, [256](#)
- Vehicle, [257](#)
- Verbal, [257](#)
- Victoria, [258](#)
- Viscosit, [259](#)
- Visual, [260](#)
- Vocab, [260](#)
- Wastewat, [261](#)
- Weather94, [262](#)
- Wheat, [263](#)
- Windmill, [264](#)
- Window, [264](#)
- Wins, [265](#)
- Wool, [266](#)
- Yearsunspot, [267](#)
- * **distribution**
 - CIsim, [53](#)
 - Combinations, [58](#)
 - normarea, [165](#)
 - ntester, [167](#)
 - SRS, [225](#)
- * **htest**
 - tsum.test, [249](#)
 - z.test, [267](#)
 - zsum.test, [270](#)
- * **univar**
 - EDA, [81](#)
 - nsize, [166](#)
- Abbey, [9](#)
- Abc, [10](#)
- Abilene, [10](#)
- Ability, [11](#)
- Abortion, [12](#)
- Absent, [13](#)
- Achieve, [14](#)
- Adsales, [14](#)
- Aggress, [15](#)
- Aid, [16](#)
- Aids, [17](#)
- Airdisasters, [18](#)
- Airline, [19](#)
- Alcohol, [20](#)
- Allergy, [20](#)
- Anesthet, [21](#)
- Anxiety, [22](#)
- Apolipop, [22](#)
- Append, [23](#)
- Appendec, [24](#)
- Aptitude, [25](#)
- Archaeo, [25](#)
- Arthriti, [26](#)
- Artifici, [27](#)
- Asprin, [28](#)
- Asthmati, [28](#)
- Attorney, [29](#)
- Autogear, [30](#)
- Backtoback, [31](#)
- Bbsalaries, [31](#)
- Bigten, [32](#)
- Biology, [33](#)

- Birth, [33](#)
Blackedu, [34](#)
Blood, [35](#)
Board, [36](#)
Bones, [37](#)
Books, [37](#)
Bookstor, [38](#)
Brain, [39](#)
Bumpers, [40](#)
Bus, [41](#)
Bypass, [41](#)

Cabinets, [42](#)
Cancer, [43](#)
Carbon, [44](#)
Cat, [45](#)
Censored, [45](#)
Challeng, [46](#)
Chemist, [47](#)
Chesapea, [48](#)
Chevy, [48](#)
Chicken, [49](#)
Chipavg, [50](#)
Chips, [51](#)
Cigar, [52](#)
Cigarette, [52](#)
CIsim, [53](#)
Citrus, [55](#)
Clean, [55](#)
Coaxial, [56](#)
Coffee, [57](#)
Coins, [57](#)
Combinations, [58](#), [225](#)
Commute, [59](#)
Concept, [60](#)
Concrete, [60](#)
Corn, [61](#)
Correlat, [62](#)
Counsel, [62](#)
Cpi, [63](#)
Crime, [64](#)

Darwin, [65](#)
Dealers, [66](#)
Defectiv, [66](#)
Degree, [67](#)
Delay, [68](#)
Depend, [69](#)
Detroit, [69](#)

Develop, [70](#)
Devmath, [71](#)
Dice, [71](#)
Diesel, [72](#)
Diplomat, [73](#)
Disposal, [74](#)
Dogs, [75](#)
Domestic, [76](#)
Dopamine, [77](#)
Dowjones, [77](#)
Drink, [78](#)
Drug, [79](#)
Dyslexia, [80](#)

Earthqk, [81](#)
EDA, [81](#)
Educat, [82](#)
Eggs, [83](#)
Elderly, [84](#)
Energy, [85](#)
Engineer, [86](#)
Entrance, [86](#)
Epaminicompact, [87](#)
Epatwoseater, [88](#)
Executiv, [89](#)
Exercise, [90](#)

Fabric, [90](#)
Faithful, [91](#)
Family, [92](#)
Ferraro1, [93](#)
Ferraro2, [93](#)
Fertility, [94](#)
Firstchi, [95](#)
Fish, [96](#)
Fitness, [97](#)
Florida2000, [98](#)
Fluid, [99](#)
Food, [100](#)
Framingh, [100](#)
Freshman, [101](#)
Funeral, [102](#)

Galaxie, [103](#)
Gallup, [103](#)
Gasoline, [104](#)
German, [105](#)
Golf, [106](#)
Governor, [107](#)

- Gpa, 108
Grades, 109
Graduate, 109
Greenriv, 110
Gnriv2, 111
Groupabc, 111
Groups, 112
Gym, 113
- Habits, 113
Haptoglo, 114
Hardware, 115
Hardwood, 116
Heat, 117
Heating, 118
Hodgkin, 118
Homes, 119
Homework, 120
Honda, 121
Hostile, 122
Housing, 122
Hurrican, 123
- Iceberg, 124
Income, 125
Independent, 126
Indian, 127
Indiapol, 128
Indy500, 128
Inflatio, 129
Inletoil, 130
Inmate, 131
Inspect, 132
Insulate, 133
Iqgpa, 134
Irises, 134
- Jdpower, 135
Jobsat, 136
- Kidsmoke, 137
Kilowatt, 138
Kinder, 138
- Laminect, 139
Lead, 140
Leader, 141
Lethal, 141
Life, 142
- Lifespan, 143
Ligntmonth, 143
Lodge, 144
Longtail, 145
Lowabil, 146
- Magnesiu, 146
Malpract, 147
Manager, 148
Marked, 148
Math, 149
Mathcomp, 150
Mathpro, 151
Maze, 152
Median, 152
Mental, 153
Mercury, 154
Metrent, 154
Miller, 155
Miller1, 156
Moisture, 156
Monoxide, 157
Movie, 158
Music, 159
- Name, 160
Nascar, 161
Nervous, 161
Newsstand, 162
Nfldraf2, 163
Nfldraft, 163
Nicotine, 164
normarea, 165
nsize, 166
ntester, 167
- Orange, 168
Orioles, 168
Oxytocin, 169
- Parented, 170
Patrol, 171
Pearson, 172
Phone, 172
Poison, 173
Politic, 174
Pollutio, 175
Porosity, 175
Poverty, 176

- Precinct, [177](#)
Prejudic, [178](#)
Presiden, [178](#)
Press, [179](#)
Prognost, [180](#)
Program, [181](#)
Pstat, [181](#)
Psych, [182](#)
Puerto, [183](#)

Quail, [183](#)
Quality, [184](#)

Rains, [185](#)
Randd, [186](#)
Rat, [186](#)
Ratings, [187](#)
Reaction, [188](#)
Reading, [189](#)
Readiq, [189](#)
Referend, [190](#)
Region, [191](#)
Register, [192](#)
Rehab, [192](#)
Remedial, [193](#)
Rentals, [194](#)
Repair, [195](#)
Retail, [195](#)
Ronbrown1, [196](#)
Ronbrown2, [197](#)
Rural, [197](#)

Salary, [198](#)
Salinity, [199](#)
Sat, [199](#)
Saving, [200](#)
Scales, [201](#)
Schizop2, [202](#)
Schizop, [203](#)
Seatbelt, [203](#)
Selfdefe, [204](#)
Senior, [205](#)
Sentence, [206](#)
Shkdrug, [206](#)
Shock, [207](#)
Shoplift, [208](#)
Short, [209](#)
Shuttle, [209](#)
SIGN.test, [210](#)

Simpson, [213](#)
Situp, [214](#)
Skewed, [214](#)
Skin, [215](#)
Slc, [216](#)
Smokyph, [216](#)
Snore, [217](#)
Snow, [218](#)
Soccer, [219](#)
Social, [220](#)
Sophomor, [220](#)
South, [221](#)
Speed, [222](#)
Spellers, [222](#)
Spelling, [223](#)
Sports, [224](#)
Spouse, [224](#)
SRS, [58](#), [225](#)
Stable, [226](#)
Stamp, [226](#)
Statclas, [227](#)
Statelaw, [228](#)
Statisti, [229](#)
Step, [229](#)
Stress, [230](#)
Study, [231](#)
Submarin, [232](#)
Subway, [233](#)
Sunspot, [233](#)
Superbowl, [234](#)
Supercar, [235](#)

Tablrock, [236](#)
Teacher, [237](#)
Tenness, [238](#)
Tensile, [239](#)
Test1, [240](#)
Thermal, [240](#)
Tiaa, [241](#)
Ticket, [242](#)
Toaster, [242](#)
Tonsils, [243](#)
Tort, [244](#)
Toxic, [245](#)
Track, [246](#)
Track15, [247](#)
Treatments, [247](#)
Trees, [248](#)
Trucks, [249](#)

tsum.test, [212](#), [249](#), [269](#), [272](#)

Tv, [253](#)

Twin, [253](#)

Undergrad, [254](#)

Vacation, [255](#)

Vaccine, [256](#)

Vehicle, [257](#)

Verbal, [257](#)

Victoria, [258](#)

Viscosit, [259](#)

Visual, [260](#)

Vocab, [260](#)

Wastewat, [261](#)

Weather94, [262](#)

Wheat, [263](#)

Windmill, [264](#)

Window, [264](#)

Wins, [265](#)

Wool, [266](#)

Yearsunspot, [267](#)

z.test, [212](#), [252](#), [267](#), [272](#)

zsum.test, [212](#), [252](#), [269](#), [270](#)