

Package ‘LearningRlab’

July 21, 2025

Version 2.4

Date 2023-8-30

Title Statistical Learning Functions

Author Carlos Javier Hellin Asensio [aut, cre],
Jose Manuel Gomez Caceres [aut],
Dennis Monheimius [aut],
Eduardo Benito [aut],
Juan Jose Cuadrado [aut],
Universidad de Alcala de Henares [aut]

Maintainer Carlos Javier Hellin Asensio <carlos.hellin@uah.es>

Depends magick, crayon

Suggests knitr, rmarkdown

Description Aids in learning statistical functions incorporating the result of calculus done with each function and how they are obtained, that is, which equation and variables are used. Also for all these equations and their related variables detailed explanations and interactive exercises are also included. All these characteristics allow to the package user to improve the learning of statistics basics by means of their use.

License Unlimited

VignetteBuilder knitr

NeedsCompilation yes

Type Package

Repository CRAN

Date/Publication 2023-08-30 22:10:03 UTC

RoxygenNote 7.1.1

Encoding UTF-8

Contents

averageDeviation_	3
binomial_	4
chisquared_	6

covariance_	7
cv_	8
drawVector	9
explain.absolute_acum_frecuency	10
explain.absolute_frecuency	11
explain.averageDeviation	12
explain.binomial	13
explain.chisquared	14
explain.covariance	15
explain.cv	16
explain.fisher	17
explain.geometricMean	18
explain.harmonicMean	19
explain.laplace	20
explain.mean	21
explain.median	22
explain.mode	23
explain.normal	24
explain.pearson	25
explain.percentile	26
explain.poisson	27
explain.quartile	28
explain.relative_acum_frecuency	29
explain.relative_frecuency	30
explain.standardDeviation	31
explain.tstudent	32
explain.variance	33
fisher_	34
frecuency_abs	35
frecuency_absolute_acum	36
frecuency_relative	37
frecuency_relative_acum	38
geometricMean_	40
getUserAction	41
harmonicMean_	41
initImages	42
interactive.absolute_acum_frecuency	43
interactive.absolute_frecuency	44
interactive.averageDeviation	45
interactive.binomial	46
interactive.chisquared	46
interactive.covariance	47
interactive.cv	48
interactive.fisher	49
interactive.geometricMean	49
interactive.harmonicMean	50
interactive.laplace	51
interactive.mean	52

averageDeviation_3

interactive.median

interactive.mode

interactive.normal

interactive.pearson

interactive.percentile

interactive.poisson

interactive.quartile

interactive.relative_acum_frecuency

interactive.relative_frecuency

interactive.standardDeviation

interactive.tstudent

interactive.variance

laplace_

LearningRlab

meanC

mean_

median_

mode_

normal_

pearson_

percentile_

poisson_

quartile_

standardDeviation_

tstudent_

variance_

52

53

54

55

55

56

57

58

58

59

60

61

61

63

63

64

65

67

68

69

70

71

72

73

74

75

Index77

averageDeviation_	Average Absolute Deviation Function
-------------------	-------------------------------------

Description

This function calculates the average absolute deviation of a numbers vector.

Usage

averageDeviation_(x)

Arguments

x Should be a numbers vector

Details

To calculate the average deviation, the user should give a numbers vector. The result is the sum of the differences in absolute value between each vector element and the mean, divided by the number of elemets. The average absolute deviation formule is the following:

$$\sigma = \frac{\sum_i^N |X_i - \bar{X}|}{N}$$

®LearningRlab

Value

Numeric, the average absolute deviation of the numbers vector.

Note

A vector is created by c(), like c(1,2,3,4,5) creates a vector with the numbers: 1,2,3,4,5

Author(s)

Dennis Monheimius, <dennis.monhemimius@edu.uah.es>
 Eduardo Benito, <eduardo.benito@edu.uah.es>
 Juan Jose Cuadrado, <jjcg@uah.es>
 Universidad de Alcala de Henares

Examples

```
#data creation
data <- c(1:20)
result = averageDeviation_(data)
```

binomial_

Binomial Distribution Calculus Function

Description

This function calculates the binomial distribution of experiment.

Usage

```
binomial_(n,x,p)
```

Arguments

x	Should be a numbers.
n	Should be a numbers.
p	Should be a numbers.

Details

To calculate the binomial distribution, the user should give three number (the number of trials, probability of success and binomial random variable). The result is a discrete probability distribution that counts the number of successes in a sequence of n independent Bernoulli trials with a fixed probability p of occurrence of success between trials. The binomial distribution formule is the following:

$$\text{Binomial Distribution} = \frac{n!}{x!(n-x)!} p^x (1-p)^{(n-x)}$$

LearningRlab ®

Value

Numeric, the binomial distribution of three variables.

Note

Each variable is a number. Example: n <- 3 | x <- 2 | p <- 0.7

Author(s)

Jose Manuel Gomez Caceres, <josemanuel.gomezc@edu.uah.es>
Juan Jose Cuadrado, <jjcg@uah.es>
Universidad de Alcala de Henares

Examples

```
#data creation
n = 3
x = 2
p = 0.7
binomial_(n,x,p)
```

chisquared_

*Chisquared Distribution Calculus Function***Description**

This function calculates the chisquared distribution of two vectors of numbers.

Usage

```
chisquared_(x,y)
```

Arguments

x	Should be a vector.
y	Should be a vector.

Details

To calculate the chisquared distribution, the user should give two vectors of numbers. The result is a sum of the squares of k independent standard normal random variables. The chisquared distribution formule is the following:

$$\textbf{Chi-Squared} = \frac{\sum (f_e - f_o)^2}{f_e}$$

LearningRlab ®

Value

Numeric, the chisquared distribution of two vectors of numbers.

Note

A vector is created by c(), like c(1,2,3,4,5) creates a vector with the numbers: 1,2,3,4,5

Author(s)

Jose Manuel Gomez Caceres, <josemanuel.gomezc@edu.uah.es>
 Juan Jose Cuadrado, <jjcg@uah.es>
 Universidad de Alcala de Henares

Examples

```
#data creation
data = c(1,4,3,3,2,5,7,12,1,2,3,12)
data2 = c(1,2,4,4,6,5,11,2,10,5,6,1)
chisquared_(data, data2)
```

covariance_

*Coariance Calculus Function***Description**

This function calculates the covariance of two vectors of numbers.

Usage

```
covariance_(x,y)
```

Arguments

x	Should be a vector
y	Should be a vector

Details

To calculate the covariance, the user should give two vectors of numbers. The result is a measure of the joint variability of two vectors of numbers. The covariance formule is the following:

$$\textbf{Covariance} = \frac{\sum_1^n (x_i - \bar{x})(y_i - \bar{y})}{n}$$

LearningRlab ®**Value**

Numeric, the covariance of two vectors of numbers.

Note

A vector is created by c(), like c(1,2,3,4,5) creates a vector with the numbers: 1,2,3,4,5

Author(s)

Dennis Monheimius, <dennis.monhemimius@edu.uah.es>
 Eduardo Benito, <eduardo.benito@edu.uah.es>
 Juan Jose Cuadrado, <jjcg@uah.es>
 Universidad de Alcala de Henares

Examples

```
#data creation
data = c(1,4,3,3,2,5,7,12,1,2,3,12)
data2 = c(1,2,4,4,6,5,11,2,10,5,6,1)
covariance_(data, data2)
```

cv_

*Coefficient of Variation Calculus Function***Description**

This function calculates the coefficient of variation of a numbers vector.

Usage

```
cv_(x)
```

Arguments

x Should be a numbers vector

Details

To calculate the coefficient of variation, the user should give a numbers vector. The result is defined as the ratio of the standard deviation to the mean. The coefficient of variation formule is the following:

$$CV = \frac{\text{std dev}}{\text{mean}} = \frac{s}{\bar{y}}$$

LearningRlab ®

Value

Numeric, the coefficient of variation of the numbers vector.

Note

A vector is created by c(), like c(1,2,3,4,5) creates a vector with the numbers: 1,2,3,4,5

Author(s)

Jose Manuel Gomez Caceres, <josemanuel.gomezc@edu.uah.es>
Juan Jose Cuadrado, <jjcg@uah.es>
Universidad de Alcala de Henares

Examples

```
#data creation  
data = c(1,4,3,3,2,5,7,12,1,2,3,12)  
  
cv_(data)
```

drawVector

Draw Vector Function

Description

This function prints all the elements of a vector

Usage

```
drawVector(buffer)
```

Arguments

buffer A vector of elements

Value

There isn't return value, prints on screen

Note

A vector is created by c(), like c(1,2,3,4,5) creates a vector with the numbers: 1,2,3,4,5 or c(true,false,false) creates a vector with the booleans: true, false, true

Author(s)

Dennis Monheimius, <dennis.monhemimius@edu.uah.es>
Eduardo Benito, <eduardo.benito@edu.uah.es>
Juan Jose Cuadrado, <jjcg@uah.es>
Universidad de Alcala de Henares

Examples

```
{  
  #data creation  
  data <- c(1:12)  
  drawVector(data)  
}
```

```
explain.absolute_acum_frecuency
```

Absolute Accumulated Frequency Calculus Explained

Description

Step by step demonstration of the absolute accumulated frequency calculus

Usage

```
explain.absolute_acum_frecuency(v, x)
```

Arguments

v	Should be a vector
x	Should be a number

Details

To calculate the absolute accumulated frequency, the user should give a vector and a number. We can saw the absolute accumulated frequency formule in the frecuency_acum_absolute help document.

Value

A demonstration of the calculus process

Note

A vector is created by c(), like c(1,2,3,4,5) creates a vector with the numbers: 1,2,3,4,5

Author(s)

Dennis Monheimius, <dennis.monhemimius@edu.uah.es>
Eduardo Benito, <eduardo.benito@edu.uah.es>
Juan Jose Cuadrado, <jjcg@uah.es>
Universidad de Alcala de Henares

Examples

```
{  
  #data creation  
  data <- c(1,2,2,5,10,4,2)  
  value = 2  
  #function execution  
  explain.absolute_acum_frecuency(data, value)  
}
```

explain.absolute_frecuency

Absolute Frecuency Calculus Explained

Description

Step by step demonstration of the absolute frecuency calculus

Usage

explain.absolute_frecuency(v,x)

Arguments

v	Should be a vector
x	Should be a number

Details

To calculate the absolute frecuency, the user should give a vector and a number. We can saw the absolute frecuency formule in the frecuency_abs help document.

Value

A demonstration of the calculus process

Note

A vector is created by c(), like c(1,2,3,4,5) creates a vector with the numbers: 1,2,3,4,5

Author(s)

Dennis Monheimius, <dennis.monhemimius@edu.uah.es>
Eduardo Benito, <eduardo.benito@edu.uah.es>
Juan Jose Cuadrado, <jjcg@uah.es>
Universidad de Alcala de Henares

Examples

```
{  
  #data creation  
  data <- c(1,2,2,5,10,4,2)  
  value = 2  
  #function execution  
  explain.absolute_frecuency(data, value)  
}
```

explain.averageDeviation

Average Absolute Deviation Function Explained

Description

Step by step demonstration of the average absolute deviation calculus.

Usage

```
explain.averageDeviation(x)
```

Arguments

x Should be a numbers vector

Details

To calculate the average absolute deviation, the user should give a numbers vector. The result is the explained process to calculate the average absolute deviation, with the data of the dataset provided like argument. We can saw the average absolute deviation formule in the averageDeviation_ help document.

Value

Numeric, the average absolute deviation of the numbers vector.

Note

A vector is created by c(), like c(1,2,3,4,5) creates a vector with the numbers: 1,2,3,4,5

Author(s)

Dennis Monheimius, <dennis.monhemimius@edu.uah.es>
Eduardo Benito, <eduardo.benito@edu.uah.es>
Juan Jose Cuadrado, <jjcg@uah.es>
Universidad de Alcala de Henares

Examples

```
#data creation
data <- c(7,2,5,7,1,4,12)

explain.averageDeviation(data)
```

`explain.binomial`*Binomial Distribution Function Explained*

Description

Step by step demonstration of the binomial distribution calculus.

Usage

```
explain.binomial(n,x,p)
```

Arguments

x	Should be a numbers.
n	Should be a numbers.
p	Should be a numbers.

Details

To calculate the binomial distribution, the user should give three number (the number of trials, probability of success and binomial random variable). The result is a discrete probability distribution that counts the number of successes in a sequence of n independent Bernoulli trials with a fixed probability p of occurrence of success between trials. We can saw the binomial distribution formule in the binomial_ help document.

Value

Numeric result and the process of this calculus explained.

Note

Each variable is a number. Example: n <- 3 | x <- 2 | p <- 0.7

Author(s)

Jose Manuel Gomez Caceres, <josemanuel.gomezc@edu.uah.es>
Juan Jose Cuadrado, <jjcg@uah.es>
Universidad de Alcala de Henares

Examples

```
#data creation
n = 3
x = 2
p = 0.7

explain.binomial(n,x,p)
```

explain.chisquared	<i>Chisquared Distribution Function Explained</i>
--------------------	---

Description

Step by step demonstration of the chisquared distribution calculus.

Usage

```
explain.chisquared(x,y)
```

Arguments

x	Should be a vector.
y	Should be a vector.

Details

To calculate the chisquared distribution, the user should give two vectors of numbers. The result is a sum of the squares of k independent standard normal random variables. We can saw the chisquared distribution formule in the chisquared_ help document.

Value

Numeric result and the process of this calculus explained.

Note

A vector is created by c(), like c(1,2,3,4,5) creates a vector with the numbers: 1,2,3,4,5

Author(s)

Jose Manuel Gomez Caceres, <josemanuel.gomezc@edu.uah.es>
Juan Jose Cuadrado, <jjcg@uah.es>
Universidad de Alcala de Henares

Examples

```
#data creation
data <- c(10,4,5,7,3,4,1)
data2 <- c(1,8,3,4,4,5,7)

explain.chisquared(data, data2)
```

explain.covariance	<i>Covariance Function Explained</i>
--------------------	--------------------------------------

Description

Step by step demonstration of the covariance calculus.

Usage

```
explain.covariance(x,y)
```

Arguments

x	Should be a vector
y	Should be a vector

Details

To calculate the covariance, the user should give two vectors of numbers. The result is the explained process to calculate the covariance, with the data of the datasets provided like argument. We can saw the harmonic mean formule in the covariance_ help document.

Value

Numeric result and the process of this calculus explained.

Note

A vector is created by c(), like c(1,2,3,4,5) creates a vector with the numbers: 1,2,3,4,5

Author(s)

Jose Manuel Gomez Caceres, <josemanuel.gomezc@edu.uah.es>
Juan Jose Cuadrado, <jjcg@uah.es>
Universidad de Alcala de Henares

Examples

```
#data creation
data <- c(10,4,5,7,3,4,1)
data2 <- c(1,8,3,4,4,5,7)

explain.covariance(data, data2)
```

explain.cv

Coefficient of Variation Function Explained

Description

Step by step demonstration of the coefficient of variation calculus.

Usage

```
explain.cv(x)
```

Arguments

x Should be a numbers vector

Details

To calculate the coefficient of variation, the user should give a numbers vector. The result is defined as the ratio of the standard deviation to the mean. We can saw the coefficient of variation formule in the cv_ help document.

Value

Numeric result and the process of this calculus explained.

Note

A vector is created by c(), like c(1,2,3,4,5) creates a vector with the numbers: 1,2,3,4,5

Author(s)

Jose Manuel Gomez Caceres, <josemanuel.gomezc@edu.uah.es>
Juan Jose Cuadrado, <jjcg@uah.es>
Universidad de Alcala de Henares

Examples

```
#data creation
data <- c(10,4,5,7,3,4,1)

explain.cv(data)
```

`explain.fisher`*F Fisher Distribution Function Explained*

Description

Step by step demonstration of the fisher distribution calculus.

Usage

```
explain.fisher(x,y)
```

Arguments

x	Should be a vector
y	Should be a vector

Details

To calculate the fisher distribution, the user should give two vectors of numbers. The result is a continuous probability distribution that arises frequently as the null distribution of a test statistic. We can saw fisher distribution formule in the fisher_ help document.

Value

Numeric result and the process of this calculus explained.

Note

A vector is created by `c()`, like `c(1,2,3,4,5)` creates a vector with the numbers: 1,2,3,4,5

Author(s)

Jose Manuel Gomez Caceres, <josemanuel.gomezc@edu.uah.es>
Juan Jose Cuadrado, <jjcg@uah.es>
Universidad de Alcala de Henares

Examples

```
#data creation
data <- c(10,4,5,7,3,4,1)
data2 <- c(1,8,3,4,4,5,7)

explain.fisher(data, data2)
```

explain.geometricMean *Geometric Mean Function Explained*

Description

Step by step demonstration of the geometric mean calculus.

Usage

```
explain.geometricMean(x)
```

Arguments

x Should be a numbers vector

Details

To calculate the geometric mean of a dataset, the user should give a vector. The result is the explained process to calculate the geometric mean, with the data of the dataset provided like argument. We can saw the geometric mean formule in the `geometricMean_` help document.

Value

Numeric result and the process of this calculus explained.

Note

A vector is created by `c()`, like `c(1,2,3,4,5)` creates a vector with the numbers: 1,2,3,4,5

Author(s)

Dennis Monheimius, <dennis.monhemimius@edu.uah.es>
Eduardo Benito, <eduardo.benito@edu.uah.es>
Juan Jose Cuadrado, <jjcg@uah.es>
Universidad de Alcala de Henares

Examples

```
{  
  #data creation  
  data <- c(5,21,12,7,3,9,1)  
  
  explain.geometricMean(data)  
}
```

explain.harmonicMean *Harmonic Mean Function Explained*

Description

Step by step demonstration of the harmonic mean calculus.

Usage

```
explain.harmonicMean(x)
```

Arguments

x Should be a numbers vector

Details

To calculate the harmonic mean, the user should give a numbers vector. The result is the explained process to calculate the harmonic mean, with the data of the dataset provided like argument. We can see the harmonic mean formula in the harmonicMean_ help document.

Value

Numeric result and the process of this calculus explained.

Note

A vector is created by c(), like c(1,2,3,4,5) creates a vector with the numbers: 1,2,3,4,5

Author(s)

Jose Manuel Gomez Caceres, <josemanuel.gomezc@edu.uah.es>
Juan Jose Cuadrado, <jjcg@uah.es>
Universidad de Alcala de Henares

Examples

```
#data creation  
data <- c(10,4,5,7,3,4,1)  
  
explain.harmonicMean(data)
```

`explain.laplace`*Laplace's Rule Function Explained*

Description

Step by step demonstration of the Laplace's rule calculus.

Usage

```
explain.laplace(x,y)
```

Arguments

x	Should be a vector
y	Should be a vector

Details

To calculate the Laplace's rule, the user should give two vector (unfavorable cases/favorable cases). The result is as the quotient between the number of favorable cases to A, and that of all possible results of the experiment. We can see the Laplace's rule correlation formula in the `laplace_` help document.

Value

Numeric result and the process of this calculus explained.

Note

A vector is created by `c()`, like `c(1,2,3,4,5)` creates a vector with the numbers: 1,2,3,4,5

Author(s)

Jose Manuel Gomez Caceres, <josemanuel.gomezc@edu.uah.es>
Juan Jose Cuadrado, <jjcg@uah.es>
Universidad de Alcala de Henares

Examples

```
#data creation
data <- 3
data2 <- c(1,2,3,4,5,6)

explain.laplace(data, data2)
```

`explain.mean`*Mean Function Explained*

Description

Step by step demonstration of the arithmetic mean calculus.

Usage

```
explain.mean(x)
```

Arguments

`x` Should be a numbers vector

Details

To calculate the arithmetic mean of a dataset, the user should give a vector. The result is the explained process to calculate the arithmetic mean, with the data of the dataset provided like argument. We can saw the arithmetic mean formule in the `mean_` help document.

Value

Numeric result and the process of this calculus explained.

Note

A vector is created by `c()`, like `c(1,2,3,4,5)` creates a vector with the numbers: 1,2,3,4,5

Author(s)

Dennis Monheimius, <dennis.monhemimius@edu.uah.es>
Eduardo Benito, <eduardo.benito@edu.uah.es>
Juan Jose Cuadrado, <jjcg@uah.es>
Universidad de Alcala de Henares

Examples

```
{  
  #data creation  
  data <- c(1,2,2,5,10,4,2)  
  
  explain.mean(data)  
}
```

explain.median*Median Function Explained*

Description

Step by step demonstration of the median calculus.

Usage

```
explain.median(x)
```

Arguments

x Should be a numbers vector

Details

To calculate the median, the user should give a numbers vector. The result is the explained process to calculate the median, with the data of the dataset provided like argument. We can saw the median formule in the median_ help document.

Value

Numeric result and the process of this calculus explained.

Note

A vector is created by c(), like c(1,2,3,4,5) creates a vector with the numbers: 1,2,3,4,5

Author(s)

Dennis Monheimius, <dennis.monhemimius@edu.uah.es>
Eduardo Benito, <eduardo.benito@edu.uah.es>
Juan Jose Cuadrado, <jjcg@uah.es>
Universidad de Alcala de Henares

Examples

```
{  
  #data creation  
  data <- c(1,2,2,5,10,4,2)  
  
  explain.median(data)  
}
```

explain.mode*Mode Function Explained*

Description

Step by step demonstration of the mode calculus.

Usage

```
explain.mode(x)
```

Arguments

x Should be a numbers vector

Details

To calculate the mode, the user should give a numbers vector. The result is the explained process to calculate the mode, with the data of the dataset provided like argument. We can saw the mode formule in the mode_ help document.

Value

Numeric result and the process of this calculus explained.

Note

A vector is created by c(), like c(1,2,3,4,5) creates a vector with the numbers: 1,2,3,4,5

Author(s)

Dennis Monheimius, <dennis.monhemimius@edu.uah.es>
Eduardo Benito, <eduardo.benito@edu.uah.es>
Juan Jose Cuadrado, <jjcg@uah.es>
Universidad de Alcala de Henares

Examples

```
{  
  #data creation  
  data <- c(1,1,2,5,2,3,1,4,1)  
  
  explain.mode(data)  
}
```

`explain.normal`*Normal Distribution Function Explained*

Description

Step by step demonstration of the normal distribution calculus.

Usage

```
explain.normal(x)
```

Arguments

`x` Should be a number.

Details

To calculate the normal distribution, the user should give a number. The result is a type of continuous probability distribution for a real-valued random variable. We can see the normal distribution correlation formulae in the `normal_help` document.

Value

Numeric result and the process of this calculus explained.

Note

The variable is a number. Example: `x <- 0.1`

Author(s)

Jose Manuel Gomez Caceres, <josemanuel.gomezc@edu.uah.es>
Juan Jose Cuadrado, <jjcg@uah.es>
Universidad de Alcala de Henares

Examples

```
#data creation  
x = 0.1  
  
explain.normal(x)
```

explain.pearson*Pearson Correlation Function Explained*

Description

Step by step demonstration of the pearson correlation calculus.

Usage

```
explain.pearson(x,y)
```

Arguments

x	Should be a vector
y	Should be a vector

Details

To calculate the pearson correlation, the user should give two vectors of numbers. The result is the covariance of the two vectors of numbers divided by the product of their standard deviations. We can see the pearson correlation formula in the pearson_ help document.

Value

Numeric result and the process of this calculus explained.

Note

A vector is created by c(), like c(1,2,3,4,5) creates a vector with the numbers: 1,2,3,4,5

Author(s)

Jose Manuel Gomez Caceres, <josemanuel.gomezc@edu.uah.es>
Juan Jose Cuadrado, <jjcg@uah.es>
Universidad de Alcala de Henares

Examples

```
#data creation
data <- c(10,4,5,7,3,4,1)
data2 <- c(1,8,3,4,4,5,7)

explain.pearson(data, data2)
```

explain.percentile	<i>Percentiles Calculus Explained</i>
--------------------	---------------------------------------

Description

Step by step demonstration of the percentiles calculus

Usage

```
explain.percentile(x)
```

Arguments

x	Should be a vector
---	--------------------

Details

To calculate the percentiles, the user should give a vector. We can saw the percentile formule in the percentile_ help document.

Value

A demonstration of the calculus process

Note

A vector is created by c(), like c(1,2,3,4,5) creates a vector with the numbers: 1,2,3,4,5

Author(s)

Dennis Monheimius, <dennis.monhemimius@edu.uah.es>
Eduardo Benito, <eduardo.benito@edu.uah.es>
Juan Jose Cuadrado, <jjcg@uah.es>
Universidad de Alcala de Henares

Examples

```
{  
  #data creation  
  data <- c(1,2,2,5,10,4,2)  
  
  explain.percentile(data)  
}
```

explain.poisson*Poisson Distribution Function Explained*

Description

Step by step demonstration of the Poisson distribution calculus.

Usage

```
explain.poisson(k,lam)
```

Arguments

k	Should be a numbers
lam	Should be a numbers

Details

To calculate the Poisson distribution, the user should give two number (the number of times the phenomenon and the number of occurrences). The result is a discrete probability distribution that expresses, from a mean frequency of occurrence, the probability that a certain number of events will occur during a certain period of time. We can saw the Poisson distribution correlation formule in the poisson_ help document.

Value

Numeric result and the process of this calculus explained.

Note

Each variable is a number. Example: lam <- 2 | k <- 3

Author(s)

Jose Manuel Gomez Caceres, <josemanuel.gomezc@edu.uah.es>
Juan Jose Cuadrado, <jjcg@uah.es>
Universidad de Alcala de Henares

Examples

```
#data creation  
lam = 2  
k = 3  
explain.poisson(k,lam)
```

`explain.quartile`*Quartiles Calculus Explained*

Description

Step by step demonstration of the quartiles calculus

Usage

```
explain.quartile(x)
```

Arguments

`x` Should be a vector

Details

To calculate the quartiles, the user should give a vector. We can saw the quartile formule in the `quartile_help` document.

Value

A demonstration of the calculus process

Note

A vector is created by `c()`, like `c(1,2,3,4,5)` creates a vector with the numbers: 1,2,3,4,5

Author(s)

Dennis Monheimius, <dennis.monhemimius@edu.uah.es>

Eduardo Benito, <eduardo.benito@edu.uah.es>

Juan Jose Cuadrado, <jjcg@uah.es>

Universidad de Alcala de Henares

Examples

```
{  
  #data creation  
  data <- c(1,2,2,5,10,4,2)  
  
  explain.quartile(data)  
}
```

`explain.relative_acum_frecuency`*Relative Accumulated Frequency Calculus Explained*

Description

Step by step demonstration of the relative accumulated frequency calculus

Usage

```
explain.relative_acum_frecuency(v,x)
```

Arguments

<code>v</code>	Should be a vector
<code>x</code>	Should be a numebr of the vector

Details

To calculate the relative accumulated frequency, the user should give a vector and a number. We can saw the relative accumulated frequency formule in the `frecuency_acum_relative` help document.

Value

A demonstration of the calculus process

Note

A vector is created by `c()`, like `c(1,2,3,4,5)` creates a vector with the numbers: 1,2,3,4,5

Author(s)

Dennis Monheimius, <dennis.monhemimius@edu.uah.es>
Eduardo Benito, <eduardo.benito@edu.uah.es>
Juan Jose Cuadrado, <jjcg@uah.es>
Universidad de Alcala de Henares

Examples

```
{  
  #data creation  
  data <- c(1,2,2,5,10,4,2)  
  value = 2  
  #function execution  
  explain.relative_acum_frecuency(data, value)  
}
```

`explain.relative_frequency`*Relative Frequency Calculus Explained*

Description

Step by step demonstration of the relative frequency calculus

Usage

```
explain.relative_frequency(v,x)
```

Arguments

<code>v</code>	Should be a vector
<code>x</code>	Should be a number

Details

To calculate the relative frequency, the user should give a vector and a number. We can see the relative frequency formula in the `frequency_relative` help document.

Value

A demonstration of the calculus process

Note

A vector is created by `c()`, like `c(1,2,3,4,5)` creates a vector with the numbers: 1,2,3,4,5

Author(s)

Dennis Monheimius, <dennis.monheimius@edu.uah.es>
Eduardo Benito, <eduardo.benito@edu.uah.es>
Juan Jose Cuadrado, <jjcg@uah.es>
Universidad de Alcala de Henares

Examples

```
{  
  #data creation  
  data <- c(1,2,2,5,10,4,2)  
  value = 2  
  #function execution  
  explain.relative_frequency(data, value)  
}
```

`explain.standardDeviation`*Standard Deviation Function Explained*

Description

Step by step demonstration of the standard deviation calculus.

Usage

```
explain.standardDeviation(x)
```

Arguments

x	Should be a numbers vector
---	----------------------------

Details

To calculate the standard deviation, the user should give a numbers vector. The result is the explained process to calculate the standard deviation, with the data of the dataset provided like argument. We can saw the standard deviation formule in the standardDeviation_ help document.

Value

Numeric result and the process of this calculus explained.

Note

A vector is created by `c()`, like `c(1,2,3,4,5)` creates a vector with the numbers: 1,2,3,4,5

Author(s)

Dennis Monheimius, <dennis.monhemimius@edu.uah.es>
Eduardo Benito, <eduardo.benito@edu.uah.es>
Juan Jose Cuadrado, <jjcg@uah.es>
Universidad de Alcala de Henares

Examples

```
#data creation
data <- c(1,5,3,7,10,4,2)

explain.standardDeviation(data)
```

explain.tstudent

T-Student Distribution Function Explained

Description

Step by step demonstration of the T-Student distribution calculus.

Usage

```
explain.tstudent(x,u,s,n)
```

Arguments

x	Should be a number
u	Should be a number
s	Should be a number
n	Should be a number

Details

To calculate the T-Student distribution, the user should give four number (sample mean, population mean, population standard deviation and sample size). The result is a probability distribution that arises from the problem of estimating the mean of a normally distributed population when the sample size is small. We can saw the T-Student distribution formule in the tstudent_help document.

Value

Numeric result and the process of this calculus explained.

Note

Each variable is a number. Example: `x <- 2 | y <- 4`

Author(s)

Jose Manuel Gomez Caceres, <josemanuel.gomezc@edu.uah.es>
Juan Jose Cuadrado, <jjcg@uah.es>
Universidad de Alcala de Henares

Examples

```
#data creation
x = 52.9
u = 50
s = 3
n = 10

explain.tstudent(x,u,s,n)
```

explain.variance	<i>Variance Function Explained</i>
------------------	------------------------------------

Description

Step by step demonstration of the variance calculus.

Usage

```
explain.variance(x)
```

Arguments

x	Should be a numbers vector
---	----------------------------

Details

To calculate the variance, the user should give a numbers vector. The result is the explained process to calculate the variance, with the data of the dataset provided like argument. We can saw the variance formule in the variance_ help document.

Value

Numeric result and the process of this calculus explained.

Note

A vector is created by c(), like c(1,2,3,4,5) creates a vector with the numbers: 1,2,3,4,5

Author(s)

Dennis Monheimius, <dennis.monhemimius@edu.uah.es>
Eduardo Benito, <eduardo.benito@edu.uah.es>
Juan Jose Cuadrado, <jjcg@uah.es>
Universidad de Alcala de Henares

Examples

```
#data creation  
data <- c(10,4,5,7,3,4,1)  
  
explain.variance(data)
```

fisher_*F Fisher Distribution Calculus Function*

Description

This function calculates the fisher distribution of a numbers vector.

Usage

fisher_(x,y)

Arguments

x	Should be a vector
y	Should be a vector

Details

To calculate the fisher distribution, the user should give two vectors of numbers. The result is a continuous probability distribution that arises frequently as the null distribution of a test statistic. The fisher distributionformule is the following:

$$\text{Fisher Distribution} = \frac{S_x^2}{S_w^2}$$

LearningRlab®

Value

Numeric, the fisher distribution.

Note

A vector is created by c(), like c(1,2,3,4,5) creates a vector with the numbers: 1,2,3,4,5

Author(s)

Jose Manuel Gomez Caceres, <josemanuel.gomezc@edu.uah.es>
Juan Jose Cuadrado, <jjcg@uah.es>
Universidad de Alcala de Henares

Examples

```
#data creation
x <- c(70,75,74,72,68,59)
y <- c(74,77,70,80,72,76)

fisher_(x,y)
```

frequency_abs	<i>Absolute Frequency Calculus</i>
---------------	------------------------------------

Description

This function calculate the number of times that a specific number appears in the data set.

Usage

```
frequency_abs(v, x)
```

Arguments

v	Should be a vector
x	Should be a number

Details

The absolute frequency formula is the following:

Absolute frequency = number of aparitions of
the examined element

LearningRlab ®

Value

An integer that represents the number of times that the value appears in the vector

Note

A vector is created by c(), like c(1,2,3,4,5) creates a vector with the numbers: 1,2,3,4,5

Author(s)

Dennis Monheimius, <dennis.monhemimius@edu.uah.es>
 Eduardo Benito, <eduardo.benito@edu.uah.es>
 Juan Jose Cuadrado, <jjcg@uah.es>
 Universidad de Alcala de Henares

Examples

```
{
  #data creation
  data = c(1,4,3,3,2,5,7,12,1,2,3,12)
  value = 12
  #function execution
  frequency_abs(data, value)
}
```

frequency_absolute_acum

Accumulated Absolute Frequency Calculus

Description

This function calculate the number of times that a specific number appears in the data set. The value depends on the elements that are lower than itself

Usage

```
frequency_absolute_acum(v,x)
```

Arguments

v	Should be a vector
x	Should be a number

Details

The accumulated absolute frequency formula is the following:

**Absolute
accumulated =
frequency (X)** $\sum F_i \text{ where } i \leq X$

LearningRlab ®

Value

A double that represents the number of times that the value appears in the vector regarding the total of elements

Note

A vector is created by `c()`, like `c(1,2,3,4,5)` creates a vector with the numbers: 1,2,3,4,5

Author(s)

Dennis Monheimius, <dennis.monhemimius@edu.uah.es>

Eduardo Benito, <eduardo.benito@edu.uah.es>

Juan Jose Cuadrado, <jjcg@uah.es>

Universidad de Alcala de Henares

Examples

```
{
  #data creation
  data = c(1,4,3,3,2,5,7,12,1,2,3,12)
  value = 12
  #function execution
  frequency_absolute_acum(data, value)
}
```

frequency_relative	<i>Relative Frequency Calculus</i>
--------------------	------------------------------------

Description

This function calculate the number of times that a specific number appears in the data set divided by the total length of the vector.

Usage

```
frequency_relative(v,x)
```

Arguments

v	Should be a vector
x	Should be a number

Details

The relative frequency formula is the following:

$$\text{Relative frequency} = \frac{\text{absolute frequency}}{\sum \text{all frequencies}}$$

LearningRlab®

Value

A double that represents the number of times that the value appears in the vector regarding the total of elements

Note

A vector is created by `c()`, like `c(1,2,3,4,5)` creates a vector with the numbers: 1,2,3,4,5

Author(s)

Dennis Monheimius, <dennis.monhemimius@edu.uah.es>
 Eduardo Benito, <eduardo.benito@edu.uah.es>
 Juan Jose Cuadrado, <jjcg@uah.es>
 Universidad de Alcala de Henares

Examples

```
{
#data creation
data <- c(1,4,3,3,2,5,7,12,1,2,3,12)
value = 12
frequency_relative(data, value)
}
```

frequency_relative_acum

Accumulated Relative Frequency Calculus

Description

This function calculate the number of times that a specific number appears in the data set divided by the total length of the vector. The value depends on the elements that are lower than itself

Usage

```
frequency_relative_acum(v,x)
```

Arguments

v	Should be a vector
x	Should be a number

Details

The accumulated relative frequency formula is the following:

$$\text{Relative accumulated frequency (X)} = \sum f_i \text{ where } i \leq X$$

LearningRlab ®

Value

A double that represents the number of times that the value appears in the vector regarding the total of elements

Note

A vector is created by `c()`, like `c(1,2,3,4,5)` creates a vector with the numbers: 1,2,3,4,5

Author(s)

Dennis Monheimius, <dennis.monhemimius@edu.uah.es>
Eduardo Benito, <eduardo.benito@edu.uah.es>
Juan Jose Cuadrado, <jjcg@uah.es>
Universidad de Alcala de Henares

Examples

```
{  
  #data creation  
  data = c(1,4,3,3,2,5,7,12,1,2,3,12)  
  value = 12  
  #function execution  
  frequency_relative_acum(data, value)  
}
```

geometricMean_	<i>Geometric Mean Function</i>
----------------	--------------------------------

Description

This function calculates the geometric mean of a numbers vector.

Usage

```
geometricMean_(x)
```

Arguments

x	Should be a numbers vector
---	----------------------------

Details

To calculate the geometric mean of a dataset, the user should give a numbers vector. The result is the product of all vector elements raise to 1 divided by the number of elements. The arithmetic mean formule is the following:

$$MG = \sqrt[n]{(X_1)(X_2)(X_3)...(X_n)}$$

©LearningRlab

Value

A numeric, the geometric mean of the numbers vector.

Note

A vector is created by `c()`, like `c(1,2,3,4,5)` creates a vector with the numbers: 1,2,3,4,5

Author(s)

Dennis Monheimius, <dennis.monhemimius@edu.uah.es>
Eduardo Benito, <eduardo.benito@edu.uah.es>
Juan Jose Cuadrado, <jjcg@uah.es>
Universidad de Alcala de Henares

Examples

```
#data creation  
data = c(1:20)  
  
geometricMean_(data)
```

getUserAction	<i>Get User Action Funcion</i>
---------------	--------------------------------

Description

This function get the buffer introduced by the user. Typically a numerical vector.

Usage

```
getUserAction()
```

Value

A vector

Note

The process is interactive with the user

Author(s)

Dennis Monheimius, <dennis.monhemimius@edu.uah.es>
Eduardo Benito, <eduardo.benito@edu.uah.es>
Juan Jose Cuadrado, <jjcg@uah.es>
Universidad de Alcala de Henares

Examples

```
{  
  ## Not run:  
  vector <- getUserAction()  
  
  ## End(Not run)  
}
```

harmonicMean_	<i>Harmonic Mean Function</i>
---------------	-------------------------------

Description

This function calculates the harmonic mean of a numbers vector.

Usage

```
harmonicMean_ (x)
```

Arguments

x Should be a numbers vector

Details

To calculate the harmonic mean, the user should give a numbers vector. The result is calculated by dividing the number of observations by the reciprocal of each number in the vector. The harmonic mean formule is the following:

$$\text{Harmonic Mean} = \frac{N}{\frac{1}{X_1} + \frac{1}{X_2} + \dots + \frac{1}{X_N}}$$

LearningRlab ®

Value

Numeric, the harmonic mean of the numbers vector.

Note

A vector is created by `c()`, like `c(1,2,3,4,5)` creates a vector with the numbers: 1,2,3,4,5

Author(s)

Jose Manuel Gomez Caceres, <josemanuel.gomezc@edu.uah.es>
 Juan Jose Cuadrado, <jjcg@uah.es>
 Universidad de Alcala de Henares

Examples

```
#data creation
data = c(1,4,3,3,2,5,7,12,1,2,3,12)

harmonicMean_(data)
```

initImages

Init Images Function

Description

This function is used to display an image.

Usage

```
initImages(path)
```

Arguments

path	An url of an image
------	--------------------

Value

There isn't return value

Note

The path should be toward an image

Author(s)

Dennis Monheimius, <dennis.monhemimius@edu.uah.es>
Eduardo Benito, <eduardo.benito@edu.uah.es>
Juan Jose Cuadrado, <jjcg@uah.es>
Universidad de Alcala de Henares

Examples

```
{  
  
  ## Not run:  
  path = "https://i.imgur.com/8237YhzJ.png"  
  initImages(path)  
  
  ## End(Not run)  
}
```

interactive.absolute_acum_frecuency

User Interactive Absolute Accumulated Frecuency Calculus

Description

Interactive function for absolute accumulated frecuency calculus.

Usage

```
interactive.absolute_acum_frecuency()
```

Details

The user provides the dataset when the function needs it. After that, the function will ask what is the correct result for this dataset. The function itself will provide the formula.

Value

An interactive process to calculate the absolute accumulated frequency

Author(s)

Dennis Monheimius, <dennis.monhemimius@edu.uah.es>
 Eduardo Benito, <eduardo.benito@edu.uah.es>
 Juan Jose Cuadrado, <jjcg@uah.es>
 Universidad de Alcala de Henares

Examples

```
## Not run:
interactive.absolute_acum_frecuency()

## End(Not run)
```

```
interactive.absolute_frecuency
```

User Interactive Absolute Frecuency Calculus

Description

Interactive function for absolute frequency calculus.

Usage

```
interactive.absolute_frecuency()
```

Details

The user provides the dataset when the function needs it. After that, the function will ask what is the correct result for this dataset. The function itself will provide the formula.

Value

An interactive process to calculate the absolute frequency

Author(s)

Dennis Monheimius, <dennis.monhemimius@edu.uah.es>
 Eduardo Benito, <eduardo.benito@edu.uah.es>
 Juan Jose Cuadrado, <jjcg@uah.es>
 Universidad de Alcala de Henares

Examples

```
## Not run:  
interactive.absolute_frecuency()  
  
## End(Not run)
```

```
interactive.averageDeviation
```

User Interactive Average Absolute Deviation Calculus

Description

Interactive function for average absolute deviation calculus.

Usage

```
interactive.averageDeviation()
```

Details

The user provides the dataset when the function needs it. After that, the function will ask what is the correct result for this dataset. The function itself will provide the average absolute deviation formule, apart from the averageDeviation_ help document.

Value

An interactive process to calculate the average absolute deviation

Author(s)

Dennis Monheimius, <dennis.monhemimius@edu.uah.es>
Eduardo Benito, <eduardo.benito@edu.uah.es>

References

https://en.wikipedia.org/wiki/Average_absolute_deviation

Examples

```
## Not run:  
interactive.averageDeviation()  
  
## End(Not run)
```

`interactive.binomial` *User Interactive Binomial Distribution Calculus*

Description

Interactive function for binomial distribution calculus.

Usage

```
interactive.binomial()
```

Details

The user provides the values when the function needs it. After that, the function will ask what is the correct result for this datas. The function itself will provide the binomial distribution formule, apart from the `binomial_` help document.

Value

An interactive process to calculate the binomial distribution.

Author(s)

Jose Manuel Gomez Caceres, <josemanuel.gomezc@edu.uah.es>
Juan Jose Cuadrado, <jjcg@uah.es>
Universidad de Alcala de Henares

Examples

```
## Not run:  
interactive.binomial()  
  
## End(Not run)
```

`interactive.chisquared` *User Interactive Chisquared Distribution Calculus*

Description

Interactive function for chisquared distribution calculus.

Usage

```
interactive.chisquared()
```

Details

The user provides the datasets when the function needs it. After that, the function will ask what is the correct result for this dataset. The function itself will provide the chisquared distribution formule, apart from the chisquared_ help document.

Value

An interactive process to calculate the chisquared distribution

Author(s)

Jose Manuel Gomez Caceres, <josemanuel.gomezc@edu.uah.es>
Juan Jose Cuadrado, <jjcg@uah.es>
Universidad de Alcala de Henares

Examples

```
## Not run:  
interactive.chisquared()  
  
## End(Not run)
```

interactive.covariance

User Interactive Covariance Calculus

Description

Interactive function for covariance calculus.

Usage

```
interactive.covariance()
```

Details

The user provides the datasets when the function needs it. After that, the function will ask what is the correct result for this dataset. The function itself will provide the covariance formule, apart from the covariance_ help document.

Value

An interactive process to calculate the covariance

Author(s)

Jose Manuel Gomez Caceres, <josemanuel.gomezc@edu.uah.es>
Juan Jose Cuadrado, <jjcg@uah.es>
Universidad de Alcala de Henares

Examples

```
## Not run:  
interactive.covariance()  
  
## End(Not run)
```

interactive.cv

User Interactive Coefficient of Variation Calculus

Description

Interactive function for Coefficient of Variation calculus.

Usage

```
interactive.cv()
```

Details

The user provides the dataset when the function needs it. After that, the function will ask what is the correct result for this dataset. The function itself will provide the coefficient of variation formule, apart from the cv_ help document.

Value

An interactive process to calculate the average absolute deviation

Author(s)

Jose Manuel Gomez Caceres, <josemanuel.gomezc@edu.uah.es>
Juan Jose Cuadrado, <jjcg@uah.es>
Universidad de Alcala de Henares

Examples

```
## Not run:  
interactive.cv()  
  
## End(Not run)
```

interactive.fisher	<i>User Interactive F Fisher Distribution Calculus</i>
--------------------	--

Description

Interactive function for fisher distribution calculus.

Usage

```
interactive.fisher()
```

Details

The user provides the datasets when the function needs it. After that, the function will ask what is the correct result for this datasets. The function itself will provide the fisher distribution formule, apart from the fisher_ help document.

Value

An interactive process to calculate the fisher distribution

Author(s)

Jose Manuel Gomez Caceres, <josemanuel.gomezc@edu.uah.es>
Juan Jose Cuadrado, <jjcg@uah.es>
Universidad de Alcala de Henares

Examples

```
## Not run:  
interactive.fisher()  
  
## End(Not run)
```

interactive.geometricMean	<i>User Interactive Geometric Mean Calculus</i>
---------------------------	---

Description

Interactive function for geometric mean calculus.

Usage

```
interactive.geometricMean()
```

Details

The user provides the dataset when the function needs it. After that, the function will ask what is the correct result for this dataset. The function itself will provide the geometric mean formule, apart from the geometricMean_ help document.

Value

An interactive process to calculate the geometric mean.

Author(s)

Dennis Monheimius, <dennis.monhemimius@edu.uah.es>
Eduardo Benito, <eduardo.benito@edu.uah.es>
Juan Jose Cuadrado, <jjcg@uah.es>
Universidad de Alcala de Henares

Examples

```
## Not run:  
interactive.geometricMean()  
  
## End(Not run)
```

interactive.harmonicMean

User Interactive Harmonic Mean Calculus

Description

Interactive function for harmonic mean calculus.

Usage

```
interactive.harmonicMean()
```

Details

The user provides the dataset when the function needs it. After that, the function will ask what is the correct result for this dataset. The function itself will provide the harmonic mean formule, apart from the harmonicMean_ help document.

Value

An interactive process to calculate the harmonic mean

Author(s)

Jose Manuel Gomez Caceres, <josemanuel.gomezc@edu.uah.es>
Juan Jose Cuadrado, <jjcg@uah.es>
Universidad de Alcala de Henares

Examples

```
## Not run:  
interactive.harmonicMean()  
  
## End(Not run)
```

interactive.laplace *User Interactive Laplace's Rule Calculus*

Description

Interactive function for Laplace's rule calculus.

Usage

```
interactive.laplace()
```

Details

The user provides the values when the function needs it. After that, the function will ask what is the correct result for this datas. The function itself will provide the Laplace's rule formule, apart from the laplace_ help document.

Value

An interactive process to calculate the Laplace's rule.

Author(s)

Jose Manuel Gomez Caceres, <josemanuel.gomezc@edu.uah.es>
Juan Jose Cuadrado, <jjcg@uah.es>
Universidad de Alcala de Henares

Examples

```
## Not run:  
interactive.laplace()  
  
## End(Not run)
```

interactive.mean	<i>User Interactive Mean Calculus</i>
------------------	---------------------------------------

Description

Interactive function for arithmetic mean calculus.

Usage

```
interactive.mean()
```

Details

The user provides the dataset when the function needs it. After that, the function will ask what is the correct result for this dataset. The function itself will provide the arithmetic mean formule, apart from the mean_ help document.

Value

An interactive process to calculate the arithmetic mean.

Author(s)

Dennis Monheimius, <dennis.monhemimius@edu.uah.es>
Eduardo Benito, <eduardo.benito@edu.uah.es>
Juan Jose Cuadrado, <jjcg@uah.es>
Universidad de Alcala de Henares

Examples

```
{  
  
  ## Not run:  
  interactive.mean()  
  
  ## End(Not run)  
}
```

interactive.median	<i>User Interactive Median Calculus</i>
--------------------	---

Description

Interactive function for median calculus.

Usage

```
interactive.median()
```

Details

The user provides the dataset when the function needs it. After that, the function will ask what is the correct result for this dataset. The function itself will provide the median formule, apart from the median_ help document.

Value

An interactive process to calculate the median

Author(s)

Dennis Monheimius, <dennis.monhemimius@edu.uah.es>
Eduardo Benito, <eduardo.benito@edu.uah.es>
Juan Jose Cuadrado, <jjcg@uah.es>
Universidad de Alcala de Henares

Examples

```
## Not run:  
interactive.median()  
  
## End(Not run)
```

interactive.mode

User Interactive Mode Calculus

Description

Interactive function for mode calculus.

Usage

```
interactive.mode()
```

Details

The user provides the dataset when the function needs it. After that, the function will ask what is the correct result for this dataset.

Value

An interactive process to calculate the mode.

Author(s)

Dennis Monheimius, <dennis.monhemimius@edu.uah.es>
Eduardo Benito, <eduardo.benito@edu.uah.es>
Juan Jose Cuadrado, <jjcg@uah.es>
Universidad de Alcala de Henares

Examples

```
## Not run:  
interactive.mode()  
  
## End(Not run)
```

interactive.normal	<i>User Interactive Normal Distribution Calculus</i>
--------------------	--

Description

Interactive function for normal distribution calculus.

Usage

```
interactive.normal()
```

Details

The user provides the values when the function needs it. After that, the function will ask what is the correct result for this data. The function itself will provide the normal distribution formule, apart from the normal_ help document.

Value

An interactive process to calculate the normal distribution.

Author(s)

Jose Manuel Gomez Caceres, <josemanuel.gomezc@edu.uah.es>
Juan Jose Cuadrado, <jjcg@uah.es>
Universidad de Alcala de Henares

Examples

```
## Not run:  
interactive.normal()  
  
## End(Not run)
```

interactive.pearson	<i>User Interactive Pearson Correlation Calculus</i>
---------------------	--

Description

Interactive function for pearson correlation calculus.

Usage

```
interactive.pearson()
```

Details

The user provides the datasets when the function needs it. After that, the function will ask what is the correct result for this dataset. The function itself will provide the pearson correlation formule, apart from the pearson_ help document.

Value

An interactive process to calculate the pearson correlation.

Author(s)

Jose Manuel Gomez Caceres, <josemanuel.gomezc@edu.uah.es>
Juan Jose Cuadrado, <jjcg@uah.es>
Universidad de Alcala de Henares

Examples

```
## Not run:  
interactive.pearson()  
  
## End(Not run)
```

interactive.percentile	<i>User Interactive Percentile Calculus</i>
------------------------	---

Description

Interactive function for percentiles calculus.

Usage

```
interactive.percentile()
```

Details

The user provides the dataset when the function needs it. After that, the function will ask what is the correct result for this dataset. The function itself will provide the formula.

Value

An interactive process to calculate the percentiles

Author(s)

Dennis Monheimius, <dennis.monhemimius@edu.uah.es>
Eduardo Benito, <eduardo.benito@edu.uah.es>
Juan Jose Cuadrado, <jjcg@uah.es>
Universidad de Alcala de Henares

Examples

```
## Not run:  
interactive.percentile()  
  
## End(Not run)
```

interactive.poisson	<i>User Interactive Poisson Distribution Calculus</i>
---------------------	---

Description

Interactive function for Poisson distribution calculus.

Usage

```
interactive.poisson()
```

Details

The user provides the values when the function needs it. After that, the function will ask what is the correct result for this datas. The function itself will provide the Poisson distribution formule, apart from the poisson_ help document.

Value

An interactive process to calculate the Poisson distribution.

Author(s)

Jose Manuel Gomez Caceres, <josemanuel.gomezc@edu.uah.es>
Juan Jose Cuadrado, <jjcg@uah.es>
Universidad de Alcala de Henares

Examples

```
## Not run:  
interactive.poisson()  
  
## End(Not run)
```

interactive.quartile *User Interactive Quartiles Calculus*

Description

Interactive function for quartiles calculus.

Usage

```
interactive.quartile()
```

Details

The user provides the dataset when the function needs it. After that, the function will ask what is the correct result for this dataset. The function itself will provide the formula.

Value

An interactive process to calculate the quartiles

Author(s)

Dennis Monheimius, <dennis.monhemimius@edu.uah.es>
Eduardo Benito, <eduardo.benito@edu.uah.es>
Juan Jose Cuadrado, <jjcg@uah.es>
Universidad de Alcala de Henares

Examples

```
## Not run:  
interactive.quartile()  
  
## End(Not run)
```

interactive.relative_acum_frecuency

User Interactive Relative Accumulated Frecuency Calculus

Description

Interactive function for relative accumulated frecuency calculus.

Usage

```
interactive.relative_acum_frecuency()
```

Details

The user provides the dataset when the function needs it. After that, the function will ask what is the correct result for this dataset. The function itself will provide the formula.

Value

An interactive process to calculate the relative accumulated frecuency

Author(s)

Dennis Monheimius, <dennis.monhemimius@edu.uah.es>

Eduardo Benito, <eduardo.benito@edu.uah.es>

Juan Jose Cuadrado, <jjcg@uah.es>

Universidad de Alcala de Henares

Examples

```
## Not run:  
interactive.relative_acum_frecuency()  
  
## End(Not run)
```

interactive.relative_frecuency

User Interactive Relative Frecuency Calculus

Description

Interactive function for relative frecuency calculus.

Usage

```
interactive.relative_frecuency()
```

Details

The user provides the dataset when the function needs it. After that, the function will ask what is the correct result for this dataset. The function itself will provide the formula.

Value

An interactive process to calculate the relative frequency

Author(s)

Dennis Monheimius, <dennis.monhemimius@edu.uah.es>
Eduardo Benito, <eduardo.benito@edu.uah.es>
Juan Jose Cuadrado, <jjcg@uah.es>
Universidad de Alcala de Henares

Examples

```
## Not run:  
interactive.relative_frequency()  
  
## End(Not run)
```

interactive.standardDeviation

User Interactive Standard Deviation Calculus

Description

Interactive function for standard deviation calculus.

Usage

```
interactive.standardDeviation()
```

Details

The user provides the dataset when the function needs it. After that, the function will ask what is the correct result for this dataset. The function itself will provide the standard deviation formule, apart from the standardDeviation_ help document.

Value

An interactive process to calculate the standard deviation

Author(s)

Dennis Monheimius, <dennis.monhemimius@edu.uah.es>
Eduardo Benito, <eduardo.benito@edu.uah.es>

References

https://en.wikipedia.org/wiki/Standard_deviation

Examples

```
## Not run:
interactive.standardDeviation()

## End(Not run)
```

interactive.tstudent *User Interactive T-Student Distribution Calculus*

Description

Interactive function for T-Student distribution calculus.

Usage

```
interactive.tstudent()
```

Details

The user provides the values when the function needs it. After that, the function will ask what is the correct result for this datas. The function itself will provide the T-Students distribution formule, apart from the tstudent_ help document.

Value

An interactive process to calculate the T-Student distribution.

Author(s)

Jose Manuel Gomez Caceres, <josemanuel.gomezc@edu.uah.es>
Juan Jose Cuadrado, <jjcg@uah.es>
Universidad de Alcala de Henares

Examples

```
## Not run:
interactive.tstudent()

## End(Not run)
```

interactive.variance	<i>User Interactive Variance Calculus</i>
----------------------	---

Description

Interactive function for variance calculus.

Usage

```
interactive.variance()
```

Details

The user provides the dataset when the function needs it. After that, the function will ask what is the correct result for this dataset. The function itself will provide the variance formule, apart from the variance_ help document.

Value

An interactive process to calculate the average absolute deviation

Author(s)

Dennis Monheimius, <dennis.monhemimius@edu.uah.es>
Eduardo Benito, <eduardo.benito@edu.uah.es>
Juan Jose Cuadrado, <jjcg@uah.es>
Universidad de Alcala de Henares

Examples

```
## Not run:  
interactive.variance()  
  
## End(Not run)
```

laplace_	<i>Laplace's Rule Calculus Function</i>
----------	---

Description

This function calculates the Laplace's rule of experiment.

Usage

```
laplace_(x,y)
```

Arguments

x	Should be a vector
y	Should be a vector

Details

To calculate the Laplace's rule, the user should give two vector (unfavorable cases/favorable cases). The result is as the quotient between the number of favorable cases to A, and that of all possible results of the experiment. The Laplace's rule formule is the following:



The graphic shows the Laplace Rule formula: **LaPlace Rule** equals the fraction of *casos favorables* over *casos posibles*. Below the formula is the LearningRlab logo.

$$\text{LaPlace Rule} = \frac{\text{casos favorables}}{\text{casos posibles}}$$

LearningRlab ®

Value

Numeric, the pearson correlation.

Note

A vector is created by c(), like c(1,2,3,4,5) creates a vector with the numbers: 1,2,3,4,5

Author(s)

Jose Manuel Gomez Caceres, <josemanuel.gomezc@edu.uah.es>
Juan Jose Cuadrado, <jjcg@uah.es>
Universidad de Alcala de Henares

Examples

```
#data creation
data = 3
data2 = c(1,2,3,4,5,6)
laplace_(data, data2)
```

LearningRlab

Statistical Learning Functions

Description

Package used to teach basic statistics to students.

Details

This package pretends to serve the user as a method of learning basic statistical functions at secondary and baccalaureate courses. The content of the package incorporate a serie of statistical functions like the calculus of the arithmetic mean or the calculus of the frequencies. There is no only calculus functions, further more, there are incorporated interactive and explicative functions to help and guide the user in the learning process.

Author(s)

Jose Manuel Gomez Caceres, <josemanuel.gomezc@edu.uah.es>
Dennis Monheimius, <dennis.monhemimius@edu.uah.es>
Eduardo Benito, <eduardo.benito@edu.uah.es>
Juan Jose Cuadrado, <jjcg@uah.es>
Universidad de Alcala de Henares
Maintainer: Eduardo Benito <eduardo.benito@edu.uah.com>

meanC

Mean Function Developed in C

Description

This function calculates the arithmetic mean of a numbers vector.

Usage

meanC(x)

Arguments

x Should be a numbers vector

Details

To calculate the arithmetic mean of a dataset, the user should give a numbers vector. The result is the addition of all vector elements divided by the number of elements. The arithmetic mean formule is the following:

$$\bar{X} = \frac{\sum_{i=1}^n X_i}{n} = \frac{X_1 + X_2 + \dots + X_n}{n}$$

@LearningRlab

Value

A numeric, the arithmetic mean of the numbers vector.

Note

A vector is created by `c()`, like `c(1,2,3,4,5)` creates a vector with the numbers: 1,2,3,4,5

Author(s)

Dennis Monheimius, <dennis.monhemimius@edu.uah.es>
 Eduardo Benito, <eduardo.benito@edu.uah.es>
 Juan Jose Cuadrado, <jjcg@uah.es>
 Universidad de Alcala de Henares

Examples

```
#data creation
vector = c(1:10)
meanC(vector)
```

mean_

Mean Calculus Function

Description

This function calculates the arithmetic mean of a numbers vector.

Usage

```
mean_(x)
```


Arguments

x Should be a numbers vector

Details

To calculate the arithmetic mean of a dataset, the user should give a numbers vector. The result is the addition of all vector elements divided by the number of elements. The arithmetic mean formule is the following:

$$\bar{X} = \frac{\sum_{i=1}^n X_i}{n} = \frac{X_1 + X_2 + \dots + X_n}{n}$$

@LearningRlab

Value

A numeric, the arithmetic mean of the numbers vector.

Note

A vector is created by c(), like c(1,2,3,4,5) creates a vector with the numbers: 1,2,3,4,5

Author(s)

Dennis Monheimius, <dennis.monhemimius@edu.uah.es>
 Eduardo Benito, <eduardo.benito@edu.uah.es>
 Juan Jose Cuadrado, <jjcg@uah.es>
 Universidad de Alcala de Henares

Examples

```
vector <- c(2,4,6,8,10,12,14,16,18)
result = mean_(vector)
result
```

 median_

Median Calculus Function

Description

This function calculates the median of a numbers vector.

Usage

```
median_(x)
```

Arguments

x Should be a numbers vector

Details

To calculate the median, the user should give a numbers vector. The result is the value separating the higher half from the lower half of the dataset, it may be thought of as the middle value. The median formule is the following:

$$\text{Median} = \frac{1}{2}(n + 1)\text{th value}$$

® LearningRlab

Value

A numeric, the median of the numbers vector.

Note

A vector is created by `c()`, like `c(1,2,3,4,5)` creates a vector with the numbers: 1,2,3,4,5

Author(s)

Dennis Monheimius, <dennis.monhemimius@edu.uah.es>
Eduardo Benito, <eduardo.benito@edu.uah.es>
Juan Jose Cuadrado, <jjcg@uah.es>
Universidad de Alcala de Henares

Examples

```
{  
  
result = median_(c(1,3,2,5,12,4,4,2,9))  
result  
}
```

mode_

*Mode Calculus Function***Description**

This function calculates the mode of a numbers vector.

Usage

```
mode_(x)
```

Arguments

x Should be a numbers vector

Details

To calculate the mode of a dataset, the user should give a numbers vector. The result is the numeric value that appears most often. In other words, it's the value that is most likely to be sampled. The mode formule is the following:

MODE FORMULA

$$\text{Mode} = l_1 + \left(\frac{f_0 - f_{-1}}{2f_0 - f_{-1} - f_1} \right) \times c$$

©LearningRlab

Value

Numeric, the mode of the numbers vector.

Note

A vector is created by c(), like c(1,2,3,4,5) creates a vector with the numbers: 1,2,3,4,5

Author(s)

Dennis Monheimius, <dennis.monhemimius@edu.uah.es>
 Eduardo Benito, <eduardo.benito@edu.uah.es>
 Juan Jose Cuadrado, <jjcg@uah.es>
 Universidad de Alcala de Henares

Examples

```
{
  #data creation
  data = c(1,2,2,3,4)

  mode_(data)
```

}

normal_

*Normal Distribution Calculus Function***Description**

This function calculates the normal distribution of experiment.

Usage

```
normal_(x)
```

Arguments

x Should be a numbers.

Details

To calculate the normal distribution, the user should give a number. The result is a type of continuous probability distribution for a real-valued random variable. The normal distribution formule is the following:

$$\text{Normal Distribution} = \frac{1}{\sqrt{2\pi}} e^{-x^2/2} dx$$

LearningRlab ®

Value

Numeric, the normal distribution.

Note

The variable is a number. Example: x <- 0.1

Author(s)

Jose Manuel Gomez Caceres, <josemanuel.gomezc@edu.uah.es>
 Juan Jose Cuadrado, <jjcg@uah.es>
 Universidad de Alcala de Henares

Examples

```
#data creation
x = 0.1

normal_(x)
```

pearson_

*Pearson Correlation Calculus Function***Description**

This function calculates the pearson correlation of two vectors of numbers.

Usage

```
pearson_(x,y)
```

Arguments

x	Should be a vector
y	Should be a vector

Details

To calculate the pearson correlation, the user should give two vectors of numbers. The result is the covariance of the two vectors of numbers divided by the product of their standard deviations. The pearson correlation formule is the following:

$$\text{Pearson Correlation} = \frac{\text{Cov}(x, y)}{S_x \times S_y}$$

LearningRlab®

Value

Numeric, the pearson correlation of two vectors of numbers.

Note

A vector is created by c(), like c(1,2,3,4,5) creates a vector with the numbers: 1,2,3,4,5

Author(s)

Jose Manuel Gomez Caceres, <josemanuel.gomezc@edu.uah.es>
 Juan Jose Cuadrado, <jjcg@uah.es>
 Universidad de Alcala de Henares

Examples

```
#data creation
data = c(1,4,3,3,2,5,7,12,1,2,3,12)
data2 = c(1,2,4,4,6,5,11,2,10,5,6,1)
pearson_(data, data2)
```

percentile_

*Percentile Calculus Function***Description**

This function calculate the percentiles of a vector of numbers

Usage

```
percentile_(x, p)
```

Arguments

x Should be a vector
 p Should be a number, $0 \Rightarrow y \leq 1$

Details

To calculate the percentiles, the user should give a vector. This function divide the dataset in 100 parts as equal as possible. The formula is the following:

$$P_i = \begin{cases} \text{elemento}(E + 1) & \text{para } D \neq 0 \\ \frac{\text{elemento}(E) + \text{elemento}(E+1)}{2} & \text{para } D = 0 \end{cases}$$

LearningRlab ®**Value**

A vector sorted with the elements divided by 100 parts

Note

A vector is created by c(), like c(1,2,3,4,5) creates a vector with the numbers: 1,2,3,4,5

Author(s)

Dennis Monheimius, <dennis.monhemimius@edu.uah.es>
Eduardo Benito, <eduardo.benito@edu.uah.es>
Juan Jose Cuadrado, <jjcg@uah.es>
Universidad de Alcala de Henares

Examples

```
{  
  #data creation  
  data = c(1,4,3,3,2,5,7,12,1,2,3,12)  
  
  percentile_(data,0.3)  
}
```

poisson_

*Poisson Distribution Calculus Function***Description**

This function calculates the Poisson distribution of experiment.

Usage

```
poisson_(k, lam)
```

Arguments

k	Should be a numbers
lam	Should be a numbers

Details

To calculate the Poisson distribution, the user should give two number (the number of times the phenomenon and the number of occurrences). The result is a discrete probability distribution that expresses, from a mean frequency of occurrence, the probability that a certain number of events will occur during a certain period of time. The Poisson distribution formule is the following:

$$\text{Poisson Distribution} = \frac{\lambda^x e^{-\lambda}}{X!}$$

Value

Numeric, the pearson correlation of two numbers.

Note

Each variable is a number. Example: lam <- 2 | k <- 3

Author(s)

Jose Manuel Gomez Caceres, <josemanuel.gomezc@edu.uah.es>
 Juan Jose Cuadrado, <jjcg@uah.es>
 Universidad de Alcala de Henares

Examples

```
#data creation
lam = 2
k = 3
poisson_(k,lam)
```

 quartile_

Quartiles Calculus

Description

Calculates the 3 Quartiles of a vector of data

Usage

```
quartile_(x)
```

Arguments

x Should be a vector

Details

To calculate the quartiles, the user should give a vector. This function divide the dataset in 4 parts as equal as possible. The formula is the following:

$$Q_i = \frac{i(N)}{4} \text{ th value}$$

LearningRlab ®

Value

A vector sorted with the elements divided by 4 parts

Note

A vector is created by `c()`, like `c(1,2,3,4,5)` creates a vector with the numbers: 1,2,3,4,5

Author(s)

Dennis Monheimius, <dennis.monhemimius@edu.uah.es>

Eduardo Benito, <eduardo.benito@edu.uah.es>

Juan Jose Cuadrado, <jjcg@uah.es>

Universidad de Alcala de Henares

Examples

```
{  
  #data creation  
  data = c(1:20)  
  
  quartile_(data)  
}
```

standardDeviation_	<i>Standard Deviation Calculus Function</i>
--------------------	---

Description

This function calculates the standard deviation of a numbers vector.

Usage

```
standardDeviation_(x)
```

Arguments

x	Should be a numbers vector
---	----------------------------

Details

To calculate the standard deviation, the user should give a numbers vector. The result is the square root of the sum of the differences between each vector element and the mean squared divided by

the number of elemets. The standard deviation formule is the following:

$$s = \sqrt{\frac{1}{N-1} \sum_{i=1}^N (x_i - \bar{x})^2}$$

©LearningRlab

Value

Numeric, the standard deviation of the numbers vector.

Note

A vector is created by c(), like c(1,2,3,4,5) creates a vector with the numbers: 1,2,3,4,5

Author(s)

Dennis Monheimius, <dennis.monhemimius@edu.uah.es>
 Eduardo Benito, <eduardo.benito@edu.uah.es>
 Juan Jose Cuadrado, <jjcg@uah.es>
 Universidad de Alcala de Henares

Examples

```
#data creation
data = c(1,4,3,3,2,5,7,12,1,2,3,12)

standardDeviation_(data)
```

tstudent_

T-Student Distribution Calculus Function

Description

This function calculates the T-Student distribution of experiment.

Usage

```
tstudent_(x,u,s,n)
```

Arguments

x	Should be a number
u	Should be a number
s	Should be a number
n	Should be a number

Details

To calculate the T-Student distribution, the user should give four number (sample mean, population mean, population standard deviation and sample size). The result is a probability distribution that arises from the problem of estimating the mean of a normally distributed population when the sample size is small. The T-Student distribution formula is the following:

$$\textbf{T-Student Distribution} = \frac{\bar{x} - \mu}{\sigma / \sqrt{n}}$$

LearningRlab ®

Value

Numeric, the T-Student distribution.

Note

Each variable is a number. Example: `x <- 2 | y <- 4`

Author(s)

Jose Manuel Gomez Caceres, <josemanuel.gomezc@edu.uah.es>

Juan Jose Cuadrado, <jjcg@uah.es>

Universidad de Alcala de Henares

Examples

```
#data creation
x = 52.9
u = 50
s = 3
n = 10

tstudent_(x,u,s,n)
```

variance_

Variance Calculus Function

Description

This function calculates the variance of a numbers vector.

Usage

```
variance_(x)
```

Arguments

x Should be a numbers vector

Details

To calculate the variance, the user should give a numbers vector. The result is the expectation of the squared deviation of all numbers vector from its mean. The variance formule is the following:

$$\text{Variance} = \frac{\sum (X - \bar{X})^2}{N-1}$$

LearningRlab ®

Value

Numeric, the variance of the numbers vector.

Note

A vector is created by c(), like c(1,2,3,4,5) creates a vector with the numbers: 1,2,3,4,5

Author(s)

Dennis Monheimius, <dennis.monhemimius@edu.uah.es>

Eduardo Benito, <eduardo.benito@edu.uah.es>

Juan Jose Cuadrado, <jjcg@uah.es>

Universidad de Alcala de Henares

Examples

```
#data creation
data = c(1,4,3,3,2,5,7,12,1,2,3,12)

variance_(data)
```

Index

- * **~absoluta**
 - averageDeviation_, 3
 - explain.averageDeviation, 12
 - frequency_abs, 35
 - frequency_absolute_acum, 36
 - interactive.averageDeviation, 45
- * **~absolute**
 - averageDeviation_, 3
 - explain.absolute_acum_frequency, 10
 - explain.absolute_frequency, 11
 - explain.averageDeviation, 12
 - frequency_abs, 35
 - frequency_absolute_acum, 36
 - interactive.absolute_acum_frequency, 43
 - interactive.absolute_frequency, 44
 - interactive.averageDeviation, 45
- * **~accumulated**
 - explain.absolute_acum_frequency, 10
 - explain.relative_acum_frequency, 29
 - frequency_absolute_acum, 36
 - frequency_relative_acum, 38
 - interactive.absolute_acum_frequency, 43
 - interactive.relative_acum_frequency, 58
- * **~acumulada**
 - explain.absolute_acum_frequency, 10
 - explain.relative_acum_frequency, 29
 - frequency_absolute_acum, 36
 - frequency_relative_acum, 38
- * **~arithmetic.mean**
 - mean_, 64
 - meanC, 63
- * **~armonica**
 - explain.harmonicMean, 19
 - harmonicMean_, 41
 - interactive.harmonicMean, 50
- * **~average**
 - averageDeviation_, 3
 - explain.averageDeviation, 12
 - interactive.averageDeviation, 45
- * **~binomial**
 - binomial_, 4
 - explain.binomial, 13
 - interactive.binomial, 46
- * **~chiccuadrado**
 - chisquared_, 6
 - explain.chisquared, 14
 - interactive.chisquared, 46
- * **~chisquared_**
 - chisquared_, 6
 - explain.chisquared, 14
- * **~chisquared**
 - interactive.chisquared, 46
- * **~coefficient**
 - cv_, 8
 - explain.cv, 16
 - interactive.cv, 48
- * **~coeficiente**
 - cv_, 8
 - explain.cv, 16
 - interactive.cv, 48
- * **~coger**
 - getUserAction, 41
- * **~correlaccion**
 - explain.pearson, 25
 - pearson_, 69
- * **~correlacion**
 - explain.pearson, 25
 - pearson_, 69
- * **~covaiance**
 - explain.covariance, 15

- * **~covariance**
 - covariance_, 7
 - interactive.covariance, 47
 - interactive.pearson, 55
- * **~covarianza**
 - covariance_, 7
 - explain.covariance, 15
 - interactive.covariance, 47
 - interactive.pearson, 55
- * **~cuartiles**
 - interactive.quartile, 57
 - quartile_, 72
- * **~cv**
 - cv_, 8
 - explain.cv, 16
 - interactive.cv, 48
- * **~desviacion**
 - averageDeviation_, 3
 - explain.averageDeviation, 12
 - explain.standardDeviation, 31
 - interactive.averageDeviation, 45
 - interactive.standardDeviation, 59
 - standardDeviation_, 73
- * **~deviation**
 - averageDeviation_, 3
 - explain.averageDeviation, 12
 - explain.standardDeviation, 31
 - interactive.averageDeviation, 45
 - interactive.standardDeviation, 59
 - standardDeviation_, 73
- * **~distribucion**
 - binomial_, 4
 - explain.binomial, 13
 - explain.fisher, 17
 - explain.normal, 24
 - explain.poisson, 27
 - explain.tstudent, 32
 - fisher_, 34
 - interactive.binomial, 46
 - interactive.fisher, 49
 - interactive.normal, 54
 - interactive.poisson, 56
 - interactive.tstudent, 60
 - normal_, 68
 - poisson_, 71
 - tstudent_, 74
- * **~distribution**
 - binomial_, 4
 - explain.binomial, 13
 - explain.fisher, 17
 - explain.normal, 24
 - explain.poisson, 27
 - explain.tstudent, 32
 - fisher_, 34
 - interactive.binomial, 46
 - interactive.fisher, 49
 - interactive.normal, 54
 - interactive.poisson, 56
 - interactive.tstudent, 60
 - normal_, 68
 - poisson_, 71
 - tstudent_, 74
- * **~draw**
 - drawVector, 9
- * **~explained**
 - explain.averageDeviation, 12
 - explain.standardDeviation, 31
- * **~explain**
 - explain.absolute_acum_frecuency, 10
 - explain.absolute_frecuency, 11
 - explain.binomial, 13
 - explain.chisquared, 14
 - explain.covariance, 15
 - explain.cv, 16
 - explain.fisher, 17
 - explain.geometricMean, 18
 - explain.harmonicMean, 19
 - explain.laplace, 20
 - explain.mean, 21
 - explain.median, 22
 - explain.mode, 23
 - explain.normal, 24
 - explain.pearson, 25
 - explain.percentile, 26
 - explain.poisson, 27
 - explain.quartile, 28
 - explain.relative_acum_frecuency, 29
 - explain.relative_frecuency, 30
 - explain.tstudent, 32
 - explain.variance, 33
- * **~explicada**
 - explain.averageDeviation, 12
 - explain.binomial, 13
 - explain.chisquared, 14

- explain.covariance, 15
- explain.cv, 16
- explain.fisher, 17
- explain.geometricMean, 18
- explain.harmonicMean, 19
- explain.laplace, 20
- explain.mean, 21
- explain.median, 22
- explain.mode, 23
- explain.normal, 24
- explain.pearson, 25
- explain.poisson, 27
- explain.standardDeviation, 31
- explain.tstudent, 32
- explain.variance, 33
- * **~fisher**
 - explain.fisher, 17
 - interactive.fisher, 49
- * **~frecuencia**
 - frecuency_abs, 35
 - frecuency_absolute_acum, 36
 - frecuency_relative, 37
 - frecuency_relative_acum, 38
 - interactive.absolute_frecuency, 44
 - interactive.relative_frecuency, 58
- * **~frecuency**
 - explain.absolute_acum_frecuency, 10
 - explain.absolute_frecuency, 11
 - explain.relative_acum_frecuency, 29
 - explain.relative_frecuency, 30
 - frecuency_abs, 35
 - frecuency_absolute_acum, 36
 - frecuency_relative, 37
 - frecuency_relative_acum, 38
 - interactive.absolute_acum_frecuency, 43
 - interactive.absolute_frecuency, 44
 - interactive.relative_acum_frecuency, 58
 - interactive.relative_frecuency, 58
- * **~geometrica**
 - explain.geometricMean, 18
 - geometricMean_, 40
 - interactive.geometricMean, 49
- * **~geometric**
 - explain.geometricMean, 18
 - geometricMean_, 40
 - interactive.geometricMean, 49
- * **~get**
 - getUserAction, 41
- * **~harmonic**
 - explain.harmonicMean, 19
 - harmonicMean_, 41
 - interactive.harmonicMean, 50
- * **~imagenes**
 - initImages, 42
- * **~images**
 - initImages, 42
- * **~iniciar**
 - initImages, 42
- * **~init**
 - initImages, 42
- * **~interactiva**
 - interactive.averageDeviation, 45
 - interactive.binomial, 46
 - interactive.chisquared, 46
 - interactive.covariance, 47
 - interactive.cv, 48
 - interactive.fisher, 49
 - interactive.harmonicMean, 50
 - interactive.laplace, 51
 - interactive.median, 52
 - interactive.mode, 53
 - interactive.normal, 54
 - interactive.pearson, 55
 - interactive.poisson, 56
 - interactive.standardDeviation, 59
 - interactive.tstudent, 60
 - interactive.variance, 61
- * **~interactive**
 - interactive.absolute_acum_frecuency, 43
 - interactive.absolute_frecuency, 44
 - interactive.averageDeviation, 45
 - interactive.binomial, 46
 - interactive.chisquared, 46
 - interactive.covariance, 47
 - interactive.cv, 48
 - interactive.fisher, 49
 - interactive.geometricMean, 49
 - interactive.harmonicMean, 50
 - interactive.laplace, 51
 - interactive.mean, 52
 - interactive.median, 52

- interactive.mode, 53
- interactive.normal, 54
- interactive.pearson, 55
- interactive.percentile, 55
- interactive.poisson, 56
- interactive.quartile, 57
- interactive.relative_acum_frecuency, 58
- interactive.relative_frecuency, 58
- interactive.standardDeviation, 59
- interactive.tstudent, 60
- interactive.variance, 61
- * **~interactivo**
 - interactive.geometricMean, 49
 - interactive.mean, 52
- * **~laplace**
 - explain.laplace, 20
 - interactive.laplace, 51
 - laplace_, 61
- * **~meanC**
 - meanC, 63
- * **~mean**
 - explain.geometricMean, 18
 - explain.harmonicMean, 19
 - explain.mean, 21
 - geometricMean_, 40
 - harmonicMean_, 41
 - interactive.geometricMean, 49
 - interactive.harmonicMean, 50
 - interactive.mean, 52
 - mean_, 64
- * **~media.aritmetica**
 - mean_, 64
 - meanC, 63
- * **~media.explicada**
 - explain.mean, 21
- * **~media.interactiva**
 - interactive.mean, 52
- * **~mediaC**
 - meanC, 63
- * **~mediana.explicada**
 - explain.median, 22
- * **~mediana.interactiva**
 - interactive.median, 52
- * **~mediana**
 - median_, 65
- * **~median**
 - explain.median, 22
- interactive.median, 52
- median_, 65
- * **~media**
 - explain.geometricMean, 18
 - explain.harmonicMean, 19
 - geometricMean_, 40
 - harmonicMean_, 41
 - interactive.geometricMean, 49
 - interactive.harmonicMean, 50
 - mean_, 64
- * **~moda.explicada**
 - explain.mode, 23
- * **~moda.interactiva**
 - interactive.mode, 53
- * **~moda**
 - mode_, 67
- * **~mode**
 - explain.mode, 23
 - interactive.mode, 53
 - mode_, 67
- * **~normal**
 - explain.normal, 24
 - interactive.normal, 54
 - normal_, 68
- * **~pearson**
 - explain.pearson, 25
 - pearson_, 69
- * **~percentiles**
 - interactive.percentile, 55
- * **~percentile**
 - explain.percentile, 26
 - interactive.percentile, 55
 - percentile_, 70
- * **~percentil**
 - interactive.percentile, 55
 - percentile_, 70
- * **~pintar**
 - drawVector, 9
- * **~poisson**
 - explain.poisson, 27
 - interactive.poisson, 56
 - poisson_, 71
- * **~quantiles**
 - interactive.quartile, 57
- * **~quantile**
 - quartile_, 72
- * **~quartile**
 - explain.quartile, 28

- interactive.quartile, 57
 - quartile_, 72
- * **~regla**
 - explain.laplace, 20
 - interactive.laplace, 51
 - laplace_, 61
- * **~relativa**
 - frecuency_relative, 37
 - frecuency_relative_acum, 38
- * **~relative**
 - explain.relative_acum_frecuency, 29
 - explain.relative_frecuency, 30
 - frecuency_relative, 37
 - frecuency_relative_acum, 38
 - interactive.relative_acum_frecuency, 58
 - interactive.relative_frecuency, 58
- * **~rule**
 - explain.laplace, 20
 - interactive.laplace, 51
 - laplace_, 61
- * **~standard**
 - explain.standardDeviation, 31
 - interactive.standardDeviation, 59
- * **~standart**
 - standardDeviation_, 73
- * **~tipica**
 - explain.standardDeviation, 31
 - interactive.standardDeviation, 59
 - standardDeviation_, 73
- * **~tstudent**
 - explain.tstudent, 32
 - fisher_, 34
 - interactive.tstudent, 60
 - tstudent_, 74
- * **~user**
 - getUserAction, 41
- * **~usuario**
 - getUserAction, 41
- * **~variacion**
 - cv_, 8
 - explain.cv, 16
 - interactive.cv, 48
- * **~variance**
 - explain.variance, 33
 - interactive.variance, 61
 - variance_, 75
- * **~varianza**
 - explain.variance, 33
 - interactive.variance, 61
 - variance_, 75
- * **~variation**
 - cv_, 8
 - explain.cv, 16
 - interactive.cv, 48
- * **~vector**
 - drawVector, 9
- averageDeviation_, 3
- binomial_, 4
- chisquared_, 6
- covariance_, 7
- cv_, 8
- drawVector, 9
- explain.absolute_acum_frecuency, 10
- explain.absolute_frecuency, 11
- explain.averageDeviation, 12
- explain.binomial, 13
- explain.chisquared, 14
- explain.covariance, 15
- explain.cv, 16
- explain.fisher, 17
- explain.geometricMean, 18
- explain.harmonicMean, 19
- explain.laplace, 20
- explain.mean, 21
- explain.median, 22
- explain.mode, 23
- explain.normal, 24
- explain.pearson, 25
- explain.percentile, 26
- explain.poisson, 27
- explain.quartile, 28
- explain.relative_acum_frecuency, 29
- explain.relative_frecuency, 30
- explain.standardDeviation, 31
- explain.tstudent, 32
- explain.variance, 33
- fisher_, 34
- frecuency_abs, 35
- frecuency_absolute_acum, 36
- frecuency_relative, 37

frequency_relative_acum, [38](#)

geometricMean_, [40](#)
getUserAction, [41](#)

harmonicMean_, [41](#)

initImages, [42](#)
interactive.absolute_acum_frequency,
 [43](#)
interactive.absolute_frequency, [44](#)
interactive.averageDeviation, [45](#)
interactive.binomial, [46](#)
interactive.chisquared, [46](#)
interactive.covariance, [47](#)
interactive.cv, [48](#)
interactive.fisher, [49](#)
interactive.geometricMean, [49](#)
interactive.harmonicMean, [50](#)
interactive.laplace, [51](#)
interactive.mean, [52](#)
interactive.median, [52](#)
interactive.mode, [53](#)
interactive.normal, [54](#)
interactive.pearson, [55](#)
interactive.percentile, [55](#)
interactive.poisson, [56](#)
interactive.quartile, [57](#)
interactive.relative_acum_frequency,
 [58](#)
interactive.relative_frequency, [58](#)
interactive.standardDeviation, [59](#)
interactive.tstudent, [60](#)
interactive.variance, [61](#)

laplace_, [61](#)
LearningRlab, [63](#)
LearningRlab-package (LearningRlab), [63](#)

mean_, [64](#)
meanC, [63](#)
median_, [65](#)
mode_, [67](#)

normal_, [68](#)

pearson_, [69](#)
percentile_, [70](#)
poisson_, [71](#)
quartile_, [72](#)
standardDeviation_, [73](#)
tstudent_, [74](#)
variance_, [75](#)