

Package ‘alphaN’

July 22, 2025

Title Set Alpha Based on Sample Size Using Bayes Factors

Version 0.1.2

Description Sets the alpha level for coefficients in a regression model as a decreasing function of the sample size through the use of Jeffreys' Approximate Bayes factor. You tell alphaN() your sample size, and it tells you to which value you must lower alpha to avoid Lindley's Paradox. For details, see Wulff and Taylor (2024) [<doi:10.1177/14761270231214429>](https://doi.org/10.1177/14761270231214429).

License MIT + file LICENSE

URL <https://github.com/jespernwulff/alphaN>

BugReports <https://github.com/jespernwulff/alphaN/issues>

Suggests knitr, rmarkdown, spelling, testthat (>= 3.0.0)

VignetteBuilder knitr

Config/testthat/edition 3

Encoding UTF-8

Language en-US

RoxygenNote 7.3.2

NeedsCompilation no

Author Jesper Wulff [aut, cre] (ORCID: [<https://orcid.org/0000-0002-7976-0939>](https://orcid.org/0000-0002-7976-0939)),
Luke Taylor [aut]

Maintainer Jesper Wulff [<jwulff@econ.au.dk>](mailto:jwulff@econ.au.dk)

Repository CRAN

Date/Publication 2025-07-13 21:00:02 UTC

Contents

alphaN	2
alphaN_plot	3
JAB	4

JABp	5
JABt	6
JAB_plot	7
Index	8

alphaN	<i>Set the alpha level based on sample size for coefficients in a regression models.</i>
--------	--

Description

Set the alpha level based on sample size for coefficients in a regression models.

Usage

```
alphaN(n, BF = 1, method = "JAB", upper = 1)
```

Arguments

n	Sample size
BF	Bayes factor you would like to match. 1 to avoid Lindley’s Paradox, 3 to achieve moderate evidence and 10 to achieve strong evidence.
method	Used for the choice of 'b'. Currently one of: • "JAB": this choice of b produces Jeffery’s approximate BF (Wagenmakers, 2022) • "min": uses the minimal training sample for the prior (Gu et al., 2018) • "robust": a robust version of "min" that prevents too small b (O’Hagan, 1995) • "balanced": this choice of b balances the type I and type II errors (Gu et al, 2016)
upper	The upper limit for the range of realistic effect sizes. Only relevant when method="balanced". Defaults to 1 such that the range of realistic effect sizes is uniformly distributed between 0 and 1, U(0,1).

Value

Numeric alpha level required to achieve the desired level of evidence.

References

Gu et al. (2016). Error probabilities in default Bayesian hypothesis testing. *Journal of Mathematical Psychology*, 72, 130–143.

Gu et al. (2018). Approximated adjusted fractional Bayes factors: A general method for testing informative hypotheses. *The British Journal of Mathematical and Statistical Psychology*, 71(2).

O'Hagan, A. (1995). Fractional Bayes Factors for Model Comparison. Journal of the Royal Statistical Society. Series B (Methodological), 57(1), 99–138.

Wagenmakers (2002). Approximate Objective Bayes Factors From PValues and Sample Size: The 3pn Rule. psyarxiv.

Wulff & Taylor (2023). How and why alpha should depend on sample size: A Bayesian-frequentist compromise for significance testing. PsyArXiv.

Examples

```
# Plot of alpha level as a function of n
seqN <- seq(50, 1000, 1)
plot(seqN, alphaN(seqN), type = "l")
```

alphaN_plot	<i>Creates a plot of alpha as function of sample size for each of the four prior options</i>
-------------	--

Description

Creates a plot of alpha as function of sample size for each of the four prior options

Usage

```
alphaN_plot(BF = 1, max = 10000)
```

Arguments

BF	Bayes factor you would like to match. 1 to avoid Lindley's Paradox, 3 to achieve moderate evidence and 10 to achieve strong evidence.
max	The maximum number of sample size. Defaults to 10,000.

Value

Prints a plot.

Examples

```
# Plot of alpha level as a function of n for a Bayes factor of 3
alphaN_plot(BF = 3)
```

JAB	<i>Transforms a t-statistic from a glm or lm object into Jeffreys' approximate Bayes factor</i>
-----	---

Description

Transforms a t-statistic from a glm or lm object into Jeffreys' approximate Bayes factor

Usage

```
JAB(glm_obj, covariate, method = "JAB", upper = 1)
```

Arguments

glm_obj	a glm or lm object.
covariate	the name of the covariate that you want a BF for as a string.
method	Used for the choice of 'b'. Currently one of: • "JAB": this choice of b produces Jeffery's approximate BF (Wagenmakers, 2022) • "min": uses the minimal training sample for the prior (Gu et al., 2018) • "robust": a robust version of "min" that prevents too small b (O'Hagan, 1995) • "balanced": this choice of b balances the type I and type II errors (Gu et al, 2016)
upper	The upper limit for the range of realistic effect sizes. Only relevant when method="balanced". Defaults to 1 such that the range of realistic effect sizes is uniformly distributed between 0 and 1, U(0,1).

Value

A numeric value for the BF in favour of H1.

Examples

```
# Simulate data

## Sample size
n <- 200

## Regressors
Z1 <- runif(n, -1, 1)
Z2 <- runif(n, -1, 1)
Z3 <- runif(n, -1, 1)
Z4 <- runif(n, -1, 1)
X <- runif(n, -1, 1)

## Error term
```

```

U <- rnorm(n, 0, 0.5)

## Outcome
Y <- X/sqrt(n) + U

# Run a GLM
LM <- glm(Y ~ X + Z1 + Z2 + Z3 + Z4)

# Compute JAB for "X" based on the regression results
JAB(LM, "X")

# Compute JAB using the minimum prior
JAB(LM, "X", method = "min")

```

JABp	<i>Title</i>
------	--------------

Description

Title

Usage

```
JABp(n, p, z = TRUE, df = NULL, method = "JAB", upper = 1)
```

Arguments

n	Sample size.
p	The p-value.
z	Is the p-value based on a z- or t-statistic? TRUE if z.
df	If z=FALSE, provide the degrees of freedom for the t-statistic.
method	Used for the choice of 'b'. Currently one of: "JAB": this choice of b produces Jeffery's approximate BF (Wagenmakers, 2022) "min": uses the minimal training sample for the prior (Gu et al., 2018) "robust": a robust version of "min" that prevents too small b (O'Hagan, 1995) "balanced": this choice of b balances the type I and type II errors (Gu et al, 2016)
upper	The upper limit for the range of realistic effect sizes. Only relevant when method="balanced". Defaults to 1 such that the range of realistic effect sizes is uniformly distributed between 0 and 1, U(0,1).

Value

A numeric value for the BF in favour of H1.

Examples

```
# Transform a p-value of 0.007038863 from a z-test into JAB
# using a sample size of 200.
JABp(200, 0.007038863)

# Transform a p-value of 0.007038863 from a t-test with 190
# degrees of freedom into JAB using a sample size of 200.
JABp(200, 0.007038863, z=FALSE, df=190)
```

JABt

Transforms a t-statistic into Jeffreys' approximate Bayes factor

Description

Transforms a t-statistic into Jeffreys' approximate Bayes factor

Usage

```
JABt(n, t, method = "JAB", upper = 1)
```

Arguments

n	Sample size.
t	The t-statistic.
method	Used for the choice of 'b'. Currently one of: "JAB": this choice of b produces Jeffery's approximate BF (Wagenmakers, 2022) "min": uses the minimal training sample for the prior (Gu et al., 2018) "robust": a robust version of "min" that prevents too small b (O'Hagan, 1995) "balanced": this choice of b balances the type I and type II errors (Gu et al, 2016)
upper	The upper limit for the range of realistic effect sizes. Only relevant when method="balanced". Defaults to 1 such that the range of realistic effect sizes is uniformly distributed between 0 and 1, U(0,1).

Value

A numeric value for the BF in favour of H1.

Examples

```
# Transform a t-statistic of 2.695 computed based on a sample size of 200 into JAB
JABt(200, 2.695)
```

JAB_plot

Plots JAB as a function of the p-value

Description

Plots JAB as a function of the p-value

Usage

```
JAB_plot(n, BF = 1, method = "JAB")
```

Arguments

- | | |
|--------|---|
| n | Sample size |
| BF | Bayes factor you would like to match. 1 to avoid the Lindley Paradox, 3 to achieve moderate evidence and 10 to achieve strong evidence. |
| method | Used for the choice of 'b'. Currently one of: <ul style="list-style-type: none">• "JAB": this choice of b produces Jeffery's approximate BF• "min": uses the minimal training sample for the prior (Gu et al., '17)• "robust": a robust version of "min" that prevents too small b (O'Hagan, '95)• "balanced": this choice of b balances the type I and type II errors |

Value

Prints a plot.

Examples

```
# Plot JAB as function of the p-value for a sample size of 2000  
JAB_plot(2000)
```

Index

alphaN, [2](#)
alphaN_plot, [3](#)

JAB, [4](#)
JAB_plot, [7](#)
JABp, [5](#)
JABt, [6](#)